2024 SAGES ORAL





The association of permanent versus absorbable fixation on developing chronic post-herniorrhaphy groin pain in patients undergoing laparoscopic inguinal hernia repair

Kimberly P. Woo¹ · Ryan C. Ellis¹ · Sara M. Maskal¹ · Daphne Remulla¹ · Priya Shukla¹ · Alexandra J. Rosen¹ · Isabella Wetzka¹ · Wilhemina Osei-Koomson¹ · Sharon Phillips² · Benjamin T. Miller¹ · Lucas R. Beffa¹ · Clayton C. Petro¹ · David M. Krpata¹ · Ajita S. Prabhu¹ · Emanuele Lo Menzo³ · Michael J. Rosen¹

Received: 19 January 2024 / Accepted: 14 April 2024 / Published online: 6 May 2024 © The Author(s) 2024

Abstract

Introduction Fixation of mesh during minimally invasive inguinal hernia repair is thought to contribute to chronic postherniorrhaphy groin pain (CGP). In contrast to permanent tacks, absorbable tacks are hypothesized to minimize the likelihood of CGP. This study aimed to compare the rates of CGP after laparoscopic inguinal hernia repair between absorbable versus permanent fixation at maximum follow-up.

Methods This is a post hoc analysis of a randomized controlled trial in patients undergoing laparoscopic inguinal hernia repair (NCT03835351). All patients were contacted at maximum follow-up after surgery to administer EuraHS quality of life (QoL) surveys. The pain and restriction of activity subdomains of the survey were utilized. The primary outcome was rate of CGP, as defined by a EuraHS QoL pain domain score ≥ 4 measured at ≥ 1 year postoperatively. The secondary outcomes were pain and restriction of activity domain scores and hernia recurrence at maximum follow-up.

Results A total of 338 patients were contacted at a mean follow-up of 28 ± 11 months. 181 patients received permanent tacks and 157 patients received absorbable tacks during their repair. At maximum follow-up, the rates of CGP (27 [15%] vs 28 [18%], P = 0.47), average pain scores (1.78 ± 4.38 vs 2.32 ± 5.40 , P = 0.22), restriction of activity scores (1.39 ± 4.32 vs 2.48 ± 7.45 , P = 0.18), and the number of patients who reported an inguinal bulge (18 [9.9%] vs 15 [9.5%], P = 0.9) were similar between patients with permanent versus absorbable tacks. On multivariable analysis, there was no significant difference in the odds of CGP between the two groups (OR 1.23, 95% CI [0.60, 2.50]).

Conclusion Mesh fixation with permanent tacks does not appear to increase the risk of CGP after laparoscopic inguinal hernia repair when compared to fixation with absorbable tacks. Prospective trials are needed to further evaluate this relationship.

Keywords Inguinal hernia · Tack · Fixation · Groin pain

In the USA, an estimated 800,000 inguinal hernia operations are done annually [1], with up to 1 in 4 men and 1 in 30 women undergoing a repair in their lifetime [2]. With the advent of minimally invasive surgery, the trend has shifted toward laparoscopic approaches [3, 4], such as the totally extraperitoneal (TEP) or trans-abdominal preperitoneal (TAPP) repair techniques, both of which commonly include placement and fixation of mesh. While postoperative complications are rare, post-herniorrhaphy groin pain is a relatively common adverse outcome.

However, the etiology of this pain is not well understood and is likely multifactorial in nature. Age, gender, BMI, preoperative pain level, hernia size, presence of postoperative complications and history of prior inguinal hernia repair have all previously been identified as potential predictors for postoperative pain [5-10]. Additionally, there exists controversy surrounding the impact of fixation on

Kimberly P. Woo Wook2@ccf.org

¹ Department of General Surgery, Digestive Disease Institute, Cleveland Clinic, 9500 Euclid Ave, Cleveland, OH 44195, USA

² Department of Biostatistics, Vanderbilt University Medical Center, Nashville, TN, USA

³ Department of General Surgery, Bariatric and Metabolic Institute, Cleveland Clinic Florida, Weston Hospital, Weston, FL, USA

post-herniorrhaphy groin pain. Permanent titanium tacks have historically been used, but a number of reported complications [11, 12] and their possible association with severe postoperative pain [13, 14] has generated concern regarding the type of tacks that should be used for fixation. Though few studies have definitively identified a significant effect of permanent fixation on pain [15], the theoretical concern has led to the development of more costly alternatives, such as absorbable tacks.

The effect of permanent versus absorbable fixation on chronic post-herniorrhaphy groin pain after inguinal hernia repair has been assessed in only a handful of studies with small patient cohorts [16, 17]. Comparatively, it has been evaluated in more studies in incisional or ventral hernias. The data there has not shown an increased risk of chronic post-herniorrhaphy groin pain with permanent tacks [18–22] and therefore, no conclusive data exist to justify the use of the more expensive absorbable tacks for fixation. Still, randomized controlled trials were relatively small and lacked long-term follow-up of pain. We aim in this study to compare the rates of chronic post-herniorrhaphy groin pain after laparoscopic inguinal hernia repair between permanent versus absorbable tack fixation at maximum follow-up of a large cohort of patients.

Methods

Patient population and data source

This study is a long-term follow-up and post hoc analysis of a multi-center, single-blinded, parallel group RCT (NCT03835351). The original trial design and outcomes were previously reported [23]. In brief, patients 18 years or older, undergoing laparoscopic, elective inguinal hernia repair were assessed for the effect of intraoperative urinary catheters on postoperative urinary retention. The trial was conducted at six academic and community hospitals with 11 participating surgeons from March 2019 to March 2021. Institutional Review Board (IRB) approval was granted at participating sites before enrollment and all study participants provided written informed consent. Patients unable to provide written informed consent or unable to tolerate general anesthesia were excluded.

All operations were performed by general surgeons with advanced training in minimally invasive inguinal hernia repair. Surgical technique and method of mesh fixation were left to the discretion of the operating surgeon. Permanent fixation was achieved with ProTackTM (Covidien, Minneapolis, MN) titanium tacks, while absorbable fixation was achieved with AbsorbaTackTM (Covidien, Minneapolis, MN).

IRB approval was obtained for the current post hoc analysis. All patients who completed the previous RCT were eligible for inclusion in this analysis. These patients were contacted at maximum follow-up after surgery to obtain EuraHS Quality of Life (QoL) and Hernia Recurrence Inventory (HRI) assessments. EuraHS is a hernia-specific OoL instrument, validated for the perioperative assessment of inguinal repair outcomes [24]. Patients score their symptoms from a range of 0 to 10 with 10 indicating the most severe effect and interpreted as worse QoL. HRI is a patientreported outcome (PRO) questionnaire that consists of three questions: (1) Do you think your hernia has come back? (2) Do you feel or see a bulge? (3) Do you have pain or physical symptoms at the site. The question "Do you feel or see a bulge?" has previously been demonstrated to be highly sensitive and specific for the diagnosis of an inguinal hernia recurrence [25] and was used in this study to measure the rates of patient-reported hernia recurrence.

Outcome of interest

Two subdomains of the EuraHS-QoL survey were included in this analysis: pain and restriction of activities. The pain domain ranges from 0 to 30 and restriction of activities from 0 to 40. The previously validated minimum clinically important differences (MCID) for each domain are 3 for the pain domain and 5 for the restriction of activity domain [26]. The primary outcome was rate of chronic post-herniorrhaphy groin pain, as defined by a EuraHS-QoL pain domain score >4 measured at maximum follow-up (at least 1 year) after surgery. A cutoff score that is indicative of persistent pain has yet to be validated for this instrument. Therefore, a score of ≥ 4 was identified based on previously employed quantitative definitions in the chronic pain literature [27-30]. Additional outcomes from the EuraHS-QoL survey were average pain and restriction of activity domain scores and HRI patient-reported hernia recurrence at maximum followup. Multivariable analysis was performed adjusting for possible risk factors of chronic post-herniorrhaphy groin pain, including age, BMI, hernia size, and number of tacks. Number of tacks was analyzed as a categorical variable, <4 tacks vs > 4 tacks. Given the standard for mesh fixation is four tacks, this cutoff was chosen to ensure the variable reflected increasing tack type from the norm.

Statistical analysis

Continuous variables were compared using Wilcoxon rank sum test and categorical variables were compared using Pearson Chi-square or Fisher's exact test where appropriate. Multivariate analysis was conducted using logistic regression. A P value < 0.05 was considered statistically significant. All statistical analyses were conducted using R software (version 4.2.2, Vienna, Austria).

Results

Of the 496 patients in the original RCT who were randomized, 486 patients (98%) remained at the conclusion of the trial and were eligible for inclusion in the current study. All eligible patients were contacted, and follow-up was successfully obtained for a total of 345 patients at a mean follow-up of 28 ± 11 months. Operative details, including fixation type, were not available for seven patients. Therefore, they were excluded, and 338 (70%) patients were included in the statistical analysis.

Patient and operative characteristics

Mesh fixation for inguinal hernia repair was achieved with permanent tacks in 181 patients (53.6%) and absorbable tacks in 157 patients (46.4%). The baseline characteristics and comorbidities between these two groups were similar (Table 1). The median (IQR) age of patients who received permanent fixation was 61 (54–67) years and 62 (50–70) years for absorbable fixation. The baseline median (IQR)reported pain scores were also similar (P=0.54); 6 (2–12) for patients with permanent fixation and 8 (3–11) for patients with absorbable fixation. However, baseline EuraHS-QoL surveys were not collected from all patients in the original 3435

RCT and were only available for 229/338 (67%) patients in this analysis.

The perioperative characteristics are listed in Table 2. More of the patients who received permanent fixation had previously undergone hernia repair with mesh than those who received absorbable fixation (12 [15.4%] vs 2 [3.3%], P = 0.023). Patients with permanent fixation tended to have hernias that were > 3 cm or > 2 fingertips compared to those with absorbable fixation (53 [29.8%] vs 26 [16.8%],P = 0.01). For the operative approach, 103 patients (30.9%) underwent TAPP and 230 (69.1%) underwent TEP inguinal hernia repair. Absorbable tacks were used almost exclusively in the TEP approach (153 [98.7%]), whereas permanent tacks were used almost equally between TAPP and TEP repairs (101 [56.7%] and 77 (43.3%)]. More patients undergoing repair with permanent fixation received greater than four tacks for mesh fixation compared to those with absorbable fixation (110 [61.8%] vs 7 [4.5%], P<0.001).

Patient-reported outcomes: EuraHS-QoL and HRI

Average follow-up for the full cohort was 26 months; 27 months for permanent fixation and 26 months for absorbable fixation (P=0.41). Maximum follow-up was collected at 1 year for 62 patients (18.3%), at 2 years for 146 patients (43.2%), at 3 years for 122 patients (36.1%), and at 4 years

	п	Permanent fixation $n = 181$	Absorbable fixation $n = 157$	P value
Age (years), median (IQR)	338	61 (54, 67)	62 (50, 70)	0.69
Sex	338			0.70
Female		6.1% (11)	5.1% (8)	
Male		93.9% (170)	94.9% (149)	
BMI	335			0.09
≤ 29.9		79.3% (142)	82.7% (129)	
30–34.9		14.5% (26)	16.0% (25)	
35–39.9		5.6% (10)	0.64% (1)	
≥ 40		0.6% (1)	0.64% (1)	
ASA class	335			0.37
1		12.3% (22)	12.8% (20)	
2		54.7% (98)	62.8% (98)	
3		32.4% (58)	23.7% (37)	
4		0.6% (1)	0.6% (1)	
Diabetes	335	4.5% (8)	5.8% (9)	0.59
Hypertension	338	38% (68)	29% (46)	0.11
COPD	335	1.7% (3)	0.0% (0)	0.10
Current smoker	335	6.2% (11)	5.8% (9)	0.88
Baseline EuraHS QOL scores				
Pain domain, median (IQR)	229	6 (2, 12)	8 (3, 11)	0.54
Restriction of activity domain, median (IQR)	227	9 (1.5, 18)	11 (2, 20)	0.22

Table 2 Perioperative characteristics

	п	Permanent fixation $n = 181$	Absorbable fixation $n = 156$	P value
Prior inguinal mesh	139			0.02
No		84.6% (66)	96.7% (59)	
Yes		15.4% (12)	3.3% (2)	
Hernia size	333			0.01
\leq 3 cm or \leq 2 fingertips		70.2% (125)	83.2% (129)	
>3 cm or>2 fingertips		29.8% (53)	16.8% (26)	
Laterality	333			0.92
Bilateral		38.2% (68)	38.7% (60)	
Unilateral		61.8% (110)	61.3% (95)	
Surgical repair approach	333			< 0.001
TAPP		56.7% (101)	1.3% (2)	
TEP		43.3% (77)	98.7% (153)	
Number of tacks	333			< 0.001
≤ 4		38.2% (68)	95.5% (147)	
>4		61.8% (110)	4.5% (7)	
Operation time	335			< 0.001
<1 h		28.5% (51)	46.8% (73)	
≥1 h		71.5% (128)	53.2% (83)	

for eight patients (2.4%). At maximum follow-up, the mean EuraHS-QoL pain domain scores were not statistically different between the patients who received permanent versus absorbable fixation $(1.78 \pm 4.38 \text{ vs } 2.32 \pm 5.40, P = 0.22)$. The restriction of activity domain scores were also similar $(1.39 \pm 4.32 \text{ vs } 2.48 \pm 7.45, P = 0.18)$. Tack type did not influence patient-reported hernia recurrence; the number of patients who reported an inguinal bulge was similar between permanent and absorbable fixation (18 [9.9%] vs 15 [9.5%], P = 0.9).

Chronic post-herniorrhaphy groin pain and predictors

The overall rate of chronic post-herniorrhaphy groin pain (pain domain score \geq 4) at maximum follow-up was 16.3% (Fig. 1). There was no statistically significant difference in the rates between permanent and absorbable fixation (27 [15%] vs 28 [18%], P=0.47). On multivariable analysis, age, BMI, hernia size, and number of tacks used were not associated with chronic post-herniorrhaphy groin pain (Table 3). Though patients with absorbable fixation had 1.3 times odds of having chronic pain at maximum follow-up compared to patients with permanent fixation, this did not reach statistical significance (95% CI 0.60, 2.50).



Fig. 1 Pain scores at maximum follow-up

Discussion

Inguinal hernia repair is a very common and relatively safe procedure with minimal complications. However, some patients may experience persistent pain after surgery, known as chronic post-herniorrhaphy groin pain, with the type of tack used for mesh fixation thought to be a contributing factor. In this post hoc analysis of patients undergoing laparoscopic inguinal hernia repair, we did not identify any difference in the pain scores at maximum follow-up

 Table 3
 Predictors of chronic post-herniorrhaphy groin pain at maximum follow-up

	OR (95% CI)	Significance (P)	
BMI	0.96 (0.71–1.30)	0.79	
Age	0.94 (0.76–1.16)	0.59	
Tack type			
Permanent	1.0 (Reference)	0.57	
Absorbable	1.23 (0.60-2.50)		
Hernia size			
\leq 3 cm	1.0 (Reference)	0.49	
> 3 cm	1.27 (0.64-2.50)		
Number of tacks used			
≤4 tacks	1.0 (Reference)	0.71	
>4 tacks	0.87 (0.40-1.88)		

or in the rates of chronic post-herniorrhaphy groin pain between absorbable versus permanent tack fixation.

The rates of chronic post-herniorrhaphy groin pain vary widely in the literature, from 11 to 53% [6-9, 31-35]. This is generally considered to be due to the lack of a uniform definition or standardized tool for assessment. The International Association for the Study of Pain (IASP) defines chronic postsurgical pain as pain that "develops or increases in intensity after a surgical procedure" and "persists beyond the healing process, i.e., at least 3 months after the surgery" [36]. Most studies of chronic pain cite this definition, with collection of data occurring at time points of six months or greater [6, 7, 31-33]. Our rate of chronic post-herniorrhaphy groin pain at 1 year or greater falls within the range of published data, though on the lower end of the spectrum. This can potentially be attributed to how pain was defined and assessed for in our analysis. A closer review of the literature reveals that our rate of 16.3% is consistent with rates of approximately 11-12% that were calculated from pooled analyses [34, 35] and with other questionnaire-based studies that employed quantitative cutoffs for reported pain [32, 33]. This is in contrast to studies that found higher rates but defined chronic pain as simply the presence of pain to some extent [6-8, 31]. By establishing a cutoff, we aimed to identify groin pain that was clinically relevant and more likely to impact patient quality of life.

Given its relatively high incidence, many studies have aimed to identify the risk factors associated with the development of chronic post-herniorrhaphy groin pain. Younger age and recurrent hernias have consistently been found to be independent risk factors for chronic pain [5-9, 31, 34]. Despite our permanent fixation cohort having a significantly greater proportion of patients with prior inguinal mesh, the rates of patients reporting chronic groin pain did not differ compared to the absorbable fixation cohort. Other risk factors, such as smaller hernia defect size [37], increased BMI [10], or increased number of tacks used for fixation [38] have also been implicated in the development of chronic pain. None of these variables were found to be statistically significant in our multivariate analysis. Of note, it has also been demonstrated that high levels of preoperative pain serve as a risk factor for chronic pain [6, 7, 39]. There were, unfortunately, insufficient baseline pain scores available to reliably include this variable in our multivariate analysis.

While no studies on chronic post-herniorrhaphy groin pain have identified tack type as an independent risk factor, there have been only a few studies, limited by small sample sizes, directly comparing the impact of permanent versus absorbable tacks in inguinal hernia repairs. This was evaluated in 2018 by Hany et al. [16] in an RCT of 30 patients undergoing TAPP repair. At six months and at 1 year, there was no statistically significant difference in the pain scores between patients with permanent versus absorbable fixation. At a mean follow-up of 12 months, there were no recurrences in any of their patients. This study was followed in a case-control study of 20 patients undergoing TEP repair by Prakash et al. [17] in 2019. They too found no difference in pain scores at any of the follow-up time points ranging from six hours to six months postoperatively and no recurrences were identified with the mean follow-up of 24 months. Consistent with these two prior studies, we did not find any difference between the two fixation types in the patient-reported pain scores or in the rates of chronic pain as previously defined. No other prospective studies with larger patient cohorts have been published comparing the two types of tacks. To our knowledge, our post hoc analysis is one of the largest studies of any type evaluating the impact of tack type on chronic post-herniorrhaphy groin pain after inguinal hernia repair.

Despite the lack of prospectively collected data in inguinal hernias, there have been several RCTs comparing outcomes of absorbable versus permanent tack fixation in ventral or incisional hernia repairs. In one of the earliest RCTs comparing the two types of tacks for fixation, Colak et al. [18] found no difference in the six-month pain scores of 51 patients. At a median follow-up of 31 months, there was also no difference in rates of recurrence. Two subsequent RCTs similarly concluded that permanent tacks did not increase the risk of chronic post-herniorrhaphy groin pain or rates of recurrence [19, 20]. Interestingly, two retrospective analyses [21, 22] found that while the rates of chronic pain were the same, early postoperative pain was found to be significantly decreased in patients receiving absorbable tack fixation. The authors posited that the lack of deep penetration into bony structures of the absorbable tacks could explain this relative decrease in pain on postoperative days 1 and 10 [21]. Nevertheless, a difference in early postoperative pain was not supported in any of the randomized trials [18–20]. Even with the compelling evidence showing no association between tack type and chronic pain in ventral hernia repairs, the differences in anatomy between the abdominal wall and groin make this data alone insufficient to conclude the same in inguinal hernia repairs.

Importantly, as healthcare spending in the USA continues to rise, there has been a lack of demonstrable improvement in quality of care and outcomes [40]. With inguinal hernia repairs, the trend toward laparoscopic, and now robotic, approaches increases cost without value added [41]. The tendency to employ newer, more expensive technology and devices without evidence of superiority further adds to this burden. While costs can vary, at our institution, permanent fixation with ProTackTM is approximately \$200 less than the cost of absorbable fixation with AbsorbaTackTM. Our finding of similar outcomes between the two types of fixation, consistent with prior published literature, provides little justification to employ the more costly option.

While acknowledging the benefits of the large sample size in this analysis, there are some limitations that deserve mention. Because this was a post hoc analysis, there may have been a selection bias in the decision of which patients were enrolled in the original RCT. Importantly, though, the inclusion criteria of the trial were very broad [23], and the CONSORT diagram indicates that only a small proportion of patients (38/534, 7.1%) undergoing inguinal hernia repair during the enrollment period were excluded from the trial. Perhaps more significantly, follow-up was only successfully obtained in 70% of eligible patients. This leaves the possibility this missing data could have impacted the results of our study. We therefore performed a sensitivity analysis assuming a minimum pain score of 0 for all missing patients and a maximum pain score of 30 for all missing patients. Neither of these analyses yielded significant associations between pain at maximum follow-up and tack type. As previously stated, preoperative pain has been implicated as a significant predictor for the development of postoperative chronic pain. However, incomplete collection of baseline pain scores precluded the ability to reliably include this variable in our multivariate analysis. Our selected definition for chronic pain may not have been powered to detect smaller differences in rates of patients experiencing chronic pain between the two types of mesh fixation. While the HRI has been validated for the detection of recurrent inguinal hernias, it has not been specifically studied in the obese population. The ability to identify a recurrence may have been limited in this group of patients within our study population. And finally, the retrospective nature of our analysis precluded the ability to guarantee data collection at specific time points. Therefore, our outcomes were measured at maximum follow-up, thereby requiring uniform analysis of a potentially heterogeneous population.

In conclusion, the fixation of mesh with permanent tacks does not appear to increase the risk of chronic

post-herniorrhaphy groin pain after laparoscopic inguinal hernia repair when compared to fixation with absorbable tacks. Appropriately powered prospective trials are needed to further evaluate this relationship.

Funding None.

Declarations

Disclosures Dr. Prabhu is on the advisory board for Surgimatix and CMR Surgical and receives consulting fees from CMR Surgical and Verb Surgical. Dr. Miller has received a research grant from the American Hernia Society. Dr. Petro serves as a consultant for Advanced Medical Solutions, Bardt-Davol, and Surgimatix and has received research grants from the American Hernia Society, the Central Surgical Association, and the Society of American Gastrointestinal and Endoscopic Surgeons. Dr. Rosen serves as the medical director of the ACHQC, received a grant to his institution for research from Telabio, and has stock options with Ariste. Dr. Maskal has received a research grant from the ACHQC. None of these disclosures are relevant to this study. All other authors have no conflicts of interest or financial ties to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- Rutkow IM (2003) Demographic and socioeconomic aspects of hernia repair in the United States in 2003. Surg Clin N Am 83(5):1045–1051
- Kingsnorth A, LeBlanc K (2003) Hernias: inguinal and incisional. Lancet 362(9395):1561–1571
- Holleran TJ, Napolitano MA, Sparks AD, Duncan JE, Garrett M, Brody FJ (2022) Trends and outcomes of open, laparoscopic, and robotic inguinal hernia repair in the veterans affairs system. Hernia 26(3):889–899
- Madion M, Goldblatt MI, Gould JC, Higgins RM (2021) Tenyear trends in minimally invasive hernia repair: a NSQIP database review. Surg Endosc 35(12):7200–7208
- Christoffersen MW, Brandt E, Helgstrand F, Westen M, Rosenberg J, Kehlet H, Strandfelt P, Bisgaard T (2015) Recurrence rate after absorbable tack fixation of mesh in laparoscopic incisional hernia repair. Br J Surg 102(5):541–547
- Kalliomäki ML, Meyerson J, Gunnarsson U, Gordh T, Sandblom G (2008) Long-term pain after inguinal hernia repair in a population-based cohort; risk factors and interference with daily activities. Eur J Pain 12(2):214–225
- Fränneby U, Sandblom G, Nordin P, Nyrén O, Gunnarsson U (2006) Risk factors for long-term pain after hernia surgery. Ann Surg 244(2):212–219

- Nienhuijs S, Staal E, Strobbe L, Rosman C, Groenewoud H, Bleichrodt R (2007) Chronic pain after mesh repair of inguinal hernia: a systematic review. Am J Surg 194(3):394–400
- Poobalan AS, Bruce J, Smith WCS, King PM, Krukowski ZH, Chambers WA (2003) A review of chronic pain after inguinal herniorrhaphy. Clin J Pain 19(1):48–54
- Niebuhr H, Wegner F, Hukauf M, Lechner M, Fortelny R, Bittner R, Schug-Pass C, Köckerling F (2018) What are the influencing factors for chronic pain following TAPP inguinal hernia repair: an analysis of 20,004 patients from the Herniamed Registry. Surg Endosc 32(4):1971–1983
- Fitzgerald HL, Orenstein SB, Novitsky YW (2010) Small bowel obstruction owing to displaced spiral tack after laparoscopic TAPP inguinal hernia repair. Surg Laparosc Endosc Percutan Tech 20(3):e132–e135
- Vujovic Z, Cuarana E, Campbell KL, Valentine N, Koch S, Ziyaie D (2015) Lumbosacral discitis following laparoscopic ventral mesh rectopexy: a rare but potentially serious complication. Tech Coloproctol 19(4):263–265
- 13. Eriksen JR, Poornoroozy P, Jørgensen LN, Jacobsen B, Friis-Andersen HU, Rosenberg J (2009) Pain, quality of life and recovery after laparoscopic ventral hernia repair. Hernia 13(1):13–21
- Wong J, Anvari M (2001) Treatment of inguinodynia after laparoscopic herniorrhaphy. Surg Laparosc Endosc Percutan Tech 11(2):148–151
- Shah NS, Fullwood C, Siriwardena AK, Sheen AJ (2014) Mesh fixation at laparoscopic inguinal hernia repair: a meta-analysis comparing tissue glue and tack fixation. World J Surg 38(10):2558–2570
- 16. Hany D, Gamal M (2018) Postoperative pain and recurrence after mesh fixation with absorbable versus non-absorbable tacks in patients undergoing laparoscopic transabdominal inguinal hernia repair: a comparative study. Ain Shams J Surg 19(2):66–74
- Prakash PS, Wijerathne S, Salgaonkar HP, Lomanto D (2019) The efficacy of absorbable versus non-absorbable fixation in laparoscopic totally extraperitoneal (TEP) repair of large inguinal hernias. Asian J Surg 42(12):995–1000
- Colak E, Ozlem N, Kucuk GO, Aktimur R, Kesmer S, Yildirim K (2015) Prospective randomized trial of mesh fixation with absorbable versus nonabsorbable tacker in laparoscopic ventral incisional hernia repair. Int J Clin Exp Med 8(11):21611–21616
- Bansal VK, Asuri K, Panaiyadiyan S, Kumar S, Subramaniam R, Ramachandran R, Sagar R, Misra MC (2016) Comparison of absorbable versus nonabsorbable tackers in terms of long-term outcomes, chronic pain, and quality of life after laparoscopic incisional hernia repair: a randomized study. Surg Laparosc Endosc Percutan Tech 26(6):476–483
- Harsløf S, Krum-Møller P, Sommer T, Zinther N, Wara P, Friis-Andersen H (2018) Effect of fixation devices on postoperative pain after laparoscopic ventral hernia repair: a randomized clinical trial of permanent tacks, absorbable tacks, and synthetic glue. Langenbecks Arch Surg 403(4):529–537
- Sikar HE, Çetin K, Eyvaz K, Gökçeimam M, Kaptanoglu L, Küçük HF (2019) Evaluation of the effects of absorbable and nonabsorbable tacks on laparoscopic suprapubic hernia repair: a retrospective cohort study. Int J Surg 63:16–21
- Cavallaro G, Campanile FC, Rizzello M, Greco F, Iorio O, Iossa A, Silecchia G (2013) Lightweight polypropylene mesh fixation in laparoscopic incisional hernia repair. Minim Invasive Ther Allied Technol 22(5):283–287
- 23. Fafaj A, Lo Menzo E, Alaedeen D, Petro C, Rosenblatt S, Szomstein S, Massier C, Prabhu AS, Krpata DM, Cha W, Montelione K, Tastaldi L, Alkhatib H, Zolin SJ, Okida LF, Rosen MJ (2022) Effect of intraoperative urinary catheter use on postoperative urinary retention after laparoscopic inguinal hernia repair: a randomized clinical trial. JAMA Surg 157(8):667

- 3439
- 24. Muysoms F, Campanelli G, Champault GG, DeBeaux AC, Dietz UA, Jeekel J, Klinge U, Köckerling F, Mandala V, Montgomery A, Morales Conde S, Puppe F, Simmermacher RKJ, Śmietański M, Miserez M (2012) EuraHS: the development of an international online platform for registration and outcome measurement of ventral abdominal wall hernia repair. Hernia 16(3):239–250
- Tastaldi L, Barros PHF, Krpata DM, Prabhu AS, Rosenblatt S, Petro CC, Alkhatib H, Szutan LA, Silva RA, Olson MA, Stewart TG, Roll S, Rosen MJ, Poulose BK (2020) Hernia recurrence inventory: inguinal hernia recurrence can be accurately assessed using patientreported outcomes. Hernia 24(1):127–135
- 26. Shukla P, Huang LC, Zhao A, Sharew B, Miller B, Beffa L, Petro CC, Krpata DM, Prabhu AS, Rosen MJ (2023) Determining the minimum clinically important difference for the European Hernia Society quality of life instrument in inguinal hernia repair patients. J Am Coll Surg 237(3):525
- Jonsson AC (2006) Prevalence and intensity of pain after stroke: a population based study focusing on patients' perspectives. J Neurol Neurosurg Psychiatry 77(5):590–595
- Collins SL, Moore AR, McQuay HJ (1997) The visual analogue pain intensity scale: what is moderate pain in millimetres? Pain 72(1):95–97
- Rago R, Forfori F, Materazzi G, Abramo A, Collareta M, Miccoli P, Giunta F (2012) Evaluation of a preoperative pain score in response to pressure as a marker of postoperative pain and drugs consumption in surgical thyroidectomy. Clin J Pain 28(5):382–386
- 30. PT Surg Portuguese Surgical Research Collaborative, Simões J, João A, Azevedo J, Peyroteo M, Cunha M, Vieira B, Gonçalves N, Costa J, Soares A, Pimenta J, Romano M, Cinza A, Miguel I, Martins A, Fialho G, Reia M, Borges F, Monteiro C, Soares A, Sousa P, Frade S, Matos L, Carvas J, Martins S, Sousa X, Rodrigues C, Carvalho J, Gil I, Castro L, Rombo N, Quintela A, Ribeiro H, Parreira R, Santos P, Caires F, Torre A, Rodrigues S, Guimarães A, Carvalho M, Pimentel M, Santos D, Ramos C, Cunha C, Azevedo C (2020) Portuguese Inguinal Hernia Cohort (PINE) Study. Surgery. https://doi.org/10.1101/2020.12.19.20247585
- Bay-Nielsen M, Perkins FM, Kehlet H (2001) Pain and functional impairment 1 year after inguinal herniorrhaphy: a nationwide questionnaire study. Ann Surg 233(1):1–7
- Lundström KJ, Holmberg H, Montgomery A, Nordin P (2017) Patient-reported rates of chronic pain and recurrence after groin hernia repair. Br J Surg 105(1):106–112
- Reinpold WMJ, Nehls J, Eggert A (2011) Nerve management and chronic pain after open inguinal hernia repair: a prospective two phase study. Ann Surg 254(1):163–168
- Aasvang E, Kehlet H (2005) Chronic postoperative pain: the case of inguinal herniorrhaphy. Br J Anaesth 95(1):69–76
- Nienhuijs SW, Boelens OBA, Strobbe LJA (2005) Pain after anterior mesh hernia repair. J Am Coll Surg 200(6):885–889
- 36. Schug SA, Lavand'homme P, Barke A, Korwisi B, Rief W, Rolf-Detleft T, The IASP Taskforce for the Classification of Chronic Pain (2019) The IASP classification of chronic pain for ICD-11: chronic postsurgical or posttraumatic pain. Pain 160(1):45–52
- Hoffmann H, Walther D, Bittner R, Köckerling F, Adolf D, Kirchhoff P (2020) Smaller inguinal hernias are independent risk factors for developing chronic postoperative inguinal pain (CPIP): a registry-based multivariable analysis of 57,999 patients. Ann Surg 271(4):756–764
- Taylor C, Layani L, Liew V, Ghusn M, Crampton N, White S (2008) Laparoscopic inguinal hernia repair without mesh fixation, early results of a large randomised clinical trial. Surg Endosc 22(3):757–762
- Liem MSL, Van Duyn EB, Van Der Graaf Y, Van Vroonhoven T (2003) Recurrences after conventional anterior and laparoscopic inguinal hernia repair: a randomized comparison. Ann Surg 237(1):136–141

- 40. Papanicolas I, Woskie LR, Jha AK (2018) Health care spending in the United States and other high-income countries. JAMA 319(10):1024
- Glasgow RE, Mulvihill SJ, Pettit JC, Young J, Smith BK, Vargo DJ, Ray DM, Finlayson SRG (2021) Value analysis of methods of inguinal hernia repair. Ann Surg 274(4):572–580

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.