

Rachel J. Pearl · Peter C. Papageorgiou ·  
Michael Goldman · Akis A. Amfilochiadis ·  
Frans Boomsma · Rasmus Rojkjaer · Denis Geary ·  
Daniel H. Osmond

## Possible role of new pressor protein in hypertensive anephric hemodialysis patients

Published online: 31 October 2003  
© IPNA 2003

### Pediatr Nephrol (2003) 18:1025–1031

The acknowledgements in the above article are erroneous.  
They should read:

**Acknowledgements** These results were presented at the American Society of Nephrology 35th Annual Meeting and Scientific Exposition, 1–4 November 2002, Philadelphia, Pa., USA, and at

the American Society of Hypertension, 18th Annual Scientific Meeting, 14–17 May 2003, New York, NY, USA. This study was supported by the Heart and Stroke Foundation of Ontario (HSFO) grants NA 3478 and T4136. Peter C. Papageorgiou is a recipient of an AHEPA CANADA and HCFO scholarship.

The online version of the original article can be found at  
<http://dx.doi.org/10.1007/s00467-003-1246-6>

R. J. Pearl (✉) · M. Goldman · D. Geary  
Division of Nephrology,  
The Hospital for Sick Children,  
555 University Avenue, Toronto, Ontario, M5G 1X8, Canada  
e-mail: rachel.pearl@sickkids.ca  
Tel.: +1-416-8137654-1458  
Fax: +1-416-8136271

P. C. Papageorgiou · A. A. Amfilochiadis · D. H. Osmond  
Department of Physiology and Medicine,  
University of Toronto,  
Canada

P. C. Papageorgiou · A. A. Amfilochiadis · D. H. Osmond  
Heart and Stroke Richard Lewar Centre of Excellence,  
University of Toronto,  
Canada

F. Boomsma  
Department of Internal Medicine,  
Erasmus MC,  
Rotterdam, The Netherlands

R. Rojkjaer  
Hemostasis Biology,  
Novo Nordisk,  
Maaloev, Denmark