

Erratum to: Normal-range albuminuria does not exclude nephropathy in diabetic children

Jacek Zachwieja · Jolanta Soltysiak · Piotr Fichna · Katarzyna Lipkowska ·
Witold Stankiewicz · Bogda Skowronska · Paweł Kroll ·
Maria Lewandowska-Stachowiak

Published online: 10 April 2010
© IPNA 2010

Erratum to: Pediatr Nephrol
DOI 10.1007/s00467-010-1443-z

It has come to our attention that the reported NGAL levels are 10-fold higher than those reported by other researchers. This is best seen from the normal values reported in Table 1. Inaccurate data is due to a decimal mistake. All reported serum and urine values for NGAL have to be divided by 10.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00467-010-1443-z>.

J. Zachwieja (✉) · J. Soltysiak · K. Lipkowska ·
M. Lewandowska-Stachowiak
Department of Pediatric Nephrology,
Poznan University of Medical Sciences,
Szpitalna 27/33 str.,
60-572 Poznan, Poland
e-mail: j.zachwieja@mp.pl

P. Fichna · W. Stankiewicz · B. Skowronska
Department of Pediatric Endocrinology and Diabetes,
Poznan University of Medical Sciences,
Poznan, Poland

P. Kroll
Department of Pediatric Urology,
Poznan University of Medical Sciences,
Poznan, Poland