

Erratum to: Hypertension in infancy: diagnosis, management and outcome

Janis M. Dionne · Carolyn L. Abitbol · Joseph T. Flynn

Published online: 10 August 2011
© IPNA 2011

Erratum to: *Pediatr Nephrol*
DOI 10.1007/s00467-010-1755-z

There was an Error in Table 2

The correct Table 2 is:

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00467-010-1755-z>.

J. M. Dionne
Division of Nephrology, Department of Pediatrics,
University of British Columbia, BC Children's Hospital,
Vancouver, BC, Canada

C. L. Abitbol
Division of Pediatric Nephrology,
Department of Pediatrics, University of Miami Miller School
of Medicine/ Holtz Children's Hospital,
Miami, FL, USA

J. T. Flynn (✉)
Division of Nephrology, Seattle Children's Hospital, Department
of Pediatrics, University of Washington School of Medicine,
4800 Sand Point Way NE, M/S A-7931,
Seattle, WA 98105, USA
e-mail: joseph.flynn@seattlechildrens.org

Table 2 Neonatal blood pressures and potential treatment parameters*

Post-conceptual age	50th percentile	95th percentile	99th percentile
44 weeks			
SBP	88	105	110
DBP	50	68	73
MAP	63	80	85
42 weeks			
SBP	85	98	102
DBP	50	65	70
MAP	62	76	81
40 weeks			
SBP	80	95	100
DBP	50	65	70
MAP	60	75	80
38 weeks			
SBP	77	92	97
DBP	50	65	70
MAP	59	74	79
36 weeks			
SBP	72	87	92
DBP	50	65	70
MAP	57	72	77
34 weeks			
SBP	70	85	90
DBP	40	55	60
MAP	50	65	70
32 weeks			
SBP	68	83	88
DBP	40	55	60
MAP	49	64	69
30 weeks			
SBP	65	80	85
DBP	40	55	60
MAP	48	63	68
28 weeks			
SBP	60	75	80
DBP	38	50	54
MAP	45	58	63
26 weeks			
SBP	55	72	77
DBP	30	50	56
MAP	38	57	63

*This Table provides estimated values for blood pressures after 2 weeks of age in infants from 26 to 44 weeks post conceptual age. The 95th and 99th percentile values are intended to serve as a reference to identify infants with persistent hypertension that may require treatment