Correction to: Mode of initial renal replacement therapy and transplant outcomes in the chronic kidney disease in children (CKiD) study

Meredith A. Atkinson¹ · Jennifer L. Roem² · Anuradha Gajjar³ · Bradley A. Warady⁴ · Susan L. Furth⁵ · Alvaro Muñoz²

Published online: 2 March 2020
© IPNA 2020

Correction to: *Pediatr Nephrol*

<https://doi.org/10.1007/s00467-019-04416-2>

The original version of this article unfortunately contained a mistake. Figures 1 and 2 contained undesired lines as a result of reproduction. The corrected figures are presented below.

The online version of the original article can be found at <https://doi.org/10.1007/s00467-019-04416-2>

✉ Meredith A. Atkinson
matkins3@jhmi.edu

¹ Division of Pediatric Nephrology, Johns Hopkins University School of Medicine, 200 N. Wolfe St., 3rd Floor, Baltimore, MD 21287, USA

² Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

³ Weill Cornell Medical College, New York, NY, USA

⁴ Children's Mercy Hospital, Kansas City, MO, USA

⁵ Children's Hospital of Philadelphia, Philadelphia, PA, USA

Fig. 1 Cumulative incidence of transplantation (solid lines) following dialysis as first RRT event in 123 CKiD subjects. The black lines represent the 81 HD participants of whom 50% were transplanted by 11 months after initiating dialysis. The gray lines are the 42 PD participants of whom 50% were transplanted 13 months after initiating dialysis. The dashed lines represent the generalized gamma curves yielding a likelihood ratio statistic with a p value = 0.028

Fig. 2 Cumulative incidence of transplant failure following transplant event in 170 CKiD subjects. Solid lines depict unadjusted functions and dash-dotted lines represent adjusted functions using inverse probability weights. The gray lines are the 79 PKT participants of whom only 5% had transplant failure by 48 months. The black lines represent the 91 dialysis-experienced participants of whom 16% had transplant failure by 48 months. Incidence of transplant failure marginally differs between dialysis-experienced and PKT when unadjusted ($p = 0.092$), but attenuates when adjusted by inverse probability weights ($p = 0.206$)