



Correction to: C3 levels and neurologic involvement in hemolytic uremic syndrome related to Shiga toxin-producing *Escherichia coli*

Alejandro Balestracci¹ · Luciana Meni Battaglia¹

Published online: 23 August 2020
© IPNA 2020

Correction to: *Pediatr Nephrol*

<https://doi.org/10.1007/s00467-020-04717-x>

The original version of the letter unfortunately contained a mistake. The unit of measurement used to express C3 concentrations in the text and in Figure 1 should be g/L instead of g/dL. The corrected Figure 1 is presented below.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.




Fig. 1 Receiver operating characteristic (ROC) curve for serum C3 levels on admission as predictor of central nervous system involvement in STEC-HUS patients. Area under the ROC curve (AUC) = 0.76, CI 95% 0.617–0.87, $p = 0.0024$. C3 level ≤ 0.85 g/L was the best cut-off value (83.78% specificity, 75% sensitivity)

The online version of the original article can be found at <https://doi.org/10.1007/s00467-020-04717-x>

✉ Alejandro Balestracci
abalestracci@yahoo.com.ar

¹ Nephrology Unit, Hospital General de Niños Pedro de Elizalde, Montes de Oca 40, 1270 Ciudad Autónoma de Buenos Aires, Argentina