

Adzo Dzifa Kokutse · Alexia Stokes · Henri Baillères ·  
Kouami Kokou · Christine Baudasse

## Decay resistance of Togolese teak (*Tectona grandis* L.f) heartwood and relationship with colour

Published online: 14 March 2006  
© Springer-Verlag 2006

---

**Trees (2006) 20:219–223**

In the abstract, the sentence “Inner heartwood was found to be the most resistant to pathogen attack and outer heartwood the least.” should be replaced by “Inner heartwood was found to be the least resistant to pathogen attack and outer heartwood the most.”

---

The online version of the original article can be found at  
<http://dx.doi.org/10.1007/s00468-005-0028-0>

---

A. D. Kokutse · A. Stokes (✉)  
Laboratoire de Rhéologie des Bois de Bordeaux (Mixed unit:  
Université Bordeaux I/INRA/CNRS),  
69 rte d’Arcachon,  
33612 Cestas cedex, France  
e-mail: stokes@lrbb3.pierroton.inra.fr  
Tel.: +33-5-57122821  
Fax: +33-5-56680713

A. D. Kokutse · K. Kokou  
Laboratoire de Botanique et Ecologie Végétale, Faculté des  
Sciences, Université de Lomé,  
B.P.1515 Togo, France

H. Baillères · C. Baudasse  
CIRAD-Forêt,  
TA 80/03,  
34398 Montpellier cedex 5, France