## **CORRECTION**



## Correction to: In vitro regeneration of mulberry plants from seedling explants of *Morus indica* cv. G4 through direct organogenesis

Tanmoy Sarkar<sup>1</sup> · K. N. Ravindra<sup>1</sup> · S. Gandhi Doss<sup>1</sup> · P. M. Pratheesh Kumar<sup>1</sup> · Pankaj Tewary<sup>1</sup>

Published online: 31 August 2021 © The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2021

**Correction to: Trees** 

https://doi.org/10.1007/s00468-021-02186-9

The table 1 has been published with corrections in the original publication. The complete corrected Table 1 is given below. The original article has been corrected.

The original article can be found online at https://doi.org/10.1007/s00468-021-02186-9.



<sup>☐</sup> Tanmoy Sarkar tanmoy.dgr@gmail.com

<sup>&</sup>lt;sup>1</sup> Central Sericultural Research & Training Institute (CSR&TI), Mysuru 570008, Karnataka, India

2144 Trees (2021) 35:2143–2144

**Table 1** Effects of various concentrations of TDZ on the formation of shoot primordia from cotyledons and hypocotyls during the two phases of pre-culturing

First pre-culturing of	f explants for 2	2 days						
Treatment	T1 MS medium without TDZ		T2 MS medium plus 0.1 mg/L TDZ		T3 MS medium plus 1.1 mg/L TDZ		T4 MS medium plus 0.1 mg/L TDZ	
Media composition								
	Cotyledon	Hypocotyl	Cotyledon	Hypocotyl	Cotyledon	Hypocotyl	Cotyledon	Hypocotyl
Induction of Shoot primordia (%)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Second pre-culturing	g of explants fo	or 3 days						
Treatment	T1		T2		T3		T4	
Media composition	MS medium without TDZ		MS medium plus 0.1 mg/L TDZ		MS medium plus 1.1 mg/L TDZ		MS medium plus 1.1 mg/L TDZ	
	Cotyledon	Hypocotyl	Cotyledon	Hypocotyl	Cotyledon	Hypocotyl	Cotyledon	Hypocotyl
Induction of Shoot primordia (%)	$0.00 \pm 0.00^{\rm e}$	$0.00 \pm 0.00^{\rm e}$	53.33 ± 3.33 <sup>b</sup>	$8.88 \pm 1.11^{\text{de}}$	$57.04 \pm 5.00^{ab}$	15.29 ± 3.56 <sup>cd</sup>	$65.55 \pm 2.94^{a}$	22.22 ± 2.22°

Data represent mean  $\pm$  standard error (SE) of three repeated experiments, each with 25 explants per explant type. The values followed by different letters for each variable are significantly different at  $p \le 0.05$  according to Tukey's multiple comparison test

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

