



Bubbling analysis for approximate Lorentzian harmonic maps from Riemann surfaces

Xiaoli Han¹ · Jürgen Jost² · Lei Liu² · Liang Zhao³

Received: 3 May 2017 / Accepted: 25 October 2017 / Published online: 7 November 2017
© The Author(s) 2017. This article is an open access publication

Abstract For a sequence of approximate harmonic maps (u_n, v_n) (meaning that they satisfy the harmonic system up to controlled error terms) from a compact Riemann surface with smooth boundary to a standard static Lorentzian manifold with bounded energy, we prove that identities for the Lorentzian energy hold during the blow-up process. In particular, in the special case where the Lorentzian target metric is of the form $g_N - \beta dt^2$ for some Riemannian metric g_N and some positive function β on N , we prove that such identities also hold for the positive energy (obtained by changing the sign of the negative part of the Lorentzian energy) and there is no neck between the limit map and the bubbles. As an application, we complete the blow-up picture of singularities for a harmonic map flow into a standard static Lorentzian manifold. We prove that the energy identities of the flow hold at both finite and infinite singular times. Moreover, the no neck property of the flow at infinite singular time is true.

Mathematics Subject Classification 53C43 · 58E20

Communicated by Luigi Ambrosio.

The research is supported by NSFC Nos. 11471014, 11471299 and the Fundamental Research Funds for the Central Universities. The authors would like to thank the referee for detailed and useful comments.

✉ Jürgen Jost
jost@mis.mpg.de

Xiaoli Han
xlhan@math.tsinghua.edu.cn

Lei Liu
leiliu@mis.mpg.de; llei1988@mail.ustc.edu.cn

Liang Zhao
liangzhao@bnu.edu.cn

¹ Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China

² Max Planck Institute for Mathematics in the Sciences, Inselstrasse 22, 04103 Leipzig, Germany

³ School of Mathematical Sciences, Laboratory of Mathematics and Complex Systems of Ministry of Education, Beijing Normal University, Beijing 100875, China

1 Introduction

Harmonic maps constitute one of the fundamental objects in the field of geometric analysis. When the domain is two-dimensional, particularly interesting features arise. The conformal invariance of the energy functional leads to non-compactness of the set of harmonic maps in dimension two, and the blow-up behavior has been studied extensively in [5, 13, 20, 23, 24, 27] for the interior case and [10, 15, 16] for the boundary case. Roughly speaking, the energy identities for harmonic maps tell us that, during the weak convergence of a sequence of harmonic maps, the loss of energy is concentrated at finitely many points and can be quantized by a sum of energies of harmonic spheres and harmonic disks. Also for many other elliptic and parabolic nonlinear variational problems arising in geometry and physics, such as J-holomorphic curves or Yang–Mills fields, to understand the convergence properties of a sequence and the emergence of singularities is of special importance.

In physics, harmonic maps arise as a mathematical representation of the nonlinear sigma model and this leads to several generalizations. For example, motivated by the supersymmetric sigma model, Dirac harmonic maps where a map is coupled with a spinor field have been extensively studied. One can refer to [4, 14, 31] and the references therein. From the perspective of general relativity, it is also natural to generalize the target of a harmonic map to a Lorentzian manifold. Recent work on minimal surfaces in anti-de-Sitter space and their applications in theoretical physics (see e.g. Alday and Maldacena [1]) shows the importance of this extension. Geometrically, the link between harmonic maps into S^4_+ and the conformal Gauss maps of Willmore surfaces in S^3 [3] also naturally leads to such harmonic maps.

Thus, in this paper, we investigate harmonic maps from Riemann surfaces into Lorentzian manifolds. In order to gain some special structure, we consider a Lorentzian manifold $N \times \mathbb{R}$ that is equipped with a warped product metric of the form

$$g = g_N - \beta(d\theta + \omega)^2, \tag{1.1}$$

where $(\mathbb{R}, d\theta^2)$ is the 1-dimensional Euclidean space, (N, g_N) is an n -dimensional compact Riemannian manifold which by Nash’s theorem can be isometrically embedded into some \mathbb{R}^K , β is a positive C^∞ function on N and ω is a smooth 1-form on N . Since N is compact, β and ω are both bounded on N . We suppose for any $p \in N$,

$$0 < \lambda_1 < \beta(p) < \lambda_2, \quad |\omega(p)| + |\nabla\omega(p)| + |\nabla\beta(p)| \leq \lambda_2.$$

A Lorentzian manifold with a metric of the form (1.1) is called a standard static manifold. For more details on such manifolds, we refer to [17, 22].

Let (M, h) be a compact Riemann surface with smooth boundary ∂M . For a map $(u, v) \in C^2(M, N \times \mathbb{R})$ with fixed boundary data $(u, v)|_{\partial M} = (\phi, \psi)$, we define the functional

$$E_g(u, v) = \frac{1}{2} \int_M \left\{ |\nabla u|^2 - \beta(u)|\nabla v + \omega_i(u)\nabla u^i|^2 \right\} dv_h, \tag{1.2}$$

which is called the *Lorentzian energy* of the map (u, v) on M . Critical points (u, v) in $C^2(M, N \times \mathbb{R})$ of the functional (1.2) are called Lorentzian harmonic maps from (M, h) into the Lorentzian manifold $(N \times \mathbb{R}, g)$. Besides the Lorentzian energy $E_g(u, v)$, we also consider

$$E(u) = \frac{1}{2} \int_M |\nabla u|^2 dv_h \quad \text{and} \quad E(u, v) = \frac{1}{2} \int_M \{ |\nabla u|^2 + |\nabla v|^2 \} dv_h \tag{1.3}$$

and call it the *positive energy* of u and (u, v) on M respectively. It is obvious that both the Lorentzian and positive energy functionals are conformally invariant when $\dim M = 2$.

Zhu [32] has derived the Euler–Lagrange equations for (1.2),

$$\Delta u + A(u)(\nabla u, \nabla u) - H^\top = 0 \text{ in } M, \tag{1.4}$$

$$\operatorname{div} \left(\beta(u)(\nabla v + \omega_i \nabla u^i) \right) = 0 \text{ in } M \tag{1.5}$$

with the boundary data

$$(u, v)|_{\partial M} = (\phi, \psi) \in C^{2+\alpha}(\partial M, N \times \mathbb{R}) \tag{1.6}$$

for some $\alpha \in (0, 1)$. Here A is the second fundamental form of N in \mathbb{R}^K , H^\top is the tangential part of $H = (H^1, \dots, H^K)$ along the map (u, v) with

$$H^j := \beta(\nabla v + \omega_i \nabla u^i) \cdot \nabla u^k \left(\frac{\partial \omega_j}{\partial y^k} - \frac{\partial \omega_k}{\partial y^j} \right) - \frac{1}{2} \frac{\partial \beta}{\partial y^j} |\nabla v + \omega_i \nabla u^i|^2, \quad j = 1, \dots, K.$$

Let us now recall some related results. The existence of geodesics in Lorentzian manifolds was studied in [2]. Variational methods for such harmonic maps were developed in [6, 7]. Recently, [8] studied the corresponding heat flow under the assumption that $\omega \equiv 0$ and proved the existence of a Lorentzian harmonic map in any given homotopic class under either some geometric conditions on N or a small energy condition of the initial maps. The regularity theory of Lorentzian harmonic maps was studied in [11, 12, 19, 32].

In [9], the authors proved identities of the Lorentzian energy for a blow-up sequence of Lorentzian harmonic maps when M is a compact Riemann surface without boundary. They showed the tangential Lorentzian energy of the sequence in the neck region has no concentration by comparing the energy with piece-wise linear functions (i.e. geodesics). Then they used the Hopf differentials to control the radial Lorentzian energy.

In any case, the analysis of Lorentzian harmonic maps is more difficult than that of standard (Riemannian) harmonic maps, because one cannot no longer use positivity properties of the target metric. This is a technical reason why we restrict ourselves to standard static Lorentzian manifolds.

In this paper, we shall prove some energy identities of an approximate Lorentzian harmonic map sequence and get the no neck property during a blow-up process when M is a compact Riemann surface with boundary. We work with approximate sequences which means that we allow for error terms in the Lorentzian harmonic maps system. The reason is that this has a direct application in studying the singularities of the parabolic version, the Lorentzian harmonic map flow (see [8]). Moreover, since we assume that the domain M is a manifold with boundary, blow-up analysis on the boundary must be included in our case. Here, we will use the method of integrating by parts (cf. [20] for harmonic maps) to prove a Pohozaev type identity instead of using the Hopf differential. The Pohozaev identity method is more general and powerful than the Hopf differential method. We first prove identities for the Lorentzian energy for a blow-up sequence of approximate Lorentzian harmonic maps. Furthermore, for the special case $\omega \equiv 0$, we show that also such identities for the positive energy and no neck properties hold.

Throughout this paper, we call a map into $N \times \mathbb{R}$ a Lorentzian map and when we have a map into the Riemannian manifold N , we just call it a map. We first give the definition of an approximate Lorentzian harmonic map.

Definition 1.1 $(u, v) \in W^{2,2}(M, N \times \mathbb{R})$ is called an approximate Lorentzian harmonic map with Dirichlet boundary data (ϕ, ψ) , if there exist fields $(\tau(u, v), \kappa(u, v)) \in L^1(M)$ such that (u, v) satisfies

$$\Delta u + A(u)(\nabla u, \nabla u) - H^\top = \tau(u, v) \text{ in } M, \tag{1.7}$$

$$\operatorname{div} \left(\beta(u)(\nabla v + \omega_i \nabla u^i) \right) = \kappa(u, v) \text{ in } M \tag{1.8}$$

with the boundary condition $(u, v)|_{\partial M} = (\phi, \psi)$.

Now we can present our first main result.

Theorem 1.1 *Let $(u_n, v_n) \in W^{2,2}(M, N \times \mathbb{R})$ be a sequence of approximate harmonic maps with Dirichlet boundary $(u_n, v_n)|_{\partial M} = (\phi, \psi) \in C^{2+\alpha}(\partial M, N \times \mathbb{R})$ satisfying*

$$E(u_n, v_n) + \|(\tau_n, \kappa_n)\|_{L^2(M)} \leq \Lambda < \infty,$$

where $\|(\tau_n, \kappa_n)\|_{L^2(M)}^2 = \|\tau_n\|_{L^2(M)}^2 + \|\kappa_n\|_{L^2(M)}^2$. After taking a subsequence, still denoted by $\{u_n, v_n\}$, we can find a finite set $\mathcal{S} = \{p_1, \dots, p_I\}$ and a limit map $(u_0, v_0) \in W^{1,2}(M, N \times \mathbb{R})$ with Dirichlet boundary data $(u_0, v_0)|_{\partial M} = (\phi, \psi)$ such that $\{(u_n, v_n)\}$ converges weakly in $W_{loc}^{2,2}(M \setminus \mathcal{S})$ to (u_0, v_0) . Moreover, there are finitely many nontrivial Lorentzian harmonic spheres $(\sigma_i^l, \xi_i^l) : \mathbb{S}^2 \rightarrow N \times \mathbb{R}$ and nontrivial Lorentzian harmonic maps $(\sigma_i^k, \xi_i^k) : \mathbb{R}_+^2 := \{(x^1, x^2) \in \mathbb{R}^2 | x^2 \geq 0\} \rightarrow N \times \mathbb{R}$ with constant boundary values, where $i = 1, \dots, I$, $l = 1, \dots, l_i$ and $k = 1, \dots, k_i$ with $l_i, k_i \geq 0$ and $l_i + k_i \geq 1$, such that

$$\lim_{n \rightarrow \infty} E_g(u_n, v_n) = E_g(u_0, v_0) + \sum_{i=1}^I \sum_{l=1}^{l_i} E_g(\sigma_i^l, \xi_i^l) + \sum_{i=1}^I \sum_{k=1}^{k_i} E_g(\sigma_i^k, \xi_i^k). \tag{1.9}$$

Here and in the sequel, “finite” includes “possibly empty”, that is, singularities need not always arise. Since this is obvious, it will not be explicitly mentioned.

When $\omega \equiv 0$, the equations for Lorentzian harmonic maps become

$$\Delta u + A(u)(\nabla u, \nabla u) - B^\top(u)|\nabla v|^2 = 0 \text{ in } M, \tag{1.10}$$

$$\operatorname{div}(\beta(u)\nabla v) = 0 \text{ in } M \tag{1.11}$$

where $B(u) := (B^1, B^2, \dots, B^K)$ with

$$B^j := -\frac{1}{2} \frac{\partial \beta(u)}{\partial y^j}$$

and B^\top is the tangential part of B along the map u . In this case, the blow-up behavior is simpler. We show that the identities for the positive energy hold and there is no neck during the process.

Theorem 1.2 *If we additionally assume $\omega \equiv 0$ in Theorem 1.1, there cannot emerge any Lorentzian harmonic maps $(\sigma_i^k, \xi_i^k) : \mathbb{R}_+^2 := \{(x^1, x^2) \in \mathbb{R}^2 | x^2 \geq 0\} \rightarrow N \times \mathbb{R}$ during the blow-up process [i.e. $k_i = 0$ in (1.9)]. Moreover, the components ξ_i^l of the maps (σ_i^l, ξ_i^l) are constant and $\sigma_i^l : \mathbb{S}^2 \rightarrow N$, $1 \leq l \leq l_i$ are nontrivial harmonic spheres. In this case, (1.9) becomes*

$$\lim_{n \rightarrow \infty} E(u_n) = E(u_0) + \sum_{i=1}^I \sum_{l=1}^{l_i} E(\sigma_i^l), \tag{1.12}$$

$$\lim_{n \rightarrow \infty} E(v_n) = E(v_0). \tag{1.13}$$

and the image $u_0(M) \cup_{i=1}^I \cup_{l=1}^{l_i} (\sigma_i^l(\mathbb{S}^2))$ is a connected set in N .

As an application of Theorem 1.2, we consider a harmonic map heat flow

$$\begin{cases} \partial_t u = \Delta u + A(u)(\nabla u, \nabla u) - B^\top(u)|\nabla v|^2, & \text{in } M \times [0, T) \\ -\operatorname{div}(\beta(u)\nabla v) = 0, & \text{in } M \times [0, T) \end{cases} \tag{1.14}$$

with the boundary-initial data

$$\begin{cases} u(x, t) = \phi_0(x) & \text{on } M \times \{t = 0\}, \\ u(x, t) = \phi(x), & \text{on } \partial M \times \{t > 0\}, \\ v(x, t) = \psi(x), & \text{on } \partial M \times \{t > 0\}, \\ \phi_0(x) = \phi(x) & \text{on } \partial M. \end{cases} \tag{1.15}$$

This kind of harmonic map heat flow is a parabolic-elliptic system and was first studied in [8]. We proved the problem (1.14) and (1.15) admits a unique solution $(u, v) \in \mathcal{V}(M_0^{T_1}; N \times \mathbb{R})$ (see the notation at the end of this section), where T_1 is the first singular time and some bubbles (nontrivial harmonic spheres) split off at $t = T_1$. In this paper, we complete the blow-up picture at the singularities of this flow. First, we have

Theorem 1.3 *Suppose the problem (1.14) and (1.15) admits a unique global solution $(u, v) \in \mathcal{V}(M_0^\infty; N \times \mathbb{R})$ which blows up at infinity, i.e. $T_1 = \infty$. By choosing some $t_n \rightarrow \infty$, there exist a smooth Lorentzian harmonic map $(u_\infty, v_\infty) : M \rightarrow N \times \mathbb{R}$ with boundary data $(u_\infty, v_\infty)|_{\partial M} = (\phi, \psi)$ and finitely many harmonic spheres $\{\sigma^i\}_{i=1}^L : \mathbb{R}^2 \cup \{\infty\} \rightarrow N$ such that*

$$\lim_{n \rightarrow \infty} E(u(t_n); M) = E(u_\infty, M) + \sum_{i=1}^L E(\sigma^i), \tag{1.16}$$

$$\lim_{t \rightarrow \infty} E_g(u(t), v(t); M) = E_g(u_\infty, v_\infty; M) + \sum_{i=1}^L E(\sigma^i). \tag{1.17}$$

Furthermore, there exist sequences $\{x_n^i\}_{i=1}^L \subset M$ and $\{r_n^i\}_{i=1}^L \subset \mathbb{R}_+$ such that

$$\lim_{n \rightarrow \infty} \|u(\cdot, t_n) - u_\infty(\cdot) - \sum_{i=1}^L \sigma_n^i(\cdot)\|_{L^\infty(M)} = 0, \tag{1.18}$$

where $\sigma_n^i(\cdot) = \sigma^i\left(\frac{\cdot - x_n^i}{r_n^i}\right) - \sigma^i(\infty)$.

When the flow blows up at finite time, we have

Theorem 1.4 *Let $(u, v) \in \mathcal{V}(M_0^{T_1}; N \times \mathbb{R})$ be a solution to (1.14) and (1.15) with $T_1 < \infty$ as its first singular time. Then there exist finitely many harmonic spheres $\{\sigma^i\}_{i=1}^L : \mathbb{R}^2 \cup \{\infty\} \rightarrow N$ such that*

$$\lim_{t \nearrow T_1} E(u(t); M) = E(u(T_1), M) + \sum_{i=1}^L E(\sigma^i), \tag{1.19}$$

$$\lim_{t \nearrow T_1} E_g(u(t), v(t); M) = E_g(u(T_1), v(T_1); M) + \sum_{i=1}^L E(\sigma^i), \tag{1.20}$$

where $(u(T_1), v(T_1))$ is the weak limit of $(u(t), v(t))$ in $W^{1,2}(M)$ as $t \rightarrow T_1$.

The paper is organized as follows. In Sect. 2, we derive some basic lemmas including the small energy regularity, a Pohozaev type identity and a removable singularity result. In Sect. 3, we prove the energy identities and no neck property for a sequence of approximate Lorentzian harmonic maps (Theorems 1.1, 1.2). In Sect. 4, we apply these two results to the harmonic map heat flow and prove Theorems 1.3 and 1.4. Throughout this paper, we use C to denote a universal constant and denote $D_1(0) := \{(x, y) \in \mathbb{R}^2 \mid |x|^2 + |y|^2 \leq 1\}$, $D_1^+(0) := \{(x, y) \in \mathbb{R}^2 \mid |x|^2 + |y|^2 \leq 1, y \geq 0\}$, $\partial^0 D_1^+(0) := \{(x, y) \in D_1^+(0) \mid y = 0\}$ and $\partial^+ D_1^+(0) := \{(x, y) \in \partial D_1^+(0) \mid y > 0\}$.

Notation We denote

$$\begin{aligned} \mathcal{V}(M_s^t; N \times \mathbb{R}) &:= \{(u, v) : M \times [s, t] \rightarrow N \times \mathbb{R}, v \in L^\infty([s, t]; C^{2+\alpha}(M)), \\ &v, \nabla v \in \cap_{s < \rho < t} C^{\alpha, \alpha/2}(M \times [s, \rho]), \\ &u \in \cap_{s < \rho < t} C^{2+\alpha, 1+\alpha/2}(M \times [s, \rho])\}. \end{aligned}$$

2 some basic lemmas

In this section, we will prove some basic lemmas for Lorentzian harmonic maps, such as the small energy regularity, a Pohozaev type identity and a removable singularity result.

First, we present two small energy regularity lemmas corresponding to the interior case and the boundary case. For harmonic maps, such results have been obtained in [5, 27] for the interior case and in [10, 15, 16] for the boundary case. We use

$$\bar{u}_\Omega = \frac{1}{|\Omega|} \int_\Omega u dx$$

to denote the average value of a function u on the domain Ω . Here and in the sequel, we shall view (ϕ, ψ) as the restriction of some $C^{2+\alpha}(M, N \times \mathbb{R})$ map on ∂M and for simplicity, we still denote it by (ϕ, ψ) .

Lemma 2.1 *Let $(u, v) \in W^{2,p}(D, N \times \mathbb{R})$, $1 < p \leq 2$ be an approximate Lorentzian harmonic map with $(\tau, \kappa) \in L^p(D)$. There exist constants $\epsilon_1 = \epsilon_1(p, \lambda_1, \lambda_2, N) > 0$ and $C = C(p, \lambda_1, \lambda_2, N) > 0$, such that if $E(u, v; D) \leq \epsilon_1$, we have*

$$\begin{aligned} &\|u - \bar{u}_{D_{1/2}}\|_{W^{2,p}(D_{1/2})} + \|v - \bar{v}_{D_{1/2}}\|_{W^{2,p}(D_{1/2})} \\ &\leq C(\|\nabla u\|_{L^2(D)} + \|\nabla v\|_{L^2(D)} + \|\tau\|_{L^p(D)} + \|\kappa\|_{L^p(D)}). \end{aligned}$$

Moreover, by the Sobolev embedding $W^{2,p} \hookrightarrow C^0$, we have

$$\|u\|_{osc(D_{1/2})} = \sup_{x, y \in D_{1/2}} |u(x) - u(y)| \leq C(\|\nabla u, \nabla v\|_{L^2(D)} + \|(\tau, \kappa)\|_{L^p(D)}).$$

For the boundary case, we have

Lemma 2.2 *Let $(u, v) \in W^{2,p}(D^+, N \times \mathbb{R})$, $1 < p \leq 2$ be an approximate Lorentzian harmonic map with $(\tau, \kappa) \in L^p(D^+)$. On the boundary we assume that $u|_{\partial^0 D^+} = \phi(x)$ and $v|_{\partial^0 D^+} = \psi(x)$ where $(\phi, \psi) \in C^{2+\alpha}(D)$. There exist constants $\epsilon_2 = \epsilon_2(p, \lambda_1, \lambda_2, N) > 0$ and $C = C(p, \lambda_1, \lambda_2, N) > 0$, such that if $E(u, v; D_1^+) \leq \epsilon_2$, we have*

$$\begin{aligned} &\|u - \bar{\phi}_{\partial^0 D^+}\|_{W^{2,p}(D_{1/2}^+)} + \|v - \bar{\psi}_{\partial^0 D^+}\|_{W^{2,p}(D_{1/2}^+)} \\ &\leq C(\|(\nabla u, \nabla v)\|_{L^2(D^+)} + \|(\nabla \phi, \nabla \psi)\|_{W^{1,p}(D^+)} + \|(\tau, \kappa)\|_{L^p(D^+)}), \end{aligned}$$

where $\bar{\phi}_{\partial^0 D^+} = \frac{1}{2} \int_{\partial^0 D^+} \phi dx$ and $\bar{\psi}_{\partial^0 D^+} = \frac{1}{2} \int_{\partial^0 D^+} \psi dx$.

Moreover, by the Sobolev embedding $W^{2,p} \hookrightarrow C^0$, we have

$$\begin{aligned} \|u\|_{osc(D^+_{1/2})} &= \sup_{x,y \in D^+_{1/2}} |u(x) - u(y)| \\ &\leq C(\|\nabla u, \nabla v\|_{L^2(D^+)} + \|(\nabla\phi, \nabla\psi)\|_{W^{1,p}(D^+)} + \|(\tau, \kappa)\|_{L^p(D^+)}). \end{aligned}$$

Since the proof of the interior case is similar to and simpler than that of the boundary case, we only prove Lemma 2.2 and omit the proof of Lemma 2.1.

Proof Without loss of generality, we assume $\bar{\phi}_{\partial^0 D^+} = \bar{\psi}_{\partial^0 D^+} = 0$. Choosing a cut-off function $\eta \in C^\infty_0(D^+)$ satisfying $0 \leq \eta \leq 1$, $\eta|_{D^+_{3/4}} \equiv 1$, $|\nabla\eta| + |\nabla^2\eta| \leq C$ and computing directly, we get

$$\begin{aligned} |\Delta(\eta u)| &= |\eta\Delta u + 2\nabla\eta\nabla u + u\Delta\eta| \\ &\leq C(|u| + |\nabla u| + (|\nabla u| + |\nabla v|)(|\eta\nabla u| + |\eta\nabla v|) + |\tau|) \\ &\leq C(|\nabla u| + |\nabla v|)(|\nabla(\eta u)| + |\nabla(\eta v)|) \\ &\quad + C(|u| + (1 + |v|)(|\nabla u| + |\nabla v|) + |\tau|). \end{aligned}$$

Similarly,

$$\begin{aligned} |\Delta(\eta v)| &= |\eta\Delta v + 2\nabla\eta\nabla v + v\Delta\eta| \\ &\leq C(|v| + |\nabla v| + (|\nabla u| + |\nabla v|)(|\eta\nabla u| + |\eta\nabla v|) + |\tau| + |\kappa|) \\ &\leq C(|\nabla u| + |\nabla v|)(|\nabla(\eta u)| + |\nabla(\eta v)|) \\ &\quad + C(|v| + (1 + |v|)(|\nabla u| + |\nabla v|) + |\tau| + |\kappa|). \end{aligned}$$

First we assume that $1 < p < 2$. By standard elliptic estimates and Poincare’s inequality, we obtain

$$\begin{aligned} &\|\eta u\|_{W^{2,p}(D)} + \|\eta v\|_{W^{2,p}(D)} \\ &\leq C\|(\nabla u, \nabla v)\|_{L^2(D^+)}\|(\nabla(\eta u), \nabla(\eta v))\|_{L^{\frac{2p}{2-p}}(D^+)} + C\|(u, v)\|_{W^{1,p}(D^+)} \\ &\quad + C\left(\|(\nabla u, \nabla v)\|_{L^2(D^+)}\|v\|_{L^{\frac{2p}{2-p}}(D^+)} + \|(\phi, \psi)\|_{W^{2,p}(D^+)} + \|(\tau, \kappa)\|_{L^p(D^+)}\right) \\ &\leq C\epsilon_2\|(\nabla(\eta u), \nabla(\eta v))\|_{L^{\frac{2p}{2-p}}(D^+)} + C(\|(\nabla u, \nabla v)\|_{L^2(D^+)} + \|(\nabla\phi, \nabla\psi)\|_{W^{1,p}(D^+)} \\ &\quad + \|(\tau, \kappa)\|_{L^p(D^+)}), \end{aligned}$$

where we use the Sobolev inequality

$$\|v\|_{L^{\frac{2p}{2-p}}(D^+)} \leq C(p)\|\nabla v\|_{L^2(D^+)}.$$

Taking $\epsilon_2 > 0$ sufficiently small, we have

$$\begin{aligned} \|u\|_{W^{2,p}(D^+_{3/4})} + \|v\|_{W^{2,p}(D^+_{3/4})} &\leq \|\eta u\|_{W^{2,p}(D^+)} + \|\eta v\|_{W^{2,p}(D^+)} \\ &\leq C(\|(\nabla u, \nabla v)\|_{L^2(D^+)} + \|(\nabla\phi, \nabla\psi)\|_{W^{1,p}(D^+)} + \|(\tau, \kappa)\|_{L^p(D^+)}). \end{aligned}$$

Thus we have proved the lemma for the case $1 < p < 2$.

If $p = 2$, one can derive the above estimate for $p = \frac{4}{3}$ at first. Such an estimate implies that ∇u and ∇v are bounded in $L^4(D^+_{3/4})$. Then one can apply the $W^{2,2}$ -boundary estimate to the equation and get the conclusion of the lemma with $p = 2$. \square

For an approximate Lorentzian harmonic map, we can prove the following Pohozaev type identity which is useful in the blow-up analysis. This kind of equality was first introduced in [20] for the interior case of harmonic maps and extended in [10, 15, 16] for some boundary cases.

Lemma 2.3 *Let $D \subset \mathbb{R}^2$ be the unit disk and $(u, v) \in W^{2,2}(D, N \times \mathbb{R})$ be an approximate Lorentzian harmonic map with $(\tau, \kappa) \in L^2(D)$, then for any $0 < \rho < \frac{1}{2}$, we have*

$$\begin{aligned} & \rho \int_{\partial D_\rho} \left(|u_r|^2 - \beta(u)|v_r + \omega_i u_r^i|^2 - \frac{1}{2}|\nabla u|^2 + \frac{1}{2}\beta(u)|\nabla v + \omega_i \nabla u^i|^2 \right) ds \\ &= \int_{D_\rho} r u_r \tau dx - \int_{D_\rho} r(v_r + \omega_i u_r^i) \kappa dx. \end{aligned} \tag{2.1}$$

where (r, θ) are polar coordinates in D centered at 0. Since we use the Euclidean metric, we have that the covariant derivative $\nabla_r u$ equals to $\frac{\partial u}{\partial r}$ and we denote them with a unified notation $\frac{\partial u}{\partial r}$ or just u_r for brevity.

Proof Multiplying (1.8) by $r(v_r + \omega_i u_r^i)$ and integrating by parts, we get

$$\begin{aligned} & \int_{D_\rho} r(v_r + \omega_i u_r^i) \kappa dx \\ &= \int_{D_\rho} \operatorname{div}(\beta(u)(\nabla v + \omega_i \nabla u^i)) \cdot r(v_r + \omega_i u_r^i) dx \\ &= \int_{\partial D_\rho} r\beta(u)|v_r + \omega_i u_r^i|^2 ds - \int_{D_\rho} \beta(u)(\nabla v + \omega_i \nabla u^i) \cdot \nabla \left(r(v_r + \omega_i u_r^i) \right) dx \\ &= \int_{\partial D_\rho} r\beta(u)|v_r + \omega_i u_r^i|^2 ds - \int_{D_\rho} \beta(u)|\nabla v + \omega_i \nabla u^i|^2 dx \\ &\quad - \int_{D_\rho} \beta(u)(\nabla v + \omega_i \nabla u^i) \cdot r \left(\frac{\partial}{\partial r}(\nabla v + \omega_i \nabla u^i) - \frac{\partial \omega_i}{\partial r} \nabla u^i + \frac{\partial u_i}{\partial r} \nabla \omega^i \right) dx. \end{aligned}$$

By direct computations, noting that

$$\begin{aligned} & - \int_{D_\rho} \beta(u)(\nabla v + \omega_i \nabla u^i) \cdot r \frac{\partial}{\partial r}(\nabla v + \omega_i \nabla u^i) dx \\ &= -\frac{1}{2} \int_{D_\rho} \beta(u)r \frac{\partial}{\partial r}|\nabla v + \omega_i \nabla u^i|^2 dx \\ &= -\frac{1}{2} \int_{\partial D_\rho} r\beta(u)|\nabla v + \omega_i \nabla u^i|^2 ds + \int_{D_\rho} \beta(u)|\nabla v + \omega_i \nabla u^i|^2 dx \\ &\quad + \frac{1}{2} \int_{D_\rho} r \frac{\partial \beta(u)}{\partial r} |\nabla v + \omega_i \nabla u^i|^2 dx, \end{aligned}$$

we have

$$\begin{aligned} \int_{D_\rho} r(v_r + \omega_i u_r^i) \kappa dx &= \int_{\partial D_\rho} r\beta(u) \left(|v_r + \omega_i u_r^i|^2 - \frac{1}{2}|\nabla v + \omega_i \nabla u^i|^2 \right) ds \\ &\quad + \int_{D_\rho} \beta(u)(\nabla v + \omega_i \nabla u^i) \cdot r \left(\frac{\partial \omega_i}{\partial r} \nabla u^i - \frac{\partial u_i}{\partial r} \nabla \omega^i \right) dx \\ &\quad + \frac{1}{2} \int_{D_\rho} r \frac{\partial \beta(u)}{\partial r} |\nabla v + \omega_i \nabla u^i|^2 dx. \end{aligned} \tag{2.2}$$

Similarly, multiplying (1.7) by ru_r and integrating by parts, we get

$$\begin{aligned}
 \int_{D_\rho} ru_r \tau dx &= \int_{D_\rho} (\Delta u - H^\top) ru_r dx \\
 &= \int_{\partial D_\rho} r |u_r|^2 ds - \int_{D_\rho} \nabla u \cdot \nabla (ru_r) dx - \int_{D_\rho} H \cdot ru_r dx \\
 &= \int_{\partial D_\rho} r \left(|u_r|^2 - \frac{1}{2} |\nabla u|^2 \right) - \int_{D_\rho} H \cdot ru_r dx.
 \end{aligned} \tag{2.3}$$

Noting that

$$\begin{aligned}
 H \cdot ru_r &= r H^j u_r^j \\
 &= r \left(\beta (\nabla v + \omega_i \nabla u^i) \cdot \nabla u^k \left(\frac{\partial \omega_j}{\partial y^k} - \frac{\partial \omega_k}{\partial y^j} \right) - \frac{1}{2} \frac{\partial \beta}{\partial y^j} |\nabla v + \omega_i \nabla u^i|^2 \right) u_r^j \\
 &= r \beta (\nabla v + \omega_i \nabla u^i) \left(u_r^j \nabla \omega_j - \frac{\partial \omega_j}{\partial r} \nabla u^j \right) - \frac{1}{2} r \frac{\partial \beta(u)}{\partial r} |\nabla v + \omega_i \nabla u^i|^2
 \end{aligned} \tag{2.4}$$

and combining (2.4) with (2.2) and (2.3), we obtain the conclusion of the lemma.

By Hölder’s inequality and integrating (2.1) about ρ from r_0 to $2r_0$, we get

Corollary 2.4 *For (u, v) in Lemma 2.3, if $\|(\nabla u, \nabla v)\|_{L^2(D)} + \|(\tau, \kappa)\|_{L^2(D)} \leq \Lambda$, then for any $0 < r_0 < \frac{1}{4}$, we have*

$$\int_{D_{2r_0} \setminus D_{r_0}} \left(|u_r|^2 - \beta(u) |v_r + \omega_i u_r^i|^2 - \frac{1}{2} |\nabla u|^2 + \frac{1}{2} \beta(u) |\nabla v + \omega_i \nabla u^i|^2 \right) dx \leq Cr_0,$$

where $C = C(\lambda_1, \lambda_2, \Lambda, N) > 0$ is a constant.

Denote $\tilde{u} = u - \phi$ and $\tilde{v} = v - \psi$. For the boundary case, we have

Lemma 2.5 *Let $D^+ \subset \mathbb{R}^2$ be the upper unit disk and $(u, v) \in W^{2,2}(D^+, N \times \mathbb{R})$ be an approximate Lorentzian harmonic map with Dirichlet boundary data $(u, v)|_{\partial^0 D^+} = (\phi, \psi) \in C^{2+\alpha}(D)$ and $(\tau, \kappa) \in L^2(D^+)$, then for any $0 < \rho < \frac{1}{2}$, we have*

$$\begin{aligned}
 &\rho \int_{\partial^+ D_\rho^+} (|u_r|^2 - \beta(u) |v_r + \omega_i u_r^i|^2 - \frac{1}{2} |\nabla u|^2 + \frac{1}{2} \beta(u) |\nabla v + \omega_i \nabla u^i|^2) ds \\
 &= \int_{D_\rho^+} r \tilde{u}_r \tau dx - \int_{D_\rho^+} r (\tilde{v}_r + \omega_i \tilde{u}_r^i) \kappa dx + \int_{\partial^+ D_\rho^+} r \frac{\partial u}{\partial r} \frac{\partial \phi}{\partial r} ds - \int_{D_\rho^+} \nabla u \nabla (r \phi_r) dx \\
 &\quad + \int_{D_\rho^+} (A(u)(\nabla u, \nabla u) - H^\top) \cdot r \phi_r dx - \int_{\partial^+ D_\rho^+} r \beta(u) (v_r + \omega_i u_r^i) (\psi_r + \omega_i \phi_r^i) ds \\
 &\quad + \int_{D_\rho^+} \beta(u) (\nabla v + \omega_i \nabla u^i) \cdot (\nabla \psi + \omega_i \nabla \phi^i) dx \\
 &\quad + \int_{D_\rho^+} r \beta(u) (\nabla v + \omega_i \nabla u^i) \cdot \nabla (\psi_r + \omega_i \phi_r^i) dx
 \end{aligned} \tag{2.5}$$

where (r, θ) are polar coordinates in D centered at 0.

Proof The proof is similar to the proof of Lemma 2.3.

Multiplying (1.8) by $r(\tilde{v}_r + \omega_i \tilde{u}_r^i)$ and integrating by parts, we get

$$\begin{aligned} \int_{D_\rho^+} r(\tilde{v}_r + \omega_i \tilde{u}_r^i) \kappa dx &= \int_{D_\rho^+} \operatorname{div}(\beta(u)(\nabla v + \omega_i \nabla u^i)) \cdot r(\tilde{v}_r + \omega_i \tilde{u}_r^i) dx \\ &= \int_{\partial^+ D_\rho^+} \beta(u)(v_r + \omega_i u_r^i) \cdot r(\tilde{v}_r + \omega_i \tilde{u}_r^i) ds \\ &\quad - \int_{D_\rho^+} \beta(u)(\nabla v + \omega_i \nabla u^i) \cdot \nabla(r(\tilde{v}_r + \omega_i \tilde{u}_r^i)) dx \\ &= \int_{\partial^+ D_\rho^+} \beta(u)(v_r + \omega_i u_r^i) \cdot r(\tilde{v}_r + \omega_i \tilde{u}_r^i) ds \\ &\quad - \int_{D_\rho^+} \beta(u)(\nabla v + \omega_i \nabla u^i) \cdot (\nabla \tilde{v} + \omega_i \nabla \tilde{u}^i) dx \\ &\quad - \int_{D_\rho^+} r\beta(u)(\nabla v + \omega_i \nabla u^i) \cdot \nabla(\tilde{v}_r + \omega_i \tilde{u}_r^i) dx = \text{II} + \text{III} + \text{III}. \end{aligned}$$

By direct computations, we have

$$\begin{aligned} \text{II} &= \int_{\partial^+ D_\rho^+} r\beta(u)|v_r + \omega_i u_r^i|^2 ds - \int_{\partial^+ D_\rho^+} r\beta(u)(v_r + \omega_i u_r^i)(\psi_r + \omega_i \phi_r^i) ds, \\ \text{III} &= - \int_{D_\rho^+} \beta(u)|\nabla v + \omega_i \nabla u^i|^2 dx + \int_{D_\rho^+} \beta(u)(\nabla v + \omega_i \nabla u^i) \cdot (\nabla \psi + \omega_i \nabla \phi^i) dx, \end{aligned}$$

and

$$\begin{aligned} \text{III} &= -\frac{1}{2} \int_{D_\rho^+} r\beta(u) \frac{\partial}{\partial r} |\nabla v + \omega_i \nabla u^i|^2 dx + \int_{D_\rho^+} r\beta(u)(\nabla v + \omega_i \nabla u^i) \cdot \nabla(\psi_r + \omega_i \phi_r^i) dx \\ &\quad + \int_{D_\rho^+} r\beta(u)(\nabla v + \omega_i \nabla u^i) \cdot \left(\frac{\partial \omega_i}{\partial r} \nabla u^i - \nabla \omega_i u_r^i \right) dx. \end{aligned}$$

Noting that

$$\begin{aligned} &-\frac{1}{2} \int_{D_\rho^+} \beta(u)r \frac{\partial}{\partial r} |\nabla v + \omega_i \nabla u^i|^2 dx \\ &= -\frac{1}{2} \int_{\partial^+ D_\rho^+} r\beta(u)|\nabla v + \omega_i \nabla u^i|^2 ds + \int_{D_\rho^+} \beta(u)|\nabla v + \omega_i \nabla u^i|^2 dx \\ &\quad + \frac{1}{2} \int_{D_\rho^+} r \frac{\partial \beta(u)}{\partial r} |\nabla v + \omega_i \nabla u^i|^2 dx, \end{aligned}$$

we have

$$\begin{aligned}
 \int_{D_\rho^+} r(\tilde{v}_r + \omega_i \tilde{u}_r^i) \kappa dx &= \int_{\partial^+ D_\rho^+} r\beta(u) \left(|v_r + \omega_i u_r^i|^2 - \frac{1}{2} |\nabla v + \omega_i \nabla u^i|^2 \right) ds \\
 &\quad + \int_{D_\rho^+} \beta(u) (\nabla v + \omega_i \nabla u^i) \cdot r \left(\frac{\partial \omega_i}{\partial r} \nabla u^i - \frac{\partial u_i}{\partial r} \nabla \omega^i \right) dx \\
 &\quad + \frac{1}{2} \int_{D_\rho^+} r \frac{\partial \beta(u)}{\partial r} |\nabla v + \omega_i \nabla u^i|^2 dx \\
 &\quad - \int_{\partial^+ D_\rho^+} r\beta(u) (v_r + \omega_i u_r^i) (\psi_r + \omega_i \phi_r^i) ds \\
 &\quad + \int_{D_\rho^+} \beta(u) (\nabla v + \omega_i \nabla u^i) \cdot (\nabla \psi + \omega_i \nabla \phi^i) dx \\
 &\quad + \int_{D_\rho^+} r\beta(u) (\nabla v + \omega_i \nabla u^i) \cdot \nabla (\psi_r + \omega_i \phi_r^i) dx. \tag{2.6}
 \end{aligned}$$

Similarly, multiplying (1.7) by $r\tilde{u}_r$ and integrating by parts, we get

$$\begin{aligned}
 \int_{D_\rho^+} r\tilde{u}_r \tau dx &= \int_{D_\rho^+} (\Delta u + A(u)(\nabla u, \nabla u) - H^\top) \cdot r\tilde{u}_r dx \\
 &= \int_{\partial^+ D_\rho^+} r(|u_r|^2 - \frac{1}{2} |\nabla u|^2) ds - \int_{\partial^+ D_\rho^+} r \frac{\partial u}{\partial r} \frac{\partial \phi}{\partial r} + \int_{D_\rho^+} \nabla u \nabla (r\phi_r) dx \\
 &\quad - \int_{D_\rho^+} (A(u)(\nabla u, \nabla u) - H^\top) \cdot r\phi_r dx - \int_{D_\rho^+} H \cdot ru_r dx. \tag{2.7}
 \end{aligned}$$

Combining (2.4) with (2.6) and (2.7), we obtain the conclusion of the lemma. □

Corollary 2.6 *For (u, v) in Lemma 2.5, if $\|(\nabla u, \nabla v)\|_{L^2(D^+)} + \|(\tau, \kappa)\|_{L^2(D^+)} \leq \Lambda$, then for any $0 < r_0 < \frac{1}{4}$, we have*

$$\int_{D_{2r_0}^+ \setminus D_{r_0}^+} (|u_r|^2 - \beta(u)|v_r + \omega_i u_r^i|^2 - \frac{1}{2} |\nabla u|^2 + \frac{1}{2} \beta(u) |\nabla v + \omega_i \nabla u^i|^2) dx \leq Cr_0,$$

where $C = C(\lambda_1, \lambda_2, \Lambda, N) > 0$ is a constant.

Proof By Hölder’s inequality, it is easy to find that the right hand side of (2.5) is bounded by

$$C\rho \int_{\partial^+ D_\rho^+} (|\nabla u| + |\nabla v|) + C\rho,$$

where $C = C(\lambda_1, \lambda_2, N, \|\phi\|_{C^2}, \|\psi\|_{C^2})$. Then the conclusion of the corollary follows by an integration about ρ from r_0 to $2r_0$. □

Similar as for harmonic maps into a Riemannian manifold, there is also an energy gap for a nontrivial Lorentzian harmonic map.

Theorem 2.7 (Gap phenomenon) *Suppose either $(u, v) : \mathbb{S}^2 \rightarrow N \times \mathbb{R}$ is a smooth Lorentzian harmonic map or $(u, v) : \mathbb{R}_+^2 \rightarrow N \times \mathbb{R}$ is a smooth Lorentzian harmonic map with Dirichlet boundary condition:*

$$(u, v)|_{\partial \mathbb{R}_+^2} = \text{constant},$$

then there exists a constant $\epsilon_0 > 0$ depending on $(N \times \mathbb{R}, g)$, such that if either

$$E(u, v; \mathbb{S}^2) \leq \epsilon_0 \text{ or } E(u, v; \mathbb{R}_+^2) \leq \epsilon_0,$$

then (u, v) is a constant map. Here \mathbb{S}^2 denotes the unit sphere in \mathbb{R}^3 .

Proof One can find the proof of the theorem in [9] for the case $(u, v) : \mathbb{S}^2 \rightarrow N \times \mathbb{R}$.

By Eqs. (1.4) and (1.5), we have

$$|\Delta u| + |\Delta v| \leq C(|\nabla u|^2 + |\nabla v|^2).$$

The standard elliptic theory tells us that

$$\begin{aligned} \|\nabla u\|_{W^{1, \frac{4}{3}}} + \|\nabla v\|_{W^{1, \frac{4}{3}}} &\leq C(\|\Delta u\|_{L^{\frac{4}{3}}} + \|\Delta v\|_{L^{\frac{4}{3}}}) \\ &\leq CE^{\frac{1}{2}}(u, v; \mathbb{S}^2)(\|\nabla u\|_{L^4} + \|\nabla v\|_{L^4}) \\ &\leq C\sqrt{\epsilon_0}(\|\nabla u\|_{W^{1, \frac{4}{3}}} + \|\nabla v\|_{W^{1, \frac{4}{3}}}). \end{aligned}$$

It is easy to get that, if ϵ_0 is small enough, (u, v) must be a constant map.

If $(u, v) : \mathbb{R}_+^2 \rightarrow N \times \mathbb{R}$ is a smooth Lorentzian harmonic map with constant Dirichlet boundary condition, choosing $\epsilon_0 \leq \epsilon_2$ where ϵ_2 is the positive constant in Lemma 2.2, then by Lemma 2.2 (taking $(\phi, \psi) = \text{constant}$, $(\tau, \kappa) = 0$ and any constant $p > 2$) and Sobolev embedding, for any $R > 0$, we have

$$R\|\nabla u\|_{L^\infty(D_{R/2}^+)} + R\|\nabla v\|_{L^\infty(D_{R/2}^+)} \leq CE^{\frac{1}{2}}(u, v; D_R^+) \leq C\epsilon_0^{\frac{1}{2}}.$$

Sending R to infinity yields that (u, v) must be a constant map. □

It is necessary for the singularities to be removable during the blow-up process. Removability of singularities for a Lorentzian harmonic map (i.e. $\tau = \kappa = 0$) is proved in [9]. By assuming additionally that $\omega \equiv 0$, for an approximate Lorentzian harmonic map (i.e. $(\tau, \kappa) \neq 0$) with singularities either in the interior or on the boundary, we can also remove them.

Theorem 2.8 *Suppose $(u, v) \in W_{loc}^{2,2}(D \setminus \{0\})$ is an approximate Lorentzian harmonic map from the punctured disk $D \setminus \{0\}$ to $(N \times \mathbb{R}, g_N - \beta d^2\theta)$. If $E(u, v; D) < \infty$ and $(\tau, \kappa) \in L^2(D)$, then (u, v) can be extended to the whole disk D in $W^{2,2}(D)$.*

For an approximate Lorentzian harmonic map $(u, v) \in W_{loc}^{2,2}(D^+ \setminus \{0\})$ which is from $D^+ \setminus \{0\}$ to $(N \times \mathbb{R}, g_N - \beta d^2\theta)$ with boundary data $(u, v)|_{\partial^0 D^+} = (\phi, \psi)$. If $E(u, v; D^+) < \infty$ and $(\tau, \kappa) \in L^2(D^+)$, then (u, v) can also be extended to D^+ in $W^{2,2}(D^+)$.

Proof We prove the theorem for the boundary case and the interior case can be proved similarly.

On the one hand, it is easy to see that (u, v) is a weak solution of (1.7) and (1.8). By Theorem 1.2 in [30] which is developed from the regularity theory for critical elliptic systems with an anti-symmetric structure in [25, 26, 28, 29, 32], we know that $v \in W^{2,p}(D_\rho^+(0))$ for some $\rho > 0$ and any $1 < p < 2$. In fact, the anti-symmetric term in the equation for v equals to zero. This implies that $\nabla v \in L^4(D^+)$.

On the other hand, since the Eq. (1.7) can be written as an elliptic system with an anti-symmetric potential ([25])

$$\Delta u = \Omega \cdot \nabla u + f$$

with $\Omega \in L^2(D^+, so(n) \otimes \mathbb{R}^2)$ and $f \in L^2(D^+)$, using Theorem 1.2 in [30] again, we have $u \in W^{2,p}(D_\rho^+(0))$ for some $\rho > 0$ and any $1 < p < 2$. Then the higher regularity can be derived by a standard bootstrap argument. \square

3 Energy identity and analysis on the neck

In this section, we shall study the behavior at blow-up points both in the interior and on the boundary for an approximate Lorentzian harmonic map sequence $\{(u_n, v_n)\}$. To this end, we first define the blow-up set and show that the blow-up points for such a sequence are finite in number. Throughout this section, we suppose that there exists a constant $\Lambda > 0$ such that the sequence satisfies

$$\|(\nabla u_n, \nabla v_n)\|_{L^2(D_1(0))} + \|(\tau_n, \kappa_n)\|_{L^2(D_1(0))} \leq \Lambda. \tag{3.1}$$

Definition 3.1 For an approximate Lorentzian harmonic map sequence $\{(u_n, v_n)\}$, define

$$S_1 := \bigcap_{r>0} \left\{ x \in M \mid \liminf_{n \rightarrow \infty} \int_{D_r(x)} (|\nabla u_n|^2 + |\nabla v_n|^2) dv_h \geq \epsilon_1 \right\},$$

and

$$S_2 := \bigcap_{r>0} \left\{ x \in \partial M \mid \liminf_{n \rightarrow \infty} \int_{D_r^+(x)} (|\nabla u_n|^2 + |\nabla v_n|^2) dv_h \geq \epsilon_2 \right\},$$

where ϵ_1 and ϵ_2 are constants in Lemmas 2.1 and 2.2. The blow-up set of $\{(u_n, v_n)\}$ is defined to be $S := S_1 \cup S_2$.

Lemma 3.1 For an approximate harmonic map sequence satisfying (3.1), the blow-up set S is a finite set.

Proof By (3.1), we can take a subsequence and still denote it by $\{(u_n, v_n)\}$, such that $\{(u_n, v_n)\}$ converges weakly in $W^{1,2}(M)$ to a limit map $(u, v) : M \rightarrow (N \times \mathbb{R}, g)$. If for any point $x \in M$,

$$\lim_{r \rightarrow 0} \limsup_{n \rightarrow \infty} \int_{D_r(x)} |\nabla u_n|^2 + |\nabla v_n|^2 < \epsilon_1, \tag{3.2}$$

Lemma 2.1 tells that the convergence is strong in $W^{1,2}(M)$. Obviously in this case, S_1 is empty. Otherwise, if there exists a point $p_1 \in M$ such that

$$\lim_{r \rightarrow 0} \limsup_{n \rightarrow \infty} \int_{D_r(p_1)} |\nabla u_n|^2 + |\nabla v_n|^2 \geq \epsilon_1, \tag{3.3}$$

By taking a subsequence, we can assume that

$$\lim_{r \rightarrow 0} \lim_{n \rightarrow \infty} \int_{D_r(p_1)} |\nabla u_n|^2 + |\nabla v_n|^2 \geq \epsilon_1.$$

If (3.2) holds for any point $x \in M \setminus \{p_1\}$, we get that $S_1 = \{p_1\}$. Otherwise, we can find a point p_2 where the energy concentration (3.3) happens. Since the energy of the sequence is bounded, this process must stop after finite steps.

For points on the boundary of M , we can proceed similarly and finally, we get a subsequence $\{(u_n, v_n)\}$ which converges strongly to some (u, v) in $W_{loc}^{1,2}(M \setminus S)$, where $S = S_1 \cup S_2 = \{p_1, p_2, \dots, p_l\}$ is a finite set. \square

We consider the case that the blow-up points are in the interior first. Since the blow-up set \mathcal{S}_1 is finite, we can find small geodesic disks D_{δ_i} (by conformal invariance, we can assume that they are flat disks) for each blow-up point p_i such that $D_{\delta_i} \cap D_{\delta_j} = \emptyset$ for $i \neq j$, $i, j = 1, 2, \dots, I$, and on $M \setminus \cup_{i=1}^I D_{\delta_i}$, (u_n, v_n) converges strongly to a limit map (u, v) . Without loss of generality, we discuss the case that there is only one blow-up point $0 \in D_1(0)$ in \mathcal{S}_1 and the sequence $\{(u_n, v_n)\}$ satisfies that there is some (u, v) such that

$$(u_n, v_n) \rightarrow (u, v) \text{ weakly in } W_{loc}^{2,2}(D_1(0) \setminus \{0\}) \text{ as } n \rightarrow \infty. \tag{3.4}$$

Lemma 3.2 *Let $\{(u_n, v_n)\} \in W^{2,2}(D_1(0), N \times \mathbb{R})$ be a sequence of approximate Lorentzian harmonic maps satisfying (3.1) and (3.4). Up to a subsequence which is still denoted by $\{(u_n, v_n)\}$, there exist a positive integer L and nontrivial Lorentzian harmonic spheres $(\sigma^i, \xi^i) : \mathbb{R}^2 \cup \{\infty\} \rightarrow N \times \mathbb{R}$, $i = 1, \dots, L$ satisfying*

$$\lim_{n \rightarrow \infty} E_g(u_n, v_n; D_1(0)) = E_g(u, v; D_1(0)) + \sum_{i=1}^L E_g(\sigma^i, \xi^i). \tag{3.5}$$

Proof According to the standard induction argument in [5], we can assume that there is only one bubble at the singular point $0 \in D_1(0)$. To prove (3.5) is equivalent to prove that there exists a Lorentzian harmonic sphere (σ, ξ) such that

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} E_g(u_n, v_n, D_\delta) = E_g(\sigma, \xi). \tag{3.6}$$

By the standard argument of blow-up analysis, for any n , there exist sequences $x_n \rightarrow 0$ and $r_n \rightarrow 0$ such that

$$E(u_n, v_n; D_{r_n/2}(x_n)) = \sup_{\substack{x \in D_\delta, r \leq r_n \\ D_r(x) \subset D_\delta}} E(u_n, v_n; D_r(x)) = \frac{\epsilon_1}{8}. \tag{3.7}$$

Without loss of generality, we may assume that $x_n = 0$ and denote $\tilde{u}_n = u_n(r_n x)$, $\tilde{v}_n = v_n(r_n x)$. Then we have

$$E(\tilde{u}_n, \tilde{v}_n; D_{1/2}) = E(u_n, v_n; D_{r_n/2}) = \frac{\epsilon_1}{8} < \epsilon_1 \tag{3.8}$$

and

$$E(\tilde{u}_n, \tilde{v}_n; D_R) = E(u_n, v_n; D_{r_n R}) < \Lambda.$$

By (3.7), we can apply Lemma 2.1 on D_R for $\{(\tilde{u}_n, \tilde{v}_n)\}$ and get that $\{(\tilde{u}_n, \tilde{v}_n)\}$ converges strongly to some Lorentzian harmonic map (σ, ξ) in $W^{1,2}(D_R, N \times \mathbb{R})$ for any $R \geq 1$. By stereographic projection and the removable singularity theorem [9], we get a nonconstant harmonic sphere (σ, ξ) . Thus we get the first bubble at the blow-up point and to prove (3.6) is equivalent to prove that

$$\lim_{R \rightarrow \infty} \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} E_g(u_n, v_n; D_\delta(0) \setminus D_{r_n R}(0)) = 0. \tag{3.9}$$

Since we assume that there is only one bubble, we have that, for any $\epsilon > 0$, there holds that

$$E(u_n, v_n; D_{4\rho} \setminus D_{\frac{\rho}{2}}) \leq \epsilon^2 \text{ for } \rho \in [r_n R, \frac{\delta}{2}] \tag{3.10}$$

as $n \rightarrow \infty$, $R \rightarrow \infty$ and $\delta \rightarrow 0$. Otherwise, we will get a second bubble and this is a contradiction to the assumption that $L = 1$. One can refer to [5, 15, 31] for details of this kind

of arguments. Then by Lemma 2.1 and a standard scaling argument, for any $\rho \in [r_n R, \frac{\delta}{2}]$, we have

$$\begin{aligned} & \|u_n\|_{osc(D_{2\rho} \setminus D_\rho)} + \|v_n\|_{osc(D_{2\rho} \setminus D_\rho)} \\ & \leq CE^{\frac{1}{2}}(u_n, v_n; D_{4\rho} \setminus D_{\frac{\rho}{2}}) + C\rho \|(\tau_n, \kappa_n)\|_{L^2(D_{4\rho} \setminus D_{\frac{\rho}{2}})}. \end{aligned} \tag{3.11}$$

Define

$$u_n^*(r) := \frac{1}{2\pi} \int_0^{2\pi} u_n(r, \theta) d\theta, \quad v_n^*(r) := \frac{1}{2\pi} \int_0^{2\pi} v_n(r, \theta) d\theta.$$

By (3.11), we know that

$$\begin{aligned} \|u_n - u_n^*\|_{L^\infty(D_\delta \setminus D_{r_n R})} &= \sup_{r_n R \leq t \leq \frac{\delta}{2}} \|u_n - u_n^*\|_{L^\infty(D_{2t} \setminus D_t)} \\ &\leq \sup_{r_n R \leq t \leq \frac{\delta}{2}} \|u_n\|_{osc(D_{2t} \setminus D_t)} \leq C(\epsilon + \delta) \end{aligned} \tag{3.12}$$

and similarly,

$$\|v_n - v_n^*\|_{L^\infty(D_\delta \setminus D_{r_n R})} \leq C(\epsilon + \delta).$$

Then we get by integrating by parts that

$$\begin{aligned} & \int_{D_\delta \setminus D_{r_n R}} -\Delta u_n (u_n - u_n^*) dx \\ &= \int_{D_\delta \setminus D_{r_n R}} \nabla u_n \nabla (u_n - u_n^*) dx - \int_{\partial D_\delta} \frac{\partial u_n}{\partial r} (u_n - u_n^*) + \int_{\partial D_{r_n R}} \frac{\partial u_n}{\partial r} (u_n - u_n^*) \\ &\geq \int_{D_\delta \setminus D_{r_n R}} |\nabla u_n|^2 dx - \int_{D_\delta \setminus D_{r_n R}} \left| \frac{\partial u_n}{\partial r} \right|^2 dx - \int_{\partial D_\delta} \frac{\partial u_n}{\partial r} (u_n - u_n^*) \\ &\quad + \int_{\partial D_{r_n R}} \frac{\partial u_n}{\partial r} (u_n - u_n^*) \\ &= \int_{D_\delta \setminus D_{r_n R}} |r^{-1} \frac{\partial u_n}{\partial \theta}|^2 dx - \int_{\partial D_\delta} \frac{\partial u_n}{\partial r} (u_n - u_n^*) + \int_{\partial D_{r_n R}} \frac{\partial u_n}{\partial r} (u_n - u_n^*). \end{aligned} \tag{3.13}$$

Since (u_n, v_n) is an approximate harmonic map, we have

$$|\Delta u_n| + |\Delta v_n| \leq C(\lambda_1, \lambda_2, N)(|\nabla u_n|^2 + |\nabla v_n|^2). \tag{3.14}$$

Then we get from (3.12), (3.13) and (3.14) that

$$\begin{aligned} & \int_{D_\delta \setminus D_{r_n R}} |r^{-1} \frac{\partial u_n}{\partial \theta}|^2 dx \\ & \leq \int_{D_\delta \setminus D_{r_n R}} -\Delta u_n (u_n - u_n^*) dx + \int_{\partial D_\delta} \frac{\partial u_n}{\partial r} (u_n - u_n^*) - \int_{\partial D_{r_n R}} \frac{\partial u_n}{\partial r} (u_n - u_n^*) \\ & \leq C(\epsilon + \delta) \left(\int_{D_\delta \setminus D_{r_n R}} (|\nabla u_n|^2 + |\nabla v_n|^2) dx + \int_{\partial D_\delta} \left| \frac{\partial u_n}{\partial r} \right| + \int_{\partial D_{r_n R}} \left| \frac{\partial u_n}{\partial r} \right| \right). \end{aligned} \tag{3.15}$$

By Lemma 2.1 and the trace theory, we obtain for the boundary term in (3.15) that

$$\begin{aligned} \int_{\partial D_\delta} \left| \frac{\partial u_n}{\partial r} \right| &\leq C(\|\nabla u_n\|_{L^2(D_{\frac{3}{2}\delta} \setminus D_\delta)} + \delta \|\nabla^2 u_n\|_{L^2(D_{\frac{3}{2}\delta} \setminus D_\delta)}) \\ &\leq C(E^{\frac{1}{2}}(u_n, v_n; D_{2\delta} \setminus D_{\frac{\delta}{2}}) + \delta \|(\tau_n, \kappa_n)\|_{L^2(D_{2\delta} \setminus D_{\frac{\delta}{2}})}) \\ &\leq C(\epsilon + \delta). \end{aligned}$$

Similarly,

$$\int_{\partial D_{r_n R}} \left| \frac{\partial u_n}{\partial r} \right| \leq C(\epsilon + \delta).$$

Combining these, we have

$$\int_{D_\delta \setminus D_{r_n R}} |r^{-1} \frac{\partial u_n}{\partial \theta}|^2 dx \leq C(\epsilon + \delta). \tag{3.16}$$

Similarly, we can obtain that

$$\int_{D_\delta \setminus D_{r_n R}} |r^{-1} \frac{\partial v_n}{\partial \theta}|^2 dx \leq C(\epsilon + \delta).$$

Without loss of generality, we may assume $\delta = 2^{m_n} r_n R$, where m_n is a positive integer which tends to ∞ as $n \rightarrow \infty$. By Corollary 2.4, for $i = 0, 1, \dots, m_n - 1$, we have

$$\begin{aligned} &\int_{D_{2^{i+1}r_n R} \setminus D_{2^i r_n R}} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \beta(u_n) \left| \frac{\partial v_n}{\partial r} \right| + \omega_j \left| \frac{\partial u_n^j}{\partial r} \right|^2 \right) dx \\ &\leq C \left(\int_{D_{2^{i+1}r_n R} \setminus D_{2^i r_n R}} |r^{-1} \frac{\partial u_n}{\partial \theta}|^2 dx + \int_{D_{2^{i+1}r_n R} \setminus D_{2^i r_n R}} |r^{-1} \frac{\partial v_n}{\partial \theta}|^2 dx + 2^i r_n R \right). \end{aligned}$$

Since

$$\sum_{i=0}^{m_n-1} 2^i r_n R = 2^{m_n} r_n R = \delta,$$

we get

$$\begin{aligned} &\int_{D_\delta \setminus D_{r_n R}} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \beta(u_n) \left| \frac{\partial v_n}{\partial r} \right| + \omega_j \left| \frac{\partial u_n^j}{\partial r} \right|^2 \right) dx \\ &= \sum_{i=0}^{m_n-1} \int_{D_{2^{i+1}r_n R} \setminus D_{2^i r_n R}} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \beta(u_n) \left| \frac{\partial v_n}{\partial r} \right| + \omega_j \left| \frac{\partial u_n^j}{\partial r} \right|^2 \right) dx \leq C(\epsilon + \delta), \end{aligned} \tag{3.17}$$

from which (3.9) follows immediately. □

When the 1-form $\omega \equiv 0$, the behavior of the sequence at the blow-up points is clearer. In fact, we can get identities for the positive energy E instead of for the Lorentzian energy E_g and there is no neck between the limit map and the bubbles. More precisely, we have

Lemma 3.3 *Assume that $\{(u_n, v_n)\}$ is an approximate Lorentzian harmonic map sequence as in Lemma 3.2 and additionally, we assume that $\omega \equiv 0$ and $\|\nabla v_n\|_{L^p} \leq \Lambda$ for some $p > 2$,*

then we have that $\sigma^i : \mathbb{R}^2 \cup \{\infty\} \rightarrow N$ is a nontrivial harmonic sphere, ξ^i is a constant map and (3.5) becomes

$$\lim_{n \rightarrow \infty} E(u_n; D_1(0)) = E(u; D_1(0)) + \sum_{i=1}^L E(\sigma^i), \tag{3.18}$$

$$\lim_{n \rightarrow \infty} E(v_n; D_1(0)) = E(v; D_1(0)). \tag{3.19}$$

Furthermore, The image

$$u(D_1(0)) \cup \bigcup_{i=1}^L \sigma^i(\mathbb{R}^2) \tag{3.20}$$

is a connected set.

Proof Similar to the proof of Lemma 3.2, to prove (3.18) and (3.19) is equivalent to proving

$$\lim_{R \rightarrow \infty} \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} E(u_n, v_n; D_\delta(0) \setminus D_{r_n R}(0)) = 0. \tag{3.21}$$

Since $\|\nabla v_n\|_{L^p(D)} \leq \Lambda$ for some $p > 2$, we get

$$\int_{D_\delta \setminus D_{r_n R}} |\nabla v_n|^2 dx \leq C \delta^{1-\frac{2}{p}} \left(\int_{D_\delta \setminus D_{r_n R}} |\nabla v_n|^p dx \right)^{\frac{2}{p}} \leq C \delta^{1-\frac{2}{p}}. \tag{3.22}$$

Since $\omega \equiv 0$, (3.17) implies that

$$\int_{D_\delta \setminus D_{r_n R}} \left| \frac{\partial u_n}{\partial r} \right|^2 dx \leq C \int_{D_\delta \setminus D_{r_n R}} |\nabla v_n|^2 dx + C(\epsilon + \delta). \tag{3.23}$$

Combining (3.16), (3.22) with (3.23), we can get (3.21).

To prove (3.20) is equivalent to prove

$$\lim_{R \rightarrow \infty} \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \|u_n\|_{osc(D_\delta(0) \setminus D_{r_n R}(0))} = 0. \tag{3.24}$$

To prove (3.24), denote $Q(s) := D_{2^{s_0+s}r_n R} \setminus D_{2^{s_0-s}r_n R}$ and consider

$$f(s) := \int_{Q(s)} |\nabla u_n|^2 dx,$$

where $0 \leq s_0 \leq m_n$ and $0 \leq s \leq \min\{s_0, m_n - s_0\}$. Integrating by parts, we get

$$\begin{aligned} & \int_{Q(s)} -\Delta u_n (u_n - u_n^*) dx \\ &= \int_{Q(s)} \nabla u_n \nabla (u_n - u_n^*) dx - \int_{\partial Q(s)} \frac{\partial u_n}{\partial r} (u_n - u_n^*) \\ &\geq \frac{1}{2} \int_{Q(s)} |\nabla u_n|^2 dx + \int_{Q(s)} \left(\frac{1}{2} |\nabla u_n|^2 - \left| \frac{\partial u_n}{\partial r} \right|^2 \right) dx - \int_{\partial Q(s)} \frac{\partial u_n}{\partial r} (u_n - u_n^*). \end{aligned}$$

By (3.12) and (3.14), we obtain

$$\begin{aligned} & \left(\frac{1}{2} - C(\epsilon + \delta)\right) \int_{Q(s)} |\nabla u_n|^2 dx \\ & \leq \int_{Q(s)} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla u_n|^2 \right) dx + C(\epsilon + \delta) \int_{Q(s)} |\nabla v_n|^2 dx \\ & \quad + \int_{\partial D_{2^{s_0+s} r_n R}} \frac{\partial u_n}{\partial r} (u_n - u_n^*) - \int_{\partial D_{2^{s_0-s} r_n R}} \frac{\partial u_n}{\partial r} (u_n - u_n^*). \end{aligned} \tag{3.25}$$

We deduce from Corollary 2.4 that

$$\begin{aligned} \int_{Q(s)} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla u_n|^2 \right) dx & \leq C \int_{Q(s)} |\nabla v_n|^2 dx + C 2^{s_0+s} r_n R \\ & \leq C (2^{s_0+s} r_n R)^{1-\frac{2}{p}} \left(\int_{Q(s)} |\nabla v_n|^p dx \right)^{\frac{2}{p}} + C 2^{s_0+s} r_n R \\ & \leq C (2^{s_0+s} r_n R)^{1-\frac{2}{p}}. \end{aligned}$$

For the boundary term in (3.25), by Hölder’s inequality and Poincaré’s inequality, we have

$$\begin{aligned} \left| \int_{\partial D_{2^{s_0+s} r_n R}} \frac{\partial u_n}{\partial r} (u_n - u_n^*) \right| & \leq \left(\int_{\partial D_{2^{s_0+s} r_n R}} \left| \frac{\partial u_n}{\partial r} \right|^2 \right)^{\frac{1}{2}} \left(\int_{\partial D_{2^{s_0+s} r_n R}} |u_n - u_n^*|^2 \right)^{\frac{1}{2}} \\ & \leq \left(\int_{\partial D_{2^{s_0+s} r_n R}} \left| \frac{\partial u_n}{\partial r} \right|^2 \right)^{\frac{1}{2}} \left(2^{s_0+s} r_n R \int_{\partial D_{2^{s_0+s} r_n R}} \left| \frac{\partial u_n}{\partial \theta} \right|^2 \right)^{\frac{1}{2}} \\ & \leq C 2^{s_0+s} r_n R \int_{\partial D_{2^{s_0+s} r_n R}} |\nabla u_n|^2. \end{aligned}$$

Similarly, we also have

$$\left| \int_{\partial D_{2^{s_0-s} r_n R}} \frac{\partial u_n}{\partial r} (u_n - u_n^*) \right| \leq C 2^{s_0-s} r_n R \int_{\partial D_{2^{s_0-s} r_n R}} |\nabla u_n|^2.$$

Taking ϵ and δ sufficiently small, we get from (3.25) that

$$f(s) \leq \frac{C}{\ln 2} f'(s) + C (2^{s_0+s} r_n R)^{1-\frac{2}{p}},$$

which implies

$$\left(2^{-\frac{s}{c}} f(s) \right)' \geq -C (2^{s_0} r_n R)^{1-\frac{2}{p}} 2^{(1-\frac{2}{p}-\frac{1}{c})s},$$

where we can take C sufficiently large such that $1 - \frac{2}{p} - \frac{1}{c} > 0$. Integrating from 2 to L , we arrive at

$$f(2) \leq C 2^{-\frac{1}{c}L} f(L) + C (2^{s_0} r_n R)^{1-\frac{2}{p}} 2^{(1-\frac{2}{p}-\frac{1}{c})L}.$$

Let $s_0 = i$ and $L = L_i := \min\{i, m_n - i\}$. Noting that $Q(L_i) \subset D_\delta \setminus D_{r_n R}$, we have

$$\begin{aligned}
 & \int_{D_{2^{i+2}r_n R} \setminus D_{2^{i-2}r_n R}} |\nabla u_n|^2 dx \\
 & \leq CE(u_n, D_\delta \setminus D_{r_n R})2^{-\frac{1}{c}L_i} + C(2^i r_n R)^{1-\frac{2}{p}}2^{\left(1-\frac{2}{p}-\frac{1}{c}\right)L_i} \\
 & \leq CE(u_n, D_\delta \setminus D_{r_n R})2^{-\frac{1}{c}L_i} + C(2^i r_n R)^{1-\frac{2}{p}}2^{\left(1-\frac{2}{p}-\frac{1}{c}\right)(m_n-i)} \\
 & = CE(u_n, D_\delta \setminus D_{r_n R})2^{-\frac{1}{c}L_i} + C(2^{m_n} r_n R)^{1-\frac{2}{p}}2^{-\frac{1}{c}(m_n-i)} \\
 & = CE(u_n, D_\delta \setminus D_{r_n R})2^{-\frac{1}{c}L_i} + C\delta^{1-\frac{2}{p}}2^{\left(-\frac{1}{c}\right)(m_n-i)} \\
 & \leq C\epsilon 2^{-\frac{1}{c}i} + C\delta^{1-\frac{2}{p}}2^{\frac{1}{c}(i-m_n)}, \tag{3.26}
 \end{aligned}$$

where the last inequality follows from the energy identity (3.21). By using Lemma 2.1, now it is easy to deduce (3.24) from (3.22) and the above estimates (3.26) for energy decay. \square

For the case that the blow-up point is on the boundary of the manifold, the behavior is similar to those in Lemmas 3.2 and 3.3. But the analysis is more complicated. More precisely, we consider an approximate Lorentzian harmonic map sequence $\{(u_n, v_n)\} \in W^{2,2}(D_1^+(0), N \times \mathbb{R})$ with the Dirichlet boundary condition

$$(u_n, v_n)|_{\partial^0 D_1^+(0)} = (\varphi, \psi) \in C^{2+\alpha}(\partial^0 D_1^+(0)) \tag{3.27}$$

for some $0 < \alpha < 1$ which satisfies that

$$\|(\nabla u_n, \nabla v_n)\|_{L^2(D_1^+(0))} + \|(\tau_n, \kappa_n)\|_{L^2(D_1^+(0))} \leq \Lambda. \tag{3.28}$$

Without loss of generality, we still suppose that there is only one blow-up point $0 \in D_1^+(0)$ and the sequence $\{(u_n, v_n)\}$ satisfies that there is some (u, v) such that

$$(u_n, v_n) \rightarrow (u, v) \text{ weakly in } W_{loc}^{2,2}(D_1^+(0) \setminus \{0\}) \text{ as } n \rightarrow \infty. \tag{3.29}$$

For such a sequence, we have

Lemma 3.4 *Let $\{(u_n, v_n)\} \in W^{2,2}(D_1^+(0), N \times \mathbb{R})$ be a sequence of approximate Lorentzian harmonic maps satisfying (3.27), (3.28) and (3.29). Up to a subsequence which is still denoted by $\{(u_n, v_n)\}$, we can find a positive integer L , points $x_n^i \in D_1^+(0)$ and $r_n^i > 0$ satisfying $x_n^i \rightarrow 0$ and $r_n^i \rightarrow 0$, $i = 1, \dots, L$ as $n \rightarrow \infty$ and both of the following two cases may appear during the blow-up process.*

- (a) *If $\frac{dist(x_n^i, \partial^0 D_1^+(0))}{r_n^i} \rightarrow a^i < \infty$, there is a nonconstant Lorentzian harmonic map $(\sigma^i, \xi^i) : \mathbb{R}_{a^i}^2 \rightarrow N \times \mathbb{R}$ with a constant boundary condition which is the weak limit of $(u_n(x_n^i + r_n^i x), v_n(x_n^i + r_n^i x))$ in $W_{loc}^{1,2}(\mathbb{R}_{a^i}^{2+})$, where*

$$\mathbb{R}_{a^i}^2 := \{(x^1, x^2) \in \mathbb{R}^2 | x^2 \geq a^i\} \text{ and } \mathbb{R}_{a^i}^{2+} := \{(x^1, x^2) \in \mathbb{R}^2 | x^2 > a^i\};$$

- (b) *If $\frac{dist(x_n^i, \partial^0 D_1^+(0))}{r_n^i} \rightarrow \infty$, there is a nontrivial Lorentzian harmonic sphere $(\sigma^i, \xi^i) : \mathbb{R}^2 \cup \{\infty\} \rightarrow N \times \mathbb{R}$ which is the weak limit of $(u_n(x_n^i + r_n^i x), v_n(x_n^i + r_n^i x))$ in $W_{loc}^{1,2}(\mathbb{R}^2)$.*

Furthermore, for both of the two cases, there holds the energy identity

$$\lim_{n \rightarrow \infty} E_g(u_n, v_n; D_1^+(0)) = E_g(u, v; D_1^+(0)) + \sum_{i=1}^L E_g(\sigma^i, \xi^i). \tag{3.30}$$

Here, L just stands for a nonnegative integer which may differ from the constant in Lemma 3.2.

Proof Similar to what we have done in the proof of Lemma 3.2, for any n , there exist sequences $x_n \rightarrow 0$ and $r_n \rightarrow 0$ such that

$$E(u_n, v_n; D_{r_n}^+(x_n)) = \sup_{\substack{x \in D_\delta^+, r \leq r_n \\ D_r^+(x) \subset D_\delta^+}} E(u_n, v_n; D_r^+(x)) = \frac{1}{8} \min\{\epsilon_1, \epsilon_2\}, \tag{3.31}$$

where $D_{r_n}^+(x_n) := \{(y = (y^1, y^2) \in \mathbb{R}^2 \mid |y - x_n| \leq r_n, y^2 \geq 0)\}$. Denote $d_n = \text{dist}(x_n, \partial^0 D^+)$. We have that either $\limsup_{n \rightarrow \infty} \frac{d_n}{r_n} < \infty$ or $\limsup_{n \rightarrow \infty} \frac{d_n}{r_n} = \infty$. We discuss these two cases respectively.

Case (a) $\limsup_{n \rightarrow \infty} \frac{d_n}{r_n} < \infty$.

By taking a subsequence, we may assume that $\lim_{n \rightarrow \infty} \frac{d_n}{r_n} = a \geq 0$. Denote

$$B_n := \{x \in \mathbb{R}^2 \mid x_n + r_n x \in D^+\}.$$

We have that as $n \rightarrow \infty$,

$$B_n \rightarrow \mathbb{R}_a^2 := \{(x^1, x^2) \mid x^2 \geq -a\}$$

and for any $x \in \{x^2 = -a\}$ on the boundary, $x_n + r_n x \rightarrow 0$.

Define

$$\tilde{u}_n(x) := u_n(x_n + r_n x), \quad \tilde{v}_n(x) := v_n(x_n + r_n x).$$

It is easy to get that $(\tilde{u}_n, \tilde{v}_n) : B_n \rightarrow N \times \mathbb{R}$ is an approximate Lorentzian harmonic map with $(\tilde{\tau}_n, \tilde{\kappa}_n) = r_n^2(\tau_n, \kappa_n)$ and

$$(\tilde{u}_n(x), \tilde{v}_n(x)) = (\varphi(x_n + r_n x), \psi(x_n + r_n x)), \text{ if } x_n + r_n x \in \partial^0 D^+.$$

Lemma 2.2 and (3.31) tell us that for any $D_R(0) \subset \mathbb{R}^2$,

$$\|\tilde{u}_n\|_{W^{2,2}(D_R(0) \cap B_n)} + \|\tilde{v}_n\|_{W^{2,2}(D_R(0) \cap B_n)} \leq C(\lambda_1, \lambda_2, \Lambda, R, N).$$

By a similar argument as in Section 4 of [8], after taking a subsequence of $(\tilde{u}_n, \tilde{v}_n)$ if necessary (still denoted by $(\tilde{u}_n, \tilde{v}_n)$), there is a Lorentzian harmonic map $(\tilde{u}, \tilde{v}) \in W^{1,2}(\mathbb{R}_a^2, N \times \mathbb{R})$ with the constant boundary condition $(\tilde{u}, \tilde{v})|_{\partial \mathbb{R}_a^2} = (\phi(0), \psi(0))$ such that, for any $R > 0$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \|\nabla \tilde{u}_n\|_{L^2(D_R(0) \cap B_n)} &= \|\nabla \tilde{u}\|_{L^2(D_R(0) \cap \mathbb{R}_a^2)}, \\ \lim_{n \rightarrow \infty} \|\nabla \tilde{v}_n\|_{L^2(D_R(0) \cap B_n)} &= \|\nabla \tilde{v}\|_{L^2(D_R(0) \cap \mathbb{R}_a^2)}. \end{aligned}$$

Moreover, since $E(\tilde{u}, \tilde{v}; D_1(0) \cap \mathbb{R}_a^2) = \frac{1}{8} \min\{\epsilon_1, \epsilon_2\}$, (\tilde{u}, \tilde{v}) is a nontrivial Lorentzian harmonic map with constant boundary $(\phi(0), \psi(0))$.

Case (b) $\limsup_{n \rightarrow \infty} \frac{d_n}{r_n} = \infty$.

In this case, $(\tilde{u}_n, \tilde{v}_n)$ lives in B_n which tends to \mathbb{R}^2 as $n \rightarrow \infty$. Moreover, for any $x \in \mathbb{R}^2$, when n is sufficiently large, by (3.31), we have

$$E(\tilde{u}_n, \tilde{v}_n; D_1(x)) \leq \frac{\epsilon_1}{8}.$$

According to Lemma 2.1, there exist a subsequence of $(\tilde{u}_n, \tilde{v}_n)$ which is still denoted by $(\tilde{u}_n, \tilde{v}_n)$ and a Lorentzian harmonic map $(\tilde{u}(x), \tilde{v}(x)) \in W^{1,2}(\mathbb{R}^2, N \times \mathbb{R})$ such that

$$\lim_{n \rightarrow \infty} (\tilde{u}_n(x), \tilde{v}_n(x)) = (\tilde{u}(x), \tilde{v}(x)) \text{ in } W_{loc}^{1,2}(\mathbb{R}^2).$$

By Theorem 2.8, (\tilde{u}, \tilde{v}) can be extended to a Lorentzian harmonic sphere and (3.31) tells us that it is nontrivial.

We call the Lorentzian harmonic map (\tilde{u}, \tilde{v}) obtained in these two cases the first bubble. Without loss of generality, we assume that there is only one bubble at the blow-up point $0 \in D_1^+(0)$. Under this assumption, similar to (3.10), we have that, for any $\epsilon > 0$, there exist constants $\delta > 0$ and $R > 0$ such that

$$E(u_n, v_n; D_{4\rho}^+(x_n) \setminus D_{\frac{\rho}{2}}^+(x_n)) \leq \epsilon^2 \text{ for any } \rho \in [r_n R, \frac{\delta}{2}] \tag{3.32}$$

when n is large enough.

Now to prove the energy identity (3.30) is equivalent to prove

$$\lim_{R \rightarrow \infty} \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} E_g(u_n, v_n; D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n)) = 0. \tag{3.33}$$

We shall prove (3.33) for the two cases respectively.

For case (a) $\lim_{n \rightarrow \infty} \frac{d_n}{r_n} = a < \infty$.

For n and R are sufficiently large, we decompose the neck domain $D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n)$ into three parts which follows the decomposition in [15, 16].

$$\begin{aligned} D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n) &= D_\delta^+(x_n) \setminus D_{\frac{\delta}{2}}^+(x'_n) \cup D_{\frac{\delta}{2}}^+(x'_n) \setminus D_{2r_n R}^+(x'_n) \cup D_{2r_n R}^+(x'_n) \setminus D_{r_n R}^+(x_n) \\ &:= \Omega_1 \cup \Omega_2 \cup \Omega_3. \end{aligned}$$

Here $x'_n \in \partial^0 D^+$ is the projection of x_n , i.e. $d_n = |x_n - x'_n|$.

Since $\lim_{n \rightarrow \infty} \frac{d_n}{r_n} = a$, when n and R are large enough, it is easy to get that

$$\Omega_1 \subset D_\delta^+(x_n) \setminus D_{\frac{\delta}{4}}^+(x_n) \quad \text{and} \quad \Omega_3 \subset D_{4r_n R}^+(x_n) \setminus D_{r_n R}^+(x_n).$$

Moreover, for any $\rho \in [r_n R, \frac{\delta}{2}]$, there holds

$$D_{2\rho}^+(x'_n) \setminus D_\rho^+(x'_n) \subset D_{4\rho}^+(x_n) \setminus D_{\rho/2}^+(x_n).$$

Then we get from (3.32) that

$$E(u_n, v_n; \Omega_1) + E(u_n, v_n; \Omega_3) \leq \epsilon^2 \tag{3.34}$$

and

$$E(u_n, v_n; D_{2\rho}^+(x'_n) \setminus D_\rho^+(x'_n)) \leq \epsilon^2 \text{ for any } \rho \in \left[r_n R, \frac{\delta}{2} \right].$$

By Lemma 2.1, we have

$$\begin{aligned} & \|u_n\|_{osc(D_{2\rho}^+(x'_n)\setminus D_\rho^+(x'_n))} + \|v_n\|_{osc(D_{2\rho}^+(x'_n)\setminus D_\rho^+(x'_n))} \\ & \leq C(\|\nabla u_n, \nabla v_n\|_{L^2(D_{4\rho}^+(x'_n)\setminus D_{\rho/2}^+(x'_n))} + \|(\nabla\phi, \nabla\psi)\|_{L^2(D_{4\rho}^+(x'_n)\setminus D_{\rho/2}^+(x'_n))} \\ & \quad + \rho\|(\nabla^2\varphi, \nabla^2\psi)\|_{L^2(D_{4\rho}^+(x'_n)\setminus D_{\rho/2}^+(x'_n))} + \rho\|(\tau_n, \kappa_n)\|_{L^2(D_{4\rho}^+(x'_n)\setminus D_{\rho/2}^+(x'_n))} \end{aligned} \tag{3.35}$$

for any $\rho \in [r_n R, \frac{\delta}{2}]$.

To estimate the energy concentration in Ω_2 , we define $\mu_n(x) := u_n(x) - \varphi(x)$ for $x \in \Omega_2$ and

$$\widehat{\mu}_n(x) := \begin{cases} \mu_n(x) & x \in \Omega_2, \\ -\mu_n(x') & x \in \widehat{\Omega}_2 \setminus \Omega_2, \end{cases}$$

where $\widehat{\Omega}_2 := D_{\frac{\delta}{2}}(x'_n) \setminus D_{2r_n R}(x'_n)$, $x = (x^1, x^2)$ and $x' = (x^1, -x^2)$. It is easy to get that $\widehat{\mu}_n(x) \in W^{2,2}(\widehat{\Omega}_2)$ and satisfies

$$|\Delta \widehat{\mu}_n(x)| \leq \begin{cases} C(|\nabla u_n(x)|^2 + |\nabla v_n(x)|^2 + |\tau_n(x)| + |\Delta\varphi(x)|, & x \in \Omega_2, \\ C(|\nabla u_n(x')|^2 + |\nabla v_n(x')|^2 + |\tau_n(x')| + |\Delta\varphi(x')|, & x \in \widehat{\Omega}_2 \setminus \Omega_2. \end{cases}$$

Define

$$\widehat{\mu}_n^*(r) := \frac{1}{2\pi} \int_0^{2\pi} \widehat{\mu}_n(r, \theta) d\theta,$$

where (r, θ) is the polar coordinates at x'_n . By (3.35), we have

$$\begin{aligned} \|\widehat{\mu}_n(x) - \widehat{\mu}_n^*(x)\|_{L^\infty(\widehat{\Omega}_2)} & \leq \sup_{r_n R \leq \rho \leq \frac{\delta}{2}} \|\widehat{\mu}_n(x)\|_{osc(D_{2\rho}(x'_n)\setminus D_\rho(x'_n))} \\ & \leq 2 \sup_{r_n R \leq \rho \leq \frac{\delta}{2}} \|\mu_n(x)\|_{osc(D_{2\rho}^+(x'_n)\setminus D_\rho^+(x'_n))} \\ & \leq C(\epsilon + \delta). \end{aligned}$$

Similar to the proof of (3.15), we can obtain

$$\begin{aligned} & \int_{\widehat{\Omega}_2} |r^{-1} \frac{\partial \widehat{\mu}_n}{\partial \theta}|^2 dx \\ & \leq C(\epsilon + \delta) \left(\int_{\widehat{\Omega}_2} |\Delta \widehat{\mu}_n| dx + \int_{\partial D_{\frac{\delta}{2}}(x'_n)} \left| \frac{\partial \widehat{\mu}_n}{\partial r} \right| + \int_{\partial D_{2r_n R}(x'_n)} \left| \frac{\partial \widehat{\mu}_n}{\partial r} \right| \right). \end{aligned} \tag{3.36}$$

By direct computations, one can get that

$$\begin{aligned} \int_{\widehat{\Omega}_2} |r^{-1} \frac{\partial \widehat{\mu}_n}{\partial \theta}|^2 dx & = 2 \int_{\Omega_2} |r^{-1} \frac{\partial \mu_n}{\partial \theta}|^2 dx \\ & = 2 \int_{\Omega_2} |r^{-1} \frac{\partial u_n}{\partial \theta}|^2 dx - 4 \int_{\Omega_2} r^{-2} \frac{\partial u_n}{\partial \theta} \frac{\partial \phi}{\partial \theta} dx + 2 \int_{\Omega_2} |r^{-1} \frac{\partial \phi}{\partial \theta}|^2 dx \\ & \geq 2 \int_{\Omega_2} |r^{-1} \frac{\partial u_n}{\partial \theta}|^2 dx - C\delta \end{aligned} \tag{3.37}$$

and

$$\begin{aligned} \int_{\widehat{\Omega}_2} |\Delta \widehat{\mu}_n| dx &\leq C \int_{\Omega_2} (|\nabla u_n|^2 + |\nabla v_n|^2) dx + \int_{\widehat{\Omega}_2} (|\tau_n| + |\Delta \phi|) dx \\ &\leq C \int_{\Omega_2} (|\nabla \bar{u}_n|^2 + |\nabla v_n|^2) dx + C\delta. \end{aligned} \tag{3.38}$$

For the boundary terms of the right hand side of (3.36), by the trace theory and Lemma 2.2, we have

$$\begin{aligned} \int_{\partial D_{\delta/2}(x'_n)} \left| \frac{\partial \widehat{\mu}_n}{\partial r} \right| &= 2 \int_{\partial^+ D_{\delta/2}(x'_n)} \left| \frac{\partial \widehat{\mu}_n}{\partial r} \right| \leq C \int_{\partial^+ D_{\delta/2}(x'_n)} (|\nabla u_n| + |\nabla \phi|) \\ &\leq C (\|(\nabla u_n, \nabla v_n)\|_{L^2(D_{\delta}^+(x'_n) \setminus D_{\frac{\delta}{4}}^+(x'_n))} + \delta \|(\nabla^2 u_n, \nabla^2 v_n)\|_{L^2(D_{\delta}^+(x'_n) \setminus D_{\frac{\delta}{4}}^+(x'_n))} + \delta) \\ &\leq C (\|(\nabla u_n, \nabla v_n)\|_{L^2(D_{\frac{4}{3}\delta}^+(x_n) \setminus D_{\frac{1}{6}\delta}^+(x_n))} + \|(\nabla \phi, \nabla \psi)\|_{L^2(D_{\frac{4}{3}\delta}^+(x_n) \setminus D_{\frac{1}{6}\delta}^+(x_n))} \\ &\quad + \delta \|(\nabla^2 \phi, \nabla^2 \psi)\|_{L^2(D_{\frac{4}{3}\delta}^+(x_n) \setminus D_{\frac{1}{6}\delta}^+(x_n))} + \delta \|(\tau_n, \kappa_n)\|_{L^2(D_{\frac{4}{3}\delta}^+(x_n) \setminus D_{\frac{1}{6}\delta}^+(x_n))} + \delta) \\ &\leq C(\epsilon + \delta). \end{aligned}$$

Similarly, we have

$$\int_{\partial D_{2r_n R}(x'_n)} \left| \frac{\partial \widehat{\mu}_n}{\partial r} \right| \leq C(\epsilon + \delta).$$

Combining these two estimates with (3.36), (3.37) and (3.38), we get

$$\int_{\Omega_2} |r^{-1} \frac{\partial u_n}{\partial \theta}|^2 dx \leq C(\epsilon + \delta). \tag{3.39}$$

Similarly, we have

$$\int_{\Omega_2} |r^{-1} \frac{\partial v_n}{\partial \theta}|^2 dx \leq C(\epsilon + \delta). \tag{3.40}$$

By Corollary 2.6, we have

$$\begin{aligned} &\int_{D_{2^{i+1}r_n R}^+(x'_n) \setminus D_{2^i r_n R}^+(x'_n)} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \beta(u_n) \left| \frac{\partial v_n}{\partial r} \right| + \omega_j \left| \frac{\partial u_n^j}{\partial r} \right|^2 \right) dx \\ &\leq C \int_{D_{2^{i+1}r_n R}^+(x'_n) \setminus D_{2^i r_n R}^+(x'_n)} |r^{-1} \frac{\partial u_n}{\partial \theta}|^2 dx \\ &\quad + C \int_{D_{2^{i+1}r_n R}^+(x'_n) \setminus D_{2^i r_n R}^+(x'_n)} |r^{-1} \frac{\partial v_n}{\partial \theta}|^2 dx + 2^i r_n R. \end{aligned} \tag{3.41}$$

Thus, we arrive at

$$\begin{aligned} &\int_{\Omega_2} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \beta(u_n) \left| \frac{\partial v_n}{\partial r} \right| + \omega_j \left| \frac{\partial u_n^j}{\partial r} \right|^2 \right) dx \\ &= \sum_{i=0}^{m_n-1} \int_{D_{2^{i+1}r_n R}^+(x'_n) \setminus D_{2^i r_n R}^+(x'_n)} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \beta(u_n) \left| \frac{\partial v_n}{\partial r} \right| + \omega_j \left| \frac{\partial u_n^j}{\partial r} \right|^2 \right) dx \\ &\leq C(\epsilon + \delta). \end{aligned} \tag{3.42}$$

Then (3.39), (3.40), (3.41), (3.42) and (3.34) imply (3.33).

For case (b) $\lim_{n \rightarrow \infty} \frac{d_n}{r_n} = \infty$.

The result for this case can be derived from **case (a)** and Lemma 3.2. In fact, in this case, for n sufficiently large, we decompose the neck domain $D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n)$ as in [15, 16] as follows

$$\begin{aligned} D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n) &= D_\delta^+(x_n) \setminus D_{\frac{\delta}{2}}^+(x'_n) \cup D_{\frac{\delta}{2}}^+(x'_n) \setminus D_{2d_n}^+(x'_n) \\ &\quad \cup D_{2d_n}^+(x'_n) \setminus D_{d_n}^+(x_n) \cup D_{d_n}^+(x_n) \setminus D_{r_n R}^+(x_n) \\ &:= \Omega_1 \cup \Omega_2 \cup \Omega_3 \cup \Omega_4. \end{aligned} \tag{3.43}$$

Since $\lim_{n \rightarrow \infty} d_n = 0$ and $\lim_{n \rightarrow \infty} \frac{d_n}{r_n} = \infty$, when n is large enough, it is easy to get that

$$\Omega_1 \subset D_\delta^+(x_n) \setminus D_{\frac{\delta}{4}}^+(x_n), \quad \text{and} \quad \Omega_3 \subset D_{4d_n}^+(x_n) \setminus D_{d_n}^+(x_n).$$

Moreover, for any $\rho \in [d_n, \frac{\delta}{2}]$, there holds

$$D_{2\rho}^+(x'_n) \setminus D_\rho^+(x'_n) \subset D_{4\rho}^+(x_n) \setminus D_{\rho/2}^+(x_n).$$

By assumption (3.32), we have

$$E(u_n; \Omega_1) + E(u_n; \Omega_3) \leq \epsilon^2$$

and

$$\int_{D_{2\rho}^+(x'_n) \setminus D_\rho^+(x'_n)} |\nabla u_n|^2 dx \leq \epsilon^2 \text{ for any } \rho \in \left(d_n, \frac{\delta}{2}\right).$$

Noting that $\Omega_4 = D_{d_n}^+(x_n) \setminus D_{r_n R}^+(x_n) = D_{d_n}(x_n) \setminus D_{r_n R}(x_n)$, by Lemma 3.2, there holds

$$\lim_{R \rightarrow \infty} \lim_{n \rightarrow 0} E_g(u_n, v_n; D_{d_n}(x_n) \setminus D_{r_n R}(x_n)) = 0.$$

To estimate the energy concentration in Ω_2 , we can use the same arguments as for **case (a)** to get that

$$E_g(u_n, v_n; \Omega_2) \leq C(\epsilon + \delta). \tag{3.44}$$

Thus we finish the proof of the lemma. □

Similar to Lemma 3.3, when $\omega \equiv 0$, we have

Lemma 3.5 *Assume that $\{(u_n, v_n)\}$ is an approximate Lorentzian harmonic map sequence as in Lemma 3.4 and additionally, we assume that $\omega \equiv 0$ and $\|\nabla v_n\|_{L^p(D^+)} \leq \Lambda$ for some $p > 2$, then **case (a)** in Lemma 3.4 will not happen and in **case (b)**, we have that $\sigma^i : \mathbb{R}^2 \cup \{\infty\} \rightarrow N$ is a nontrivial harmonic sphere, ξ^i is a constant map and (3.30) becomes*

$$\lim_{n \rightarrow \infty} E(u_n; D_1^+(0)) = E(u; D_1^+(0)) + \sum_{i=1}^L E(\sigma^i), \tag{3.45}$$

$$\lim_{n \rightarrow \infty} E(v_n; D_1^+(0)) = E(v; D_1^+(0)). \tag{3.46}$$

Furthermore, the image

$$u(D_1^+(0)) \cup \bigcup_{i=1}^L \sigma^i(\mathbb{R}^2) \tag{3.47}$$

is a connected set.

Proof We use the same symbols as in Lemma 3.4. First, let us show that if $\omega \equiv 0$, **Case (a)** will not happen. In fact, since \tilde{v} satisfies

$$div(\beta(\tilde{u})\nabla\tilde{v}) = 0 \text{ in } \mathbb{R}_a^2$$

and $\tilde{v}|_{\partial\mathbb{R}_a^2} \equiv \psi(0)$, \tilde{v} must be a constant map. Thus, \tilde{u} is a harmonic map from \mathbb{R}_a^2 with constant boundary data $\tilde{u}|_{\partial\mathbb{R}_a^2} = \phi(0)$ which implies that \tilde{u} is a constant map [18]. This is a contradiction with $E(\tilde{u}, \tilde{v}; \mathbb{R}_a^2) \geq \frac{1}{8} \min\{\epsilon_1, \epsilon_2\}$.

For **case (b)**, when $\omega \equiv 0$, it is clear that \tilde{v} satisfies the equation

$$div(\beta(\tilde{u})\nabla\tilde{v}) = 0$$

in \mathbb{S}^2 with finite energy $\|\nabla\tilde{v}\|_{L^2(\mathbb{S}^2)} \leq C$ which implies that \tilde{v} must be a constant map. Therefore $\tilde{u} : \mathbb{S}^2 \rightarrow N$ is a nontrivial harmonic sphere.

Now to prove the energy identities (3.45) and (3.46) is equivalent to prove

$$\lim_{R \rightarrow \infty} \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} E(u_n, v_n; D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n)) = 0. \tag{3.48}$$

To prove the no neck result (3.47) is equivalent to prove

$$\lim_{R \rightarrow \infty} \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \|u_n\|_{osc(D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n))} = 0. \tag{3.49}$$

We decompose the neck domain as (3.43). Since $\lim_{n \rightarrow \infty} d_n = 0$ and $\lim_{n \rightarrow \infty} \frac{d_n}{r_n} = \infty$, when n is large enough, it is easy to see that

$$\Omega_1 \subset D_\delta^+(x_n) \setminus D_{\frac{\delta}{4}}^+(x_n) \quad \text{and} \quad \Omega_3 \subset D_{4d_n}^+(x_n) \setminus D_{d_n}^+(x_n).$$

By (3.32), we have

$$E(u_n; \Omega_1) + E(u_n; \Omega_3) \leq \epsilon^2, \tag{3.50}$$

which implies that there is no energy loss on $\Omega_1 \cup \Omega_3$. By Lemmas 2.1, 2.2 and (3.32), we get

$$\begin{aligned} & \|u_n\|_{osc(D_\delta^+(x_n) \setminus D_{\frac{\delta}{4}}^+(x_n))} \\ & \leq \|u_n\|_{osc(D_\delta^+(x_n) \setminus D_{\frac{\delta}{5}}^+(x_n))} \\ & \leq C(\|(\nabla u_n, \nabla v_n)\|_{L^2(D_{\frac{4\delta}{3}}^+(x_n) \setminus D_{\frac{\delta}{6}}^+(x_n))} + \|(\nabla\phi, \nabla\psi)\|_{L^2(D_{\frac{4\delta}{3}}^+(x_n) \setminus D_{\frac{\delta}{6}}^+(x_n))} \\ & \quad + \delta\|(\nabla^2\phi, \nabla^2\psi)\|_{L^2(D_{\frac{4\delta}{3}}^+(x_n) \setminus D_{\frac{\delta}{6}}^+(x_n))} + \delta\|(\tau_n, \kappa_n)\|_{L^2(D_{\frac{4\delta}{3}}^+(x_n) \setminus D_{\frac{\delta}{6}}^+(x_n))}) \\ & \leq C(\epsilon + \delta) \end{aligned} \tag{3.51}$$

and

$$\begin{aligned} & \|u_n\|_{osc(D_{2d_n}^+(x_n) \setminus D_{d_n}^+(x_n))} \\ & \leq \|u_n\|_{osc(D_{3d_n}^+(x_n) \setminus D_{d_n}^+(x_n))} \\ & \leq C(\|(\nabla u_n, \nabla v_n)\|_{L^2(D_{4d_n}^+(x_n) \setminus D_{\frac{d_n}{2}}^+(x_n))} + \|(\nabla\phi, \nabla\psi)\|_{L^2(D_{4d_n}^+(x_n) \setminus D_{\frac{d_n}{2}}^+(x_n))} \\ & \quad + d_n\|(\nabla^2\phi, \nabla^2\psi)\|_{L^2(D_{4d_n}^+(x_n) \setminus D_{\frac{d_n}{2}}^+(x_n))} + d_n\|(\tau_n, \kappa_n)\|_{L^2(D_{4d_n}^+(x_n) \setminus D_{\frac{d_n}{2}}^+(x_n))}) \\ & \leq C(\epsilon + \delta), \end{aligned} \tag{3.52}$$

when n, R are large and δ is small, which implies that there is no neck on $\Omega_1 \cup \Omega_3$.

Moreover, for any $d_n \leq \rho \leq \frac{\delta}{2}$, there holds

$$D_{2\rho}^+(x'_n) \setminus D_{\rho}^+(x'_n) \subset D_{4\rho}^+(x_n) \setminus D_{\rho/2}^+(x_n).$$

when n is big enough, then (3.32) tells us

$$\int_{D_{2\rho}^+(x'_n) \setminus D_{\rho}^+(x'_n)} |\nabla u_n|^2 dx \leq \epsilon^2 \text{ for any } \rho \in (d_n, \frac{\delta}{2}).$$

Combining this with Lemma 2.2, we get

$$\begin{aligned} & \|u_n\|_{osc(D_{2\rho}^+(x'_n) \setminus D_{\rho}^+(x'_n))} \\ & \leq C(\|\nabla u_n, \nabla v_n\|_{L^2(D_{4\rho}^+(x'_n) \setminus D_{\rho/2}^+(x'_n))} + \|(\nabla \phi, \nabla \psi)\|_{L^2(D_{4\rho}^+(x'_n) \setminus D_{\rho/2}^+(x'_n))}) \\ & \quad + \rho\|(\nabla^2 \phi, \nabla^2 \psi)\|_{L^2(D_{4\rho}^+(x'_n) \setminus D_{\rho/2}^+(x'_n))} + \rho\|(\tau_n, \kappa_n)\|_{L^2(D_{4\rho}^+(x'_n) \setminus D_{\rho/2}^+(x'_n))} \end{aligned}$$

for any $\rho \in (d_n, \frac{\delta}{2})$.

Noting that $\Omega_4 = D_{d_n}^+(x_n) \setminus D_{r_n R}^+(x_n) = D_{d_n}(x_n) \setminus D_{r_n R}(x_n)$, the proofs of (3.48) and (3.49) are reduced to the case in Lemma 3.3 and we have

$$\lim_{R \rightarrow \infty} \lim_{n \rightarrow 0} E(u_n, v_n; D_{d_n}(x_n) \setminus D_{r_n R}(x_n)) = 0 \tag{3.53}$$

and

$$\lim_{R \rightarrow \infty} \lim_{n \rightarrow 0} osc(u_n)_{D_{d_n}(x_n) \setminus D_{r_n R}(x_n)} = 0. \tag{3.54}$$

To prove that there is no energy loss on Ω_2 , noting that $\|\nabla v_n\|_{L^p(D^+)} \leq \Lambda$ for some $p > 2$, we get

$$\int_{D_{\delta}^+(x_n) \setminus D_{r_n R}^+(x_n)} |\nabla v_n|^2 dx \leq C\delta^{1-\frac{2}{p}} \left(\int_{D_{\delta}^+(x_n) \setminus D_{r_n R}^+(x_n)} |\nabla v_n|^p dx \right)^{\frac{2}{p}} \leq C\delta^{1-\frac{2}{p}}. \tag{3.55}$$

Combining this with (3.44), we obtain

$$E(u_n, \Omega_2) \leq CE(v_n, \Omega) + C(\epsilon + \delta) \leq C(\epsilon + \delta^{1-\frac{2}{p}}). \tag{3.56}$$

Then, (3.48) follows from (3.50), (3.53), (3.55) and (3.56).

Now we only need to analyze the neck on Ω_2 .

We denote $Q(s) := D_{2^{s_0+s} 2r_n R}^+(x'_n) \setminus D_{2^{s_0-s} 2r_n R}^+(x'_n)$ and $\widehat{Q}(s) := D_{2^{s_0+s} 2r_n R}(x'_n) \setminus D_{2^{s_0-s} 2r_n R}(x'_n)$, where $0 \leq s_0 \leq m_n$ and $0 \leq s \leq \min\{s_0, m_n - s_0\}$. Let

$$f(s) := \int_{Q(s)} |\nabla u_n|^2 dx.$$

Similar to the derivation of (3.15), we can obtain

$$\begin{aligned} & \int_{\widehat{Q}(s)} |\nabla \widehat{\mu}_n|^2 dx - \int_{\widehat{Q}(s)} \left| \frac{\partial \widehat{\mu}_n}{\partial r} \right|^2 dx \\ & \leq C(\epsilon + \delta) \int_{\widehat{Q}(s)} |\Delta \widehat{\mu}_n| dx + \int_{\partial \widehat{Q}(s)} \frac{\partial \widehat{\mu}_n}{\partial r} (\widehat{\mu}_n - \widehat{\mu}_n^*). \end{aligned} \tag{3.57}$$

By direct computations, we obtain

$$\begin{aligned} & \int_{\widehat{Q}(s)} |\nabla \widehat{\mu}_n|^2 dx - \int_{\widehat{Q}(s)} \left| \frac{\partial \widehat{\mu}_n}{\partial r} \right|^2 dx = \int_{Q(s)} |\nabla \mu_n|^2 dx - 2 \int_{Q(s)} \left(\left| \frac{\partial \mu_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla \mu_n|^2 \right) dx \\ & = \int_{Q(s)} |\nabla u_n|^2 dx - 2 \int_{Q(s)} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla u_n|^2 \right) dx + 4 \int_{Q(s)} \left(\frac{\partial u_n}{\partial r} \frac{\partial \phi}{\partial r} - \nabla u_n \nabla \phi \right) dx \\ & \quad + 2 \int_{Q(s)} \left(|\nabla \phi|^2 - \left| \frac{\partial \phi}{\partial r} \right|^2 \right) dx \\ & \geq \int_{Q(s)} |\nabla u_n|^2 dx - 2 \int_{Q(s)} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla u_n|^2 \right) dx - C 2^{s_0+s} r_n R. \end{aligned}$$

It is easy to check that (3.38) still holds on Ω_2 . Combining this with $\|\nabla v_n\|_{L^p(D^+)} \leq C$, we have

$$\begin{aligned} \int_{\widehat{Q}(s)} |\Delta \widehat{\mu}_n| dx & \leq C \int_{Q(s)} (|\nabla u_n|^2 + |\nabla v_n|^2) dx + C 2^{s_0+s} r_n R \\ & \leq C \int_{Q(s)} |\nabla u_n|^2 dx + C (2^{s_0+s} r_n R)^{1-\frac{2}{p}}. \end{aligned}$$

Then (3.57) implies

$$\begin{aligned} & (1 - C(\epsilon + \delta)) \int_{Q(t)} |\nabla u_n|^2 dx \\ & \leq \int_{\partial \widehat{Q}(s)} \frac{\partial \widehat{\mu}_n}{\partial r} (\widehat{\mu}_n - \widehat{\mu}_n^*) + 2 \int_{Q(s)} \left(\left| \frac{\partial u_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla u_n|^2 \right) dx + C (2^{s_0+s} r_n R)^{1-\frac{2}{p}} \\ & \leq \int_{\partial \widehat{Q}(s)} \frac{\partial \widehat{\mu}_n}{\partial r} (\widehat{\mu}_n - \widehat{\mu}_n^*) + C (2^{s_0+s} r_n R)^{1-\frac{2}{p}}, \end{aligned} \tag{3.58}$$

where the last inequality follows from Corollary 2.6 and (3.22).

For the boundary term on the right hand side of (3.58), by Hölder’s inequality and Poincaré’s inequality, we have

$$\begin{aligned} \int_{\partial D_{2^{s_0+s} 2r_n R}(x'_n)} \frac{\partial \widehat{\mu}_n}{\partial r} (\widehat{\mu}_n - \widehat{\mu}_n^*) & \leq \left(\int_{\partial D_{2^{s_0+s} 2r_n R}(x'_n)} \left| \frac{\partial \widehat{\mu}_n}{\partial r} \right|^2 \int_{\partial^+ D_{2^{s_0+s} 2r_n R}(x'_n)} |\widehat{\mu}_n - \widehat{\mu}_n^*|^2 \right)^{\frac{1}{2}} \\ & \leq C \left(\int_{\partial D_{2^{s_0+s} 2r_n R}(x'_n)} \left| \frac{\partial \widehat{\mu}_n}{\partial r} \right|^2 \right)^{\frac{1}{2}} \left(2^{s_0+s} r_n R \int_0^{2\pi} \left| \frac{\partial \widehat{\mu}_n}{\partial \theta} \right|^2 \right)^{\frac{1}{2}} \\ & \leq C 2^{s_0+s} d_n \int_{\partial D_{2^{s_0+s} 2r_n R}(x'_n)} |\nabla \widehat{\mu}_n|^2 \\ & \leq C 2^{s_0+s} d_n \int_{\partial^+ D_{2^{s_0+s} 2r_n R}^+(x'_n)} |\nabla \mu_n|^2 \\ & \leq C 2^{s_0+s} d_n \int_{\partial^+ D_{2^{s_0+s} 2r_n R}^+(x'_n)} |\nabla u_n|^2 + C (2^{s_0+s} r_n R)^2. \end{aligned}$$

Similarly, we can obtain

$$\int_{\partial D_{2^{s_0-s} 2r_n R}(x'_n)} \frac{\partial \widehat{\mu}_n}{\partial r} (\widehat{\mu}_n - \widehat{\mu}_n^*) \leq C 2^{s_0-s} d_n \int_{\partial^+ D_{2^{s_0-s} 2r_n R}^+(x'_n)} |\nabla u_n|^2 + C (2^{s_0-s} r_n R)^2.$$

Taking ϵ and δ sufficiently small, we have

$$\begin{aligned} \int_{Q(s)} |\nabla u_n|^2 dx &\leq C 2^{s_0+s} 2d_n \int_{\partial^+(D_{2^{s_0+s} 2r_n R}^+(x'_n))} |\nabla u_n|^2 \\ &\quad + C 2^{s_0-s} 2d_n \int_{\partial^+(D_{2^{s_0-s} 2r_n R}^+(x'_n))} |\nabla u_n|^2 + C (2^{s_0+s} r_n R)^{1-\frac{2}{p}}, \end{aligned}$$

which gives us

$$f(s) \leq \frac{C}{\log 2} f'(s) + C (2^{s_0+s} r_n R)^{1-\frac{2}{p}}. \tag{3.59}$$

(3.59) implies that

$$(2^{-\frac{s}{c}} f(s))' \geq -C (2^{s_0} r_n R)^{1-\frac{2}{p}} 2^{(1-\frac{2}{p}-\frac{1}{c})s}.$$

Integrating from 2 to L , we arrive at

$$f(2) \leq C 2^{-\frac{1}{c}L} f(L) + C (2^{s_0} r_n R)^{1-\frac{2}{p}} 2^{(1-\frac{2}{p}-\frac{1}{c})L}.$$

The rest proof is the same as the proof in Lemma 3.3. Thus we finish the analysis of energy loss and no neck property on $\Omega_1 \cup \Omega_3, \Omega_4$ and Ω_2 and get (3.48) and (3.49). \square

We can now prove Theorems 1.1 and 1.2.

Proof of Theorems 1.1 and 1.2 Theorem 1.1 is a direct conclusion of Lemmas 3.2 and 3.4.

If $\omega \equiv 0, \tilde{v} = v - \psi$ satisfies

$$div(\beta(u)\nabla\tilde{v}) = -div(\beta(u)\nabla\psi) + \kappa$$

with the boundary condition $\tilde{v}|_{\partial M} = 0$. By Theorem 1 in [21], for any $1 < p < \infty$, we have

$$\|\nabla\tilde{v}\|_{L^p(M)} \leq C(\|\nabla\psi\|_{L^p(M)} + \|\kappa\|_{L^2(M)}).$$

Thus we have

$$\|\nabla v\|_{L^p(M)} \leq C(\|\nabla\psi\|_{L^p(M)} + \|\kappa\|_{L^2(M)}).$$

Then, Theorem 1.2 is a direct conclusion of Lemmas 3.3 and 3.5. \square

4 Applications to the Lorentzian harmonic map flow

At the beginning of this section, let us recall a lemma in [8] which is useful in this part.

Lemma 4.1 (Lemmas 2.1, 2.4 in [8]) *Suppose $(u, v) \in \mathcal{V}(M_0^{T_1}; N \times \mathbb{R})$ is a solution of (1.14) and (1.15), then the Lorentzian energy $E_g(u(t), v(t))$ is non-increasing on $[0, T_1]$ and for any $0 \leq s \leq t < T_1$, there holds*

$$E_g(u(t), v(t)) + \int_s^t \int_M |\partial_t u|^2 dx dt \leq E_g(u(s), v(s)).$$

Moreover, for any $1 < p < \infty, t > 0$, there holds

$$\int_M |\nabla u(\cdot, t)|^2 dx + \int_M |\nabla v(\cdot, t)|^p dx + \int_0^t \int_M |\partial_t u|^2 dx dt \leq C(p, \lambda_1, \lambda_2, \phi, \psi).$$

Lemma 4.2 *Let $(u, v) \in \mathcal{V}(M_0^{T_1}; N \times \mathbb{R})$ be a solution to (1.14) and (1.15). There exists a positive constant $R_0 < 1$ such that, for any $x_0 \in M$, $0 \leq R \leq R_0$ and $0 < s \leq t < T_1$, there hold*

$$E(u(t); B_R^M(x_0)) \leq E(u(s); B_{2R}^M(x_0)) + C \frac{t-s}{R^2}, \tag{4.1}$$

and

$$E(u(s); B_R^M(x_0)) \leq E(u(t); B_{2R}^M(x_0)) + C \frac{t-s}{R^2} + C \int_s^t \int_M |\partial_t u|^2 dx dt, \tag{4.2}$$

where $B_R^M(x_0) \subset M$ is the geodesic ball centered at point x_0 with radius R , C is a positive constant depending on $\lambda_1, \lambda_2, M, N, E(\phi), \|\psi\|_{W^{1,4}(M)}$.

Proof Let $\eta \in C_0^\infty(B_{2R}^M(x_0))$ be a cut-off function such that $\eta(x) = \eta(|x - x_0|)$, $0 \leq \eta \leq 1$, $\eta|_{B_R^M(x_0)} \equiv 1$ and $|\nabla \eta| \leq \frac{C}{R}$. By direct computations, we get

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \int_M |\nabla u|^2 \eta^2 &= \int_M \langle \nabla u, \nabla u_t \rangle \eta^2 \\ &= \int_{\partial B_{2R}^M(x_0)} \frac{\partial u}{\partial r} \cdot u_t \eta^2 - \int_M \langle \Delta u, u_t \rangle \eta^2 - 2 \int_M \nabla u \cdot \nabla \eta \eta u_t \\ &= \int_M \langle -u_t - B^\top(u) |\nabla v|^2, u_t \rangle \eta^2 - 2 \int_M \nabla u \cdot \nabla \eta \eta u_t \\ &= - \int_M |u_t|^2 \eta^2 - \int_M B^\top(u) |\nabla v|^2 \cdot u_t \eta^2 - 2 \int_M \nabla u \cdot \nabla \eta \eta u_t. \end{aligned}$$

On the one hand, by Lemma 4.1 and Young's inequality, we have

$$\frac{d}{dt} \frac{1}{2} \int_M |\nabla u|^2 \eta^2 \leq -\frac{1}{2} \int_M |u_t|^2 \eta^2 + C \int_M |\nabla u|^2 |\nabla \eta|^2 + C \int_M |\nabla v|^4 \eta^2 \leq \frac{C}{R^2}.$$

By integrating the above inequality from s to t , we can get (4.1).

On the other hand, by Lemma 4.1 and Young's inequality, we also have

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \int_M |\nabla u|^2 \eta^2 &\geq -\frac{3}{2} \int_M |u_t|^2 \eta^2 - C \int_M |\nabla u|^2 |\nabla \eta|^2 - C \int_M |\nabla v|^4 \eta^2 \\ &\geq -\frac{3}{2} \int_M |u_t|^2 \eta^2 - \frac{C}{R^2}. \end{aligned}$$

Then (4.2) follows immediately from integrating the above inequality from s to t . □

With the help of Lemma 4.2, we can apply the standard argument (see Lemma 6.4.10 in [20]) to obtain

Lemma 4.3 *Let $(u, v) \in \mathcal{V}(M_0^{T_1}; N \times \mathbb{R})$ be a solution to (1.14) and (1.15). Assume that there is only one singular point $x_0 \in M$ at time T_1 . Then there exists a positive number $m > 0$ such that, as $t \uparrow T_1$,*

$$|\nabla u|^2(x, t) dx \rightarrow m \delta_{x_0} + |\nabla u|^2(x, T_1) dx \text{ as Radon measures.} \tag{4.3}$$

Here δ_{x_0} denotes the δ -mass at x_0 .

Now we shall prove Theorems 1.3 and 1.4.

Proof of Theorem 1.3 In fact, Theorem 1.3 is a consequence of Lemma 4.1, Theorems 1.1 and 1.2.

By Lemma 4.1, we can find a positive sequence $t_n \rightarrow \infty$, such that

$$\lim_{n \rightarrow \infty} \int_M |\partial_t u|^2(\cdot, t_n) dx = 0 \quad \text{and} \quad E(u(\cdot, t_n), v(\cdot, t_n)) \leq C.$$

Taking the sequence to be $(u_n, v_n) = (u(\cdot, t_n), v(\cdot, t_n))$ with $(\tau_n, h_n) = (\partial_t u(\cdot, t_n), 0)$ in Theorems 1.1 and 1.2, the conclusions of Theorem 1.3 follow immediately. \square

Proof of Theorem 1.4 With the help of Lemmas 4.2, 4.3, Theorems 1.1 and 1.2, the proof of (1.19) is almost the same as the proof for the harmonic map flow and we omit the details here. One can refer to [20] for the interior case and to [15, 16] for the boundary case.

It is not hard to prove that there is a unique weak limit $(u(T_1), v(T_1)) \in W^{1,2}(M, N \times \mathbb{R})$ of $(u(t), v(t))$ in $W^{1,2}(M)$ as $t \rightarrow T_1$ (one can refer to the proof of Theorem 1.2 in [14] for a similar argument). Moreover, by Lemma 4.1,

$$v(t) \rightharpoonup v(T_1) \text{ weakly in } W^{1,4}(M).$$

Then, we have

$$\begin{aligned} & \int_M \beta(u(t)) |\nabla v(t)|^2 dx - \int_M \beta(u(T_1)) |\nabla v(T_1)|^2 dx \\ &= \int_M \beta(u(t)) \nabla v(t) \nabla (v(t) - v(T_1)) + (\beta(u(t)) \nabla v(t) - \beta(u(T_1)) \nabla v(T_1)) \nabla v(T_1) dx \\ &= \int_M (\beta(u(t)) - \beta(u(T_1))) \nabla v(t) \nabla v(T_1) + \beta(u(T_1)) (\nabla v(t) - \nabla v(T_1)) \nabla v(T_1) dx \\ &= \text{II} + \text{III}, \end{aligned}$$

where the first term of the second line is zero by integrating by parts and Eq. (1.14). Noting that

$$\text{II} \leq C \|\nabla v(t)\|_{L^4(M)} \|\nabla v(T_1)\|_{L^4(M)} \|u(t) - u(T_1)\|_{L^2(M)},$$

by weak convergence, we have

$$\lim_{t \rightarrow T_1} \int_M \beta(u(t)) |\nabla v(t)|^2 dx = \int_M \beta(u(T_1)) |\nabla v(T_1)|^2 dx.$$

Combining this with (1.19), we get (1.20). \square

Acknowledgements Open access funding provided by Max Planck Society.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Alday, L., Maldacena, J.: Null polygonal Wilson loops and minimal surfaces in anti-de-Sitter space. *J. High Energy Phys.* **082**, 59 (2009)
2. Benci, V., Fortunato, D., Giannoni, F.: On the existence of multiple geodesics in static space-times. *Ann. Inst. H. Poincaré* **8**(1), 79–102 (1991)

3. Bryant, R.: A duality theorem for Willmore surfaces. *J. Differ. Geom.* **20**(1), 23–53 (1984)
4. Chen, Q., Jost, J., Li, J., Wang, G.: Dirac-harmonic maps. *Math. Z.* **254**(2), 409–432 (2006)
5. Ding, W., Tian, G.: Energy identity for a class of approximate harmonic maps from surfaces. *Commun. Anal. Geom.* **3**(4), 543–554 (1995)
6. Greco, C.: The Dirichlet-problem for harmonic maps from the disk in to a lorentzian warped product. *Ann. Inst. H. Poincaré* **10**(2), 239–252 (1993)
7. Greco, C.: A multiplicity result for the Dirichlet problem for harmonic maps from the disk into a Lorentzian warped product. *Nonlinear Anal.* **28**(10), 1661–1668 (1997)
8. Han, X., Jost, J., Liu, L., Zhao, L.: Global Existence of Harmonic Map Heat Flow Into Lorentzian Manifolds. MPI MIS Preprint: 70/2016. <https://www.mis.mpg.de/publications/preprints/2016/prepr2016-70.html>. Accessed 11 Oct 2016
9. Han, X., Zhao, L., Zhu, M.: Energy identity for harmonic maps into standard stationary Lorentzian manifolds. *J. Geom. Phys.* **114**, 621–630 (2017)
10. Huang, T., Wang, C.: Boundary Bubbling Analysis of Approximate Harmonic Maps Under Either Weak or Strong Anchoring Conditions in Dimension Two. [arXiv:1607.05357](https://arxiv.org/abs/1607.05357)
11. Isobe, T.: Optimal regularity of harmonic maps from a Riemannian manifold into a static Lorentzian manifold. *Pac. J. Math.* **178**(1), 71–93 (1997)
12. Isobe, T.: Regularity of harmonic maps into a static Lorentzian manifold. *J. Geom. Anal.* **8**(3), 447–463 (1998)
13. Jost, J.: *Two-Dimensional Geometric Variational Problems*. Wiley, New York (1991)
14. Jost, J., Liu, L., Zhu, M.: A global weak solution of the Dirac-harmonic map flow. *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2017). <https://doi.org/10.1016/j.anihpc.2017.01.002>
15. Jost, J., Liu, L., Zhu, M.: The Qualitative Behavior at the Free Boundary for Approximate Harmonic Maps from Surfaces. MPI MIS Preprint: 26/2016. <https://www.mis.mpg.de/publications/preprints/2016/prepr2016-26.html>. Accessed 21 Mar 2016
16. Jost, J., Liu, L., Zhu, M.: Bubbling Analysis Near the Dirichlet Boundary for Approximate Harmonic Maps from Surfaces. MPI MIS Preprint: 38/2016 (to appear in *Comm. Anal. Geom.*). <https://www.mis.mpg.de/publications/preprints/2016/prepr2016-38.html>. Accessed 29 June 2016
17. Kramer, D., Stephani, H., Hertl, E., MacCallum, M.: *Exact Solutions of Einsteins Field Equations*. Cambridge University Press, Cambridge (1980)
18. Lemaire, L.: Applications harmoniques de surfaces riemanniennes. *J. Differ. Geom.* **13**(1), 51–78 (1978)
19. Li, J., Liu, L.: Partial Regularity of Harmonic Maps from a Riemannian Manifold Into a Lorentzian Manifold. [arXiv:1704.08673](https://arxiv.org/abs/1704.08673)
20. Lin, F., Wang, C.: *The Analysis of Harmonic Maps and Their Heat Flows*, p. xii+267. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ (2008)
21. Meyers, G.: An L^p -estimate for the gradient of solutions of second order elliptic divergence equations. *Annali Della Scuola Normale superiore di Pisa-Classe di Scienze* **17**(3), 189–206 (1963)
22. O’Neill, B.: *Semi-Riemannian Geometry: With Applications to Relativity Pure and Applied Mathematics*, vol. 103, p. xiii+468. Academic Press Inc./Harcourt Brace Jovanovich Publishers, New York (1983)
23. Parker, T.: Bubble tree convergence for harmonic maps. *J. Differ. Geom.* **44**(3), 595–633 (1996)
24. Qing, J., Tian, G.: Bubbling of the heat flows for harmonic maps from surfaces. *Commun. Pure Appl. Math.* **50**(4), 295–310 (1997)
25. Rivière, T.: Conservation laws for conformally invariant variational problems. *Invent. Math.* **168**(1), 1–22 (2007)
26. Rivière, T., Struwe, M.: Partial regularity for harmonic maps and related problems. *Commun. Pure Appl. Math.* **61**(4), 451–463 (2007)
27. Sacks, J., Uhlenbeck, K.: The existence of minimal immersions of 2-spheres. *Ann. Math.* **113**, 1–24 (1981)
28. Sharp, B.: Higher integrability for solutions to a system of critical elliptic PDE. *Methods Appl. Anal.* **21**(2), 221–240 (2014)
29. Sharp, B., Topping, P.: Decay estimates for Riviere’s equation, with applications to regularity and compactness. *Trans. Am. Math. Soc.* **365**(5), 2317–2339 (2013)
30. Sharp, B., Zhu, M.: Regularity at the free boundary for Dirac-harmonic maps from surfaces. *Calc. Var. Partial Differ. Equ.* **55**(27), 1–30 (2016)
31. Zhao, L.: Energy identities for Dirac-harmonic maps. *Calc. Var. Partial Differ. Equ.* **28**(1), 121–138 (2007)
32. Zhu, M.: Regularity for harmonic maps into certain pseudo-Riemannian manifolds. *J. Math. Pures Appl.* **99**(1), 106–123 (2013)