

Management of postoperative arterial hemorrhage after pancreato-biliary surgery according to the site of bleeding: re-laparotomy or interventional radiology

Fumihiko Miura · Takehide Asano · Hodaka Amano · Masahiro Yoshida · Naoyuki Toyota · Keita Wada · Kenichiro Kato · Eriko Yamazaki · Susumu Kadowaki · Makoto Shibuya · Sawako Maeno · Shigeru Furui · Koji Takeshita · Yutaka Kotake · Tadahiro Takada

Received: 14 December 2007 / Accepted: 16 January 2008 / Published online: 26 December 2008
© Springer 2008

Abstract

Background/Purpose Intra-abdominal arterial hemorrhage is still one of the most serious complications after pancreato-biliary surgery. We retrospectively analyzed our experiences with 15 patients in order to establish a therapeutic strategy for postoperative arterial hemorrhage following pancreato-biliary surgery.

Methods Between August 1981 and November 2007, 15 patients developed massive intra-abdominal arterial bleeding after pancreato-biliary surgery. The initial surgery of these 15 patients were pylorus-preserving pancreatoduodenectomy (PPPD) (7 patients), hemihepatectomy and caudate lobectomy with extrahepatic bile duct resection or PPPD (4 patients), Whipple's pancreatoduodenectomy (PD) (3 patients), and total pancreatectomy (1 patient). Twelve patients were managed by transcatheter arterial embolization and three patients underwent re-laparotomy.

Results Patients were divided into two groups according to the site of bleeding: SMA group, superior mesenteric artery (4 patients); HA group, stump of gastroduodenal artery, right hepatic artery, common hepatic artery, or proper hepatic artery (11 patients). In the SMA group, re-laparotomy and coil embolization for pseudoaneurysm were performed in three and one patients, respectively, but none of the patients survived. In the HA group, all 11

patients were managed by transcatheter arterial embolization. None of four patients who had major hepatectomy with extrahepatic bile duct resection survived. Six of seven patients (85.7%) who had pancreatectomy survived, although hepatic infarction occurred in four.

Conclusions Management of postoperative arterial hemorrhage after pancreato-biliary surgery should be done according to the site of bleeding and the initial operative procedure. Careful consideration is required for indication of interventional radiology for bleeding from SMA after pancreatectomy and hepatic artery after major hepatectomy with bilioenteric anastomosis.

Keywords Complications · Postoperative bleeding · Pancreato-biliary surgery

Introduction

Despite a declining mortality rate after pancreato-biliary surgery [1–4], intra-abdominal arterial hemorrhage is still one of the most serious complications [5–8]. Arterial bleeding has a disastrous clinical outcome, with a reported mortality rate of 20–50% [6, 9, 10]. With the recent advances in interventional radiology (IVR), transcatheter arterial embolization (TAE) has been widely used, gaining acceptance for the treatment of visceral aneurysm or pseudoaneurysm. Various articles regarding TAE for postoperative ruptured pseudoaneurysm have reported success rates of 50–100% [6, 11–13]. To our knowledge, management of postoperative arterial hemorrhage after pancreato-biliary surgery according to the site of bleeding and clinical condition including the type of operative procedure has not yet been advocated and, therefore, a detailed indication for TAE for this condition has still to be

F. Miura · T. Asano (✉) · H. Amano · M. Yoshida · N. Toyota · K. Wada · K. Kato · E. Yamazaki · S. Kadowaki · M. Shibuya · S. Maeno · T. Takada
Department of Surgery, Teikyo University School of Medicine, 2-11-1, Kaga, Itabashi-ku, Tokyo 173-8605, Japan
e-mail: asn@med.teikyo-u.ac.jp

S. Furui · K. Takeshita · Y. Kotake
Department of Radiology,
Teikyo University School of Medicine, Tokyo, Japan

established. In this article, we retrospectively analyzed our experiences with 15 patients in order to establish a therapeutic strategy for postoperative arterial hemorrhage following hepato-biliary-pancreatic surgery.

Patients and methods

Patients

Between August 1981 and November 2007, 708 patients underwent major pancreato-biliary surgery at the Department of Surgery, Teikyo University Hospital. Fifteen of these 708 patients (2.1%) (11 men and 4 women, 50–82 [mean 65.2] years old) developed massive intra-abdominal arterial bleeding after the surgery. These 15 patients were divided into two groups according to the site of bleeding: SMA group, bleeding from the superior mesenteric artery (SMA) (4 patients); HA group, bleeding from the stump of the gastroduodenal artery (GDA), right hepatic artery (RHA), common hepatic artery (CHA), or proper hepatic artery (PHA) (11 patients).

Patient characteristics are listed in Table 1. Diagnoses at initial surgery were pancreatic head carcinoma (4 patients), lower bile duct carcinoma (4 patients), perihilar cholangiocarcinoma (2 patients), ampullary adenocarcinoma (2 patients), intrahepatic cholangiocarcinoma (1 patient), gallbladder carcinoma (1 patient), and chronic pancreatitis (1 patient). The procedures of the initial surgery were pylorus-

preserving pancreatoduodenectomy (PPPD) (7 patients), Whipple's pancreatoduodenectomy (PD) (3 patients), right hemihepatectomy and caudate lobectomy with extrahepatic bile duct resection (2 patients), right hemihepatectomy and caudate lobectomy with PPPD (1 patient), left hemihepatectomy and caudate lobectomy with extrahepatic bile duct resection (1 patient), and total pancreatectomy (1 patient). In the patients with malignant disease, skeletonization of the hepatic artery was routinely performed to remove neural and lymphatic tissue of the hepatoduodenal ligament. In patients with pancreatic head carcinoma, the SMA was right-semicircularly exposed to resect the surrounding right-sided nerve plexus. Sentinel bleeding was defined as minimal bleeding from either abdominal drains or the gastrointestinal tract without evidence of peripheral circulatory impairment or transfusion requirement.

TAE technique

Patients underwent emergency angiography using the standard Seldinger technique and an angiography catheter. At first, superior mesenteric arteriography, arterial portography, and celiac arteriography were performed with a 4 F preshaped catheter. When a pseudoaneurysm or extravasation of contrast media was identified, transcatheter microcoil embolotherapy was performed. A 2.9 F microcatheter (Sniper; Clinical Supply, Gifu, Japan, or Progreat; Terumo, Tokyo, Japan) was then inserted via the angiography catheter. Coil embolization, both proximal and distal of the

Table 1 Profiles of 15 patients encountering intra-abdominal arterial bleeding after pancreato-biliary surgery

Pt. no.	Age (year)/gender	Diagnosis	Initial operation
<i>SMA group</i>			
1	50/M	Ampullary carcinoma	PPPD
2	53/M	Pancreatic head carcinoma	PPPD
3	59/M	Pancreatic head carcinoma	PD
4	64/M	Pancreatic head carcinoma	PPPD
<i>HA group</i>			
5	78/M	Hilar bile duct carcinoma	Right hepatic lobectomy + S1 + BDR
6	73/F	Gall bladder carcinoma	Right hepatic lobectomy + S1 + PPPD + PVR
7	63/F	Hilar bile duct carcinoma	Right hepatic lobectomy + S1 + BDR
8	69/M	Intrahepatic cholangiocarcinoma	Left hepatic lobectomy + S1 + BDR + PVR
9	68/M	Pancreatic head carcinoma	PD
10	66/M	Ampullary carcinoma	PPPD
11	82/M	Lower bile duct carcinoma	PPPD
12	77/F	Lower bile duct carcinoma	PD
13	56/M	Lower bile duct carcinoma	PPPD
14	60/F	Chronic pancreatitis	TP
15	60/M	Lower bile duct carcinoma	PPPD

PPPD Pylorus preserving pancreatoduodenectomy; PD pancreatoduodenectomy; S1 caudate lobectomy; BDR extrahepatic bile duct resection; PVR portal vein resection; TP total pancreatectomy

bleeding site, if possible, or pseudoaneurysm was performed with fibered platinum microcoils of various diameters and shapes (VortX; Boston Scientific, Natick, MA, USA, Truffill; Cordis, Miami Lakes, FL, USA; and Complex Helical Fibered Platinum Coil-18; Target Vascular, Fremont, CA, USA) and interlocking detachable coil (IDC) (Boston-Scientific, Natick, MA, USA) via the microcatheter.

Complication definitions

Leakage of a pancreatojejunostomy and pancreatic fistula was defined as amylase-rich exudate (amylase more than three times normal serum concentration) from the drainage tube placed at the pancreatojejunostomy and near the pancreas, respectively. Leakage of a hepaticojejunostomy was defined as the persistent presence of bile from the drainage tube placed at the hepaticojejunostomy. Diagnosis of hepatic infarction was based on the coexistence of CT findings (geographic, low attenuation perfusion defects without mass effect) seen in association with an acute increase in the serum alanine aminotransaminase level to more than 1,000 IU.

Results

The results are shown in Table 2. The interval between initial surgery and the first intervention in the 15 patients was 5–39 days (mean 22.6 days). All patients had intra-abdominal infections, the cause being leakage of pancreatojejunostomy in six patients, leakage of hepaticojejunostomy in five, and leakage of pancreatojejunostomy and hepaticojejunostomy, leakage of gastrojejunostomy, pancreatic fistula, and perforation of the jejunal limb in one patient each. Sentinel bleeding was noted in seven patients (46.7%) for 1–10 (mean 2.2) days before massive bleeding. Sentinel bleeding was identified as bleeding from drain in all seven patients. When sentinel bleeding occurred, contrast-enhanced CT and angiography were performed in five and four patients, respectively, but neither pseudoaneurysm nor extravasation was demonstrated on either examination. On the occurrence of massive bleeding, contrast-enhanced CT was performed before angiography in eight patients. Intra-abdominal pseudoaneurysms and hematomas, hematomas only, and fluid collection were demonstrated in five, two, and one patient, respectively. Angiography was performed in 14 patients, and pseudoaneurysms or extravasations were demonstrated in all of them.

SMA group

The SMA group consisted of four patients in whom bleeding occurred from SMA. Three patients required re-

laparotomy, and coil embolization was performed in one. None of the patients in this group survived.

One patient (patient 2) was transferred to the operating room as soon as bleeding from drain was identified, without first performing angiography. The bleeding site could not be reached at re-laparotomy due to dense adhesion, serious inflammation, and massive bleeding; therefore ligation of the fistula was performed. Complete hemostasis was not obtained and he was died of bleeding shortly after the second surgery. The site of bleeding was detected at autopsy.

Angiography was performed in three patients. Extravasations from the right wall of SMA were demonstrated in two patients (patients 1 and 3) and a pseudoaneurysm of a tributary of SMA was identified in one patient (patient 4). Two patients (patients 1 and 3) were transferred to the operating room because TAE was avoided for fear of ischemia of the small intestine. In patient 1, SMA was cut both proximal and distal to the bleeding point, and re-arterialization was performed with a 6 mm polytetrafluoroethylene (Gore-Tex) graft. Because the jejunal limb lapsed into necrosis during the operation, the jejunal limb was resected with disassembling pancreatojejunostomy and hepaticojejunostomy. Tube hepaticostomy and pancreateostomy were performed. Although hemostasis was obtained, he did not recover from circulatory failure and died one day after the second operation. In patient 3, bleeding was stopped by suturing. The jejunal limb was resected disassembling pancreatojejunostomy and hepaticojejunostomy in order to detect the bleeding point by improving the operative field, although insufficiency of pancreatojejunostomy was not noted. Tube hepaticostomy and pancreateostomy were performed. The upper half of the midline incision was not closed, allowing open drainage. After the second operation, bleeding occurred several times and was stopped by suturing at bedside. The general state deteriorated gradually, and he died of hepatic failure 58 days after the initial surgery.

In one patient (patient 4), bleeding from CHA was also identified prior to bleeding from SMA. For the CHA bleeding, coil embolization was performed for CHA pseudoaneurysm using interlocking detachable coils 34 days after the initial surgery. Re-bleeding from CHA was not noted, but bleeding from a pseudoaneurysm of a tributary of SMA occurred 46 days after the surgery, and coil embolization was performed on the same day (Fig. 1a). He encountered re-bleeding from pseudoaneurysms of SMA at days 52, 63, 103 post surgery (Fig. 1b). For each re-bleeding, coil-embolization was performed for the pseudoaneurysms using interlocking detachable coils. The patency of SMA was preserved in order to prevent ischemia of the small intestine. He died of multiple organ failure 110 days after the initial surgery.

Table 2 Results of 15 patients encountering intra-abdominal arterial bleeding after pancreato-biliary surgery

Pt. no.	Sentinel bleeding	Bleeding site	Postoperative day of bleeding	Cause of intra-abdominal infection	Intervention for bleeding	Complication or result of coil embolization	Outcome
<i>SMA group</i>							
1	Yes	SMA	27	PJ leakage	Replacement of artificial vessel, choledocostomy, and pancreatostomy	Not performed	Died of hemorrhage
2	No	SMA	20	HJ leakage	Surgical hemostasis	Not performed	Died of hemorrhage
3	No	SMA	28	HJ leakage	Surgical hemostasis, choledocostomy, and pancreatostomy	Not performed	Died of hepatic failure
4	Yes	CHA SMA	34 46	PJ leakage	Coil embolization of pseudoaneurysm Coil embolization of pseudoaneurysm 4 times	Re-bleeding	Died of hemorrhage
<i>HA group</i>							
5	Yes	CHA	9	Pancreatic fistula	TAE of CHA	Unsuccessful	Died of hemorrhage
6	No	GDA	26	PJ leakage	TAE of CHA	Hepatic failure	Died of hepatic failure
7	Yes	PHA	23	HJ leakage	TAE of PHA	Hepatic failure	Died of hepatic failure
8	No	PHA	6	HJ leakage	TAE of PHA	Hepatic failure	Died of hepatic failure
9	Yes	GDA	26	HJ leakage	TAE of CHA	Hepatic infarction	Died of DIC
10	Yes	GDA	31	PJ leakage	TAE of CHA(LHA)	Success	Recovered
11	No	RHA	9	PJ and HJ leakage	TAE of PHA	Hepatic infarction	Recovered
12	Yes	RHA	25	GJ leakage	TAE of RHA	Success	Recovered
13	No	GDA	21	PJ leakage	TAE of CHA	Hepatic infarction	Recovered
14	No	GDA	27	Jejunal limb perforation	TAE of CHA	Hepatic infarction	Recovered
15	No	GDA	27	PJ leakage	TAE of CHA(LHA)	Success	Recovered

SMA Superior mesenteric artery; *PJ* pancreaticojejunostomy; *HJ* hepaticojejunostomy; *CHA* common hepatic artery; *TAE* transcatheter arterial embolization; *PHA* proper hepatic artery; *GDA* gastroduodenal artery; *RHA* right hepatic artery

HA group

The HA group consisted of 11 patients in whom bleeding sites were major branches of the celiac artery: stump of GDA in six patients, PHA in two, RHA in two, and CHA in one. TAE was performed in all 11 patients. The sites of embolization were CHA in seven patients, PHA in three, and RHA in one. Portal vein patency was confirmed by superior mesenteric arteriography before embolization in all patients. In 10 of 11 patients, complete hemostasis was obtained and rebleeding was not noted in any of the patients.

All four patients undergoing hepatic lobectomy for biliary hilar malignancy (patients 5–8) died, three of hepatic failure, and one of persistent bleeding within 24 h. On angiography, there were collateral pathways for the arterial supply to hepatic parenchyma in two patients: accessory

left hepatic artery arising from left gastric artery (patient 6) and right inferior phrenic artery (patient 7). In two of the four patients, serum total bilirubin concentrations before bleeding were more than 10 mg/dl, and in the other two patients they were less than 3.0 mg/dl.

In seven patients undergoing pancreatectomy (patients 9–15), one patient (patient 9) died of disseminated intravascular coagulation (DIC) 10 days after TAE, and six patients recovered. Hepatic infarction was encountered in four patients. The sites of infarction on CT were lateral segment and S5 in three patients and lateral segment in one. Three of the four patients encountering hepatic infarction improved with lateral segmentectomy performed 3 days after TAE in one patient (patient 11) (Fig. 2), plasma exchange in one (patient 13), and medical treatment in one (patient 14).



Fig. 1 Superior mesenteric arteriography of patient 4. **a** Coil embolization for pseudoaneurysm of tributary of the superior mesenteric artery was performed. Temporary hemostasis was obtained using interlocking detachable coils. The *white arrow* indicates interlocking detachable coils in the pseudoaneurysm of the stump of the GDA. The *black arrow* indicates interlocking detachable coils in the aneurysm of the tributary of the superior mesenteric

artery. **b** Angiography revealed enlargement of pseudoaneurysm (*arrow*) after 6 days of coil embolization for a pseudoaneurysm of a tributary of the superior mesenteric artery. Re-bleeding from pseudoaneurysms of SMA occurred 52, 63, and 103 days after the initial surgery. For each re-bleeding, coil embolizations were performed for the pseudoaneurysms with interlocking detachable coils

In three patients without hepatic infarction, either right (RHA) or left hepatic artery (LHA) was preserved, as the sites of embolization were RHA (patient 8) and CHA (LHA) accompanied by replaced RHA arising from SMA (patients 10 and 15).

Discussion

Sentinel bleeding was first described by Brodsky and Turnbull [6] in 1991 as hemorrhage of drainage fluid or in the gastrointestinal tract. Sato et al. [11] emphasized the significance of early angiography promptly after the recognition of sentinel bleeding, since all 10 patients with massive bleeding after pancreatectomy had sentinel bleeding. However, the efficacy of early angiography for sentinel bleeding was not evaluated. In this study, on the occurrence of sentinel bleeding, CT and angiography failed to demonstrate the site of bleeding in five and four patients, respectively. Sentinel bleeding is less likely to be visualized by angiography, possibly due to the intermittent character of hemorrhage [10, 14]. Okuno et al. [15] reported that 6 patients with episodes of sentinel bleeding after hepato-biliary-pancreatic surgery obtained hemostasis through medical treatment. Although sentinel bleeding does not always accompany subsequent massive bleeding, surgeons should prepare for possible massive bleeding when sentinel bleeding is identified.

There have been a few reports describing postoperative hemorrhage from SMA in detail. Blockage of blood flow



Fig. 2 Lateral segmentectomy was performed for hepatic infarction 3 days after coil embolization for CHA in patient 11. At laparotomy the entire portion of the lateral segment of the liver seemed to be necrotic (*arrows*)

from SMA reduces ischemic necrosis of the small intestine. Therefore, we did not perform TAE for SMA in any of the patients. Unfortunately, none of the patients with bleeding from SMA was successfully treated in this study. Recently, a few case reports described the successful treatment of

patients with chronic pancreatitis encountering bleeding from SMA by TAE of pseudoaneurysm using interlocking detachable coils [16, 17]. Because the design of the interlocking detachable coil allows controlled release of the device and its accurate placement, it may also be an effective means of treating pseudoaneurysms [17]. In one patient (patient 4), we performed coil embolization for pseudoaneurysm of SMA by using interlocking detachable coils, but repeated bleeding occurred due to sepsis from pancreatojejunal anastomotic insufficiency. The long-term success of embolization of the pseudoaneurysm with interlocking detachable coils is unknown. One case report has described coil compaction after embolization of SMA pseudoaneurysm [18]. In cerebral aneurysm, coil compaction following successful embolization often occurs [16]. Even if an aneurysm is completely embolized, the “water-hammer effect” of blood pulsation transmitted into the coil-thrombus complex may induce eventful coil compaction or aneurysm regrowth with potential for late bleeding [19]. In aneurysms of SMA, abdominal movements and pancreatic enzymes may also increase the risk of coil compaction [18]. There is a report that continuous intra-arterial infusion of vasopressin for bleeding from a branch of SMA after PD obtained complete hemostasis [15]. However, whether this procedure using vasopressin would be effective for massive bleeding from SMA remains unclear.

The first line of treatment for massive bleeding from hepatic artery after pancreatectomy is TAE. However, the long-term result of this treatment is unknown, and liver infarction or ischemia has been reported as a mortal complication of TAE for pseudoaneurysms arising from hepatic arteries [11, 13, 20]. It is commonly believed that the dual blood supply and extensive collateral pathways of the liver serve to protect this organ from ischemic insult. Occlusion of the hepatic artery is usually well-tolerated and has been performed safely in some cases following PD. However, recently Yoon et al. [21] revealed a 30% incidence of liver abscess after hepatic artery occlusion for bleeding pseudoaneurysms after PD. Kim et al. [22] found that six out of seven patients had hepatic abscesses following hepatic artery embolization after having undergone a prior Whipple procedure. In our series, liver infarction occurred in five patients. Ligation of the PHA results in the development of small collateral vessels in the hepatic ligaments and around the common bile duct [23]. These collateral vessels may be noted immediately after temporary occlusion of hepatic arterial flow [23]. However, collateral arteries around the common bile duct were disrupted after bile duct resection. The fact that infarction of the entire liver did not occur in any patients must be attributable to recruitment of other collateral pathways including inferior phrenic arteries,

intercostal arteries, and gastric arteries that were presumably not ligated at the initial surgery.

There have been a few detailed reports in the literature concerning postoperative hemorrhage after major hepatectomy for biliary hilar malignancy [15, 24]. Nagino et al. [24] treated nine patients who encountered postoperative hemorrhage after hepatectomy for hilar cholangiocarcinoma. Re-laparotomy was performed in eight patients, and seven survived. In only one patient was TAE used for rupture of the RHA after extended left hepatic lobectomy. The patient suffered from liver failure following embolization, but he recovered. In our series, all four patients after major hepatectomy with extrahepatic bile duct resection died of hepatic failure or persistent bleeding within 24 h after TAE of hepatic artery. These results imply that TAE for postoperative hemorrhage after major hepatectomy with extrahepatic bile duct resection is nearly contraindicated. Experimental and clinical studies elucidated that partial arterialization of the portal vein for hepatic artery occlusion after hepatic resection could preclude hepatic necrosis or failure [25, 26]. Iseki et al. [27] reported three successful cases of concomitant hepatic artery resection with arterioportal shunt for pancreato-biliary malignancy, advocating that mesenteric arterioportal shunt has the advantages of choice of arterial size and a site removed from the primary operative field.

The recent refinement of IVR has been solving the most serious complications of TAE, namely, mesenteric ischemia and hepatic infarction or failure after TAE for SMA and hepatic artery, respectively. In recent years, covered stent-grafts have been used in the endovascular repair of abdominal aneurysm. Obliteration of the aneurysmal neck results in effective obliteration of the aneurysm without block of arterial flow [28]. In 1998, McGraw et al. [29] first reported the successful use of covered stent for management of a SMA pseudoaneurysm occurring after PD. In 2000, Burger et al. [30] first described the successful stent-graft repair of a ruptured hepatic artery aneurysm after PD. Some authors actually consider this therapeutic option as the first-line treatment for a bleeding pseudoaneurysm in critical situations where the consequences of superior mesenteric or hepatic artery occlusion may be disastrous [20, 30–32]. The procedure remains risky due to the possibility of rupturing the artery because of its fragile vascular wall and infection involving the covered stent itself, since the vessel is affected by septic condition [33]. It has to be mentioned, however, that this angiographic intervention in branches of the celiac trunk is technically demanding, even in experienced centers, and is not always possible for anatomical reasons [20]. In addition, long-term results of endovascular covered stent placement have yet to be evaluated; therefore further investigation will be necessary to determine the feasibility of this treatment. In

Japan, no covered stent for visceral arteries is commercially available; thus we do not perform this procedure.

Re-laparotomy is sometimes required for control of bleeding that was technically impossible to stop by angiography, or for recurrent bleeding after coil embolization [20]. Some authors claimed that surgical intervention was the first-line treatment for postoperative massive hemorrhage following pancreato-biliary surgery [34–36]. There are two major advantages of surgical intervention over TAE. Hemostasis by suturing could stop arterial bleeding without blocking blood flow. Control of infection could be achieved by completion pancreatectomy, tube pancreatectomy, or appropriate drainage tube placement [34, 35]. However, a surgical approach to bleeding pseudoaneurysm is sometimes unsuccessful because of dense adhesion and a poor operative view filled with blood. In such situations the aortic occlusion balloon catheter method is helpful for controlling oligemic shock and obtaining a good view of the operative field [34].

Randomized controlled trials (RCTs) are needed to elucidate the appropriate management for massive bleeding after pancreato-biliary surgery. However, it is difficult to plan RCTs in an ethically acceptable manner, and besides that, the number of patients encountering postoperative hemorrhage is fortunately quite limited. A drawback of the present study, of course, is this very fact, the limited number of patients.

In any event, we think that postoperative arterial hemorrhage after pancreato-biliary surgery should be managed according to the site of bleeding and the initial operative procedure. The first-line treatment for bleeding from major branches of the celiac artery after pancreatectomy is TAE, in spite of the fact that the possibility of hepatic infarction does exist. Careful consideration is required for the indication of IVR for bleeding from SMA after pancreatectomy and hepatic artery after major hepatectomy with bilioenteric anastomosis. Treatment options including partial arterialization of the portal vein and re-laparotomy should be considered for the prevention of hepatic failure and re-bleeding. Angiographic stenting with preservation of superior mesenteric and hepatic arterial flow, if technically possible, might represent the best treatment option.

References

- Cameron JL, Pitt HA, Yeo CJ, Lillemoe KD, Kaufman HS, Coleman J. One hundred and forty-five consecutive pancreaticoduodenectomies without mortality. *Ann Surg.* 1993;217:430–5. Discussion 5–8.
- Pellegrini CA, Heck CF, Raper S, Way LW. An analysis of the reduced morbidity and mortality rates after pancreaticoduodenectomy. *Arch Surg.* 1989;124:778–81.
- Grace PA, Pitt HA, Tompkins RK, DenBesten L, Longmire WP Jr. Decreased morbidity and mortality after pancreaticoduodenectomy. *Am J Surg.* 1986;151:141–9.
- Yokoyama Y, Nagino M, Nishio H, Ebata T, Igami T, Nimura Y. Recent advances in the treatment of hilar cholangiocarcinoma: portal vein embolization. *J Hepatobiliary Pancreat Surg.* 2007;14:447–54.
- Watanabe M, Usui S, Kajiwara H, Nakamura M, Sumiyama Y, Takada T, et al. Current pancreatogastrointestinal anastomotic methods: results of a Japanese survey of 3109 patients. *J Hepatobiliary Pancreat Surg.* 2004;11:25–33.
- Brodsky JT, Turnbull AD. Arterial hemorrhage after pancreaticoduodenectomy. The 'sentinel bleed'. *Arch Surg.* 1991;126:1037–40.
- Miedema BW, Sarr MG, van Heerden JA, Nagorney DM, McIlrath DC, Ilstrup D. Complications following pancreaticoduodenectomy. Current management. *Arch Surg.* 1992;127:945–9. Discussion 9–50.
- Yamaguchi K, Tanaka M, Chijiwa K, Nagakawa T, Imamura M, Takada T. Early and late complications of pylorus-preserving pancreaticoduodenectomy in Japan 1998. *J Hepatobiliary Pancreat Surg.* 1999;6:303–11.
- Lerut JP, Gianello PR, Otte JB, Kestens PJ. Pancreaticoduodenal resection. Surgical experience and evaluation of risk factors in 103 patients. *Ann Surg.* 1984;199:432–7.
- van Berge Henegouwen MI, Allema JH, van Gulik TM, Verbeek PC, Obertop H, Gouma DJ. Delayed massive haemorrhage after pancreatic and biliary surgery. *Br J Surg.* 1995;82:1527–31.
- Sato N, Yamaguchi K, Shimizu S, Morisaki T, Yokohata K, Chijiwa K, et al. Coil embolization of bleeding visceral pseudoaneurysms following pancreatectomy: the importance of early angiography. *Arch Surg.* 1998;133:1099–102.
- Shibata T, Sagoh T, Ametani F, Maetani Y, Itoh K, Konishi J. Transcatheter microcoil embolotherapy for ruptured pseudoaneurysm following pancreatic and biliary surgery. *Cardiovasc Intervent Radiol.* 2002;25:180–5.
- Reber PU, Baer HU, Patel AG, Wildi S, Triller J, Buchler MW. Superselective microcoil embolization: treatment of choice in high-risk patients with extrahepatic pseudoaneurysms of the hepatic arteries. *J Am Coll Surg.* 1998;186:325–30.
- Tien YW, Lee PH, Yang CY, Ho MC, Chiu YF. Risk factors of massive bleeding related to pancreatic leak after pancreaticoduodenectomy. *J Am Coll Surg.* 2005;201:554–9.
- Okuno A, Miyazaki M, Ito H, Ambiru S, Yoshidome H, Shimizu H, et al. Nonsurgical management of ruptured pseudoaneurysm in patients with hepatobiliary pancreatic diseases. *Am J Gastroenterol.* 2001;96:1067–71.
- Tan BS, Reidy JF. Case report: transcatheter embolization of a superior mesenteric artery pseudoaneurysm with interlocking detachable coils. *Clin Radiol.* 1998;53:455–7.
- Hama Y, Kaji T, Iwasaki Y, Kyoto Y, Kawarabayashi N, Hatsuse K, et al. Transcatheter embolization of a superior mesenteric artery pseudoaneurysm. A case report. *Acta Radiol.* 1999;40:649–51.
- Hama Y, Iwasaki Y, Kaji T, Hatsuse K, Kusano S. Coil compaction after embolization of the superior mesenteric artery pseudoaneurysm. *Eur Radiol.* 2002;12(Suppl 3):S189–91.
- Redekop GJ, Durity FA, Woodhurst WB. Management-related morbidity in unselected aneurysms of the upper basilar artery. *J Neurosurg.* 1997;87:836–42.
- Makowiec F, Riediger H, Euringer W, Uhl M, Hopt UT, Adam U. Management of delayed visceral arterial bleeding after pancreatic head resection. *J Gastrointest Surg.* 2005;9:1293–9.
- Yoon YS, Kim SW, Her KH, Park YC, Ahn YJ, Jang JY, et al. Management of postoperative hemorrhage after pancreaticoduodenectomy. *Hepatogastroenterology.* 2003;50:2208–12.

22. Kim W, Clark TW, Baum RA, Soulen MC. Risk factors for liver abscess formation after hepatic chemoembolization. *J Vasc Interv Radiol*. 2001;12:965–8.
23. Redman HC, Reuter SR. Arterial collaterals in the liver hilus. *Radiology*. 1970;94:575–9.
24. Nagino M, Kamiya J, Uesaka K, Sano T, Yamamoto H, Hayakawa N, et al. Complications of hepatectomy for hilar cholangiocarcinoma. *World J Surg*. 2001;25:1277–83.
25. Shimizu Y, Miyazaki M, Shimizu H, Ito H, Nakagawa K, Ambiru S, et al. Beneficial effects of arterialization of the portal vein on extended hepatectomy. *Br J Surg*. 2000;87:784–9.
26. Iseki J, Takagi M, Touyama K, Sano K, Nakagami K, Ori T, et al. Editorial comment: feasibility of central gastrectomy for gastric cancer. *Surgery*. 2003;133:68–73.
27. Iseki J, Noie T, Touyama K, Nakagami K, Takagi M, Ori T, et al. Mesenteric arteriportal shunt after hepatic artery interruption. *Surgery*. 1998;123:58–66.
28. Cragg AH, Dake MD. Treatment of peripheral vascular disease with stent-grafts. *Radiology*. 1997;205:307–14.
29. McGraw JK, Patzik SB, Gale SS, Dodd JT, Boorstein JM. Autogenous vein-covered stent for the endovascular management of a superior mesenteric artery pseudoaneurysm. *J Vasc Interv Radiol*. 1998;9:779–82.
30. Burger T, Halloul Z, Meyer F, Grote R, Lippert H. Emergency stent-graft repair of a ruptured hepatic artery secondary to local postoperative peritonitis. *J Endovasc Ther*. 2000;7:324–7.
31. Nyman U, Svendsen P, Jivegard L, Klingenstierna H, Risberg B. Multiple pancreaticoduodenal aneurysms: treatment with superior mesenteric artery stent-graft placement and distal embolization. *J Vasc Interv Radiol*. 2000;11:1201–5.
32. Rami P, Williams D, Forauer A, Cwikiel W. Stent-graft treatment of patients with acute bleeding from hepatic artery branches. *Cardiovasc Intervent Radiol*. 2005;28:153–8.
33. Mansueto G, D’Onofrio M, Iacono C, Rozzanigo U, Serio G, Procacci C. Gastroduodenal artery stump haemorrhage following pylorus-sparing Whipple procedure: treatment with covered stents. *Dig Surg*. 2002;19:237–40.
34. Miura F, Takada T, Ochiai T, Asano T, Kenmochi T, Amano H, et al. Aortic occlusion balloon catheter technique is useful for uncontrollable massive intraabdominal bleeding after hepato-pancreato-biliary surgery. *J Gastrointest Surg*. 2006;10:519–22.
35. Farley DR, Schwall G, Trede M. Completion pancreatectomy for surgical complications after pancreaticoduodenectomy. *Br J Surg*. 1996;83:176–9.
36. de Castro SM, Kuhlmann KF, Busch OR, van Delden OM, Lameris JS, van Gulik TM, et al. Delayed massive hemorrhage after pancreatic and biliary surgery: embolization or surgery? *Ann Surg*. 2005;241:85–91.