CORRECTION

Correction to: Basophils activated via TLR signaling may contribute to pathophysiology of type 1 autoimmune pancreatitis

Masato Yanagawa¹ · Kazushige Uchida¹ · Yugo Ando¹ · Takashi Tomiyama¹ · Takashi Yamaguchi¹ · Tsukasa Ikeura¹ · Toshiro Fukui¹ · Akiyoshi Nishio¹ · Yoshiko Uemura² · Takayuki Miyara³ · Hiroyuki Okamoto⁴ · Souhei Satoi⁵ · Kazuichi Okazaki¹

Published online: 26 February 2018
© Japanese Society of Gastroenterology 2018

Correction to: J Gastroenterol

<https://doi.org/10.1007/s00535-017-1390-6>

In the original publication of this article, the Table 2 was published incorrectly.

The revised Table 2 is given in the following page.

The original article can be found online at <https://doi.org/10.1007/s00535-017-1390-6>.

✉ Kazuichi Okazaki
okazaki@hirakata.kmu.ac.jp

¹ The Third Department of Internal Medicine, Division of Gastroenterology and Hepatology, Kansai Medical University, 2-5-1 Shinmachi, Hirakata, Osaka 573-1010, Japan

² Department of Pathology, Kansai Medical University, Hirakata, Japan

³ The First Department of Internal Medicine, Division of Hematology, Respiratory Medicine and Rheumatology, Kansai Medical University, Hirakata, Japan

⁴ Department of Dermatology, Kansai Medical University, Hirakata, Japan

⁵ Department of Surgery, Kansai Medical University, Hirakata, Japan

Table 2 TLR expression on basophils infiltrated into the pancreatic tissue with type 1 AIP

Case no.	Age	Gender	TLR1	TLR2	TLR3	TLR4	TLR5	TLR6	TLR7	TLR8	TLR9
1	63	M	–	+	–	–	–	–	–	–	–
2	71	F	–	+	–	–	–	–	–	–	–
3	78	F	–	+	–	+	–	–	–	–	–
4	67	M	–	+	–	+	–	–	–	–	–
5	67	M	–	–	–	+	–	–	–	–	–
6	56	M	–	–	–	+	–	–	–	–	–
7	57	F	–	–	–	+	–	–	–	–	–
8	60	F	–	–	–	+	–	–	–	–	–
9	60	F	–	–	–	+	–	–	–	–	–
10	71	M	–	–	–	+	–	–	–	–	–