



Correction to: The intraoperative motor-evoked potential when propofol was changed to remimazolam during general anesthesia: a case series

Shoto Yamada¹ · Yukinori Akiyama² · Shunsuke Tachibana³ · Kengo Hayamizu³ · Yusuke Kimura² · Shuichi Hashimoto¹ · Michiaki Yamakage³ · Nobuhiro Mikuni²

Published online: 30 November 2022

© The Author(s) under exclusive licence to Japanese Society of Anesthesiologists 2022

Correction to: Journal of Anesthesia

<https://doi.org/10.1007/s00540-022-03112-0>

In the original publication of the article, affiliations were published incorrectly. The correct affiliations are provided in this article.

The correct affiliations are,

1. Division of Clinical Engineering, Sapporo Medical University School of Medicine, Sapporo, Japan.

2. Department of Neurosurgery, Sapporo Medical University School of Medicine, Sapporo, Japan.

3. Department of Anesthesiology, Sapporo Medical University School of Medicine, Sapporo, Japan.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00540-022-03112-0>.

✉ Yukinori Akiyama
akiyuki@sapmed.ac.jp

¹ Division of Clinical Engineering, Sapporo Medical University School of Medicine, Sapporo, Japan

² Department of Neurosurgery, Sapporo Medical University School of Medicine, Sapporo, Japan

³ Department of Anesthesiology, Sapporo Medical University School of Medicine, Sapporo, Japan