



Letter to the article by Juri T, et al.

Betul Kozanhan¹ · Munise Yıldız¹ · Mahmut Sami Tutar¹

Received: 14 February 2024 / Accepted: 18 February 2024
© The Author(s) under exclusive licence to Japanese Society of Anesthesiologists 2024

To the Editor:

We have read with great interest the recent study “Changes in the corrected carotid flow time can predict spinal anesthesia-induced hypotension in patients undergoing cesarean delivery: an observational study”, published in the *J Anesth* [1]. However, the article overlooks a critical aspect.

The study mentions that carotid ultrasonography was performed unilaterally. Since carotid arteries are symmetrical structures in the body and blood flow on each side can vary due to different physiological or pathological conditions, conducting measurements on both sides would provide a more comprehensive evaluation.

Bilateral measurement, endorsed by established guidelines, yields a more thorough dataset by considering variations and potential pathologies between both carotid arteries. Such a detailed approach is crucial for precise clinical evaluations and research outcomes [2]. The methodology section of the study does not convincingly justify its preference for measuring one side, a crucial detail to prevent bias. A transparent explanation, whether for superior acoustic windows or a specific criterion like consistently selecting the right side, is essential for the study’s credibility and repeatability.

The study’s dependability is compromised by not providing a solid rationale for bypassing bilateral measurements,

potentially leading to misleading clinical interpretations. We urge the authors to reflect on these methodological issues and encourage future studies to adopt a bilateral approach or thoroughly explain any unilateral choices.

Funding No.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

References

1. Juri T, Suehiro K, Yasuda S, Kimura A, Fujimoto Y, Mori T. Changes in the corrected carotid flow time can predict spinal anesthesia-induced hypotension in patients undergoing cesarean delivery: an observational study. *J Anesth*. 2024;38(1):108–13.
2. Oates CP, Naylor AR, Hartshorne T, Charles SM, Fail T, Humphries K, Aslam M, Khodabakhsh P. Joint recommendations for reporting carotid ultrasound investigations in the United Kingdom. *Eur J Vasc Endovasc Surg*. 2009;37(3):251–61.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This comment refers to the article available online at <https://doi.org/10.1007/s00540-023-03293-2>.

✉ Mahmut Sami Tutar
masatu42@gmail.com
Betul Kozanhan
betulkozan@gmail.com
Munise Yıldız
drmunise@hotmail.com

¹ Department of Anesthesiology and Reanimation, Konya City Hospital, University of Health Sciences, Akabe, Adana Çevre Yolu Cd. No: 135/1, Karatay, 42020 Konya, Turkey