Ectomycorrhizal fungal communities are dominated by mammalian dispersed truffle-like taxa in north-east Australian woodlands

S. J. Nuske^{1,2} · S. Anslan³ · L. Tedersoo⁴ · B. C. Congdon⁵ · S. E. Abell¹

Received: 7 November 2018 / Accepted: 5 March 2019 / Published online: 21 March 2019
© The Author(s) 2019

Abstract

Mycorrhizal fungi are very diverse, including those that produce truffle-like fruiting bodies. Truffle-like fungi are hypogeous and sequestrate (produced below-ground, with an enclosed hymenophore) and rely on animal consumption, mainly by mammals, for spore dispersal. This dependence links mycophagous mammals to mycorrhizal diversity and, assuming truffle-like fungi are important components of mycorrhizal communities, to plant nutrient cycling and ecosystem health. These links are largely untested as currently little is known about mycorrhizal fungal community structure and its dependence on mycophagous mammals. We quantified the mycorrhizal fungal community in the north-east Australian woodland, including the portion interacting with ten species of mycophagous mammals. The study area is core habitat of an endangered fungal specialist marsupial, *Bettongia tropica*, and as such provides baseline data on mycorrhizal fungi-mammal interactions in an area with no known mammal declines. We examined the mycorrhizal fungi in root and soil samples via high-throughput sequencing and compared the observed taxa to those dispersed by mycophagous mammals at the same locations. We found that the dominant root-associating ectomycorrhizal fungal taxa (> 90% sequence abundance) included the truffle-like taxa *Mesophellia*, *Hysterangium* and *Chondrogaster*. These same taxa were also present in mycophagous mammalian diets, with *Mesophellia* often dominating. Altogether, 88% of truffle-like taxa from root samples were shared with the fungal specialist diet and 52% with diets from generalist mammals. Our data suggest that changes in mammal communities, particularly the loss of fungal specialists, could, over time, induce reductions to truffle-like fungal diversity, causing ectomycorrhizal fungal communities to shift with unknown impacts on plant and ecosystem health.

Keywords Sequestrate fungi · Truffle-like fungi · Mycophagy · Ecosystem interactions · *Bettongia tropica* · Ectomycorrhizal fungi

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00572-019-00886-2>) contains supplementary material, which is available to authorized users.

✉ S. J. Nuske
susan.nuske@slu.se

¹ College of Science and Engineering, Centre for Tropical Environmental and Sustainability Science, Australian Tropical Herbarium, James Cook University, Cairns, QLD 4878, Australia

² Present address: Department of Forest Ecology and Management, Swedish University of Agricultural Sciences, 90183 Umeå, Sweden

³ Zoological Institute, Braunschweig University of Technology, Mendelssohnstr. 4, 38106 Braunschweig, Germany

⁴ Natural History Museum and Institute of Ecology and Earth Sciences, University of Tartu, 14A Ravila, EE-50411 Tartu, Estonia

⁵ College of Science and Engineering, Centre for Tropical Environmental and Sustainability Science, James Cook University, Cairns, QLD 4878, Australia

Introduction

Hypogeous, sequestrate (truffle-like) mycorrhizal fungi are an important component of forest ecosystems and they rely on animals, particularly mammals, for their spore dispersal (Claridge and May 1994). This implies that truffle-like fungal diversity is likely linked to mammal diversity (Vernes 2007). Disruption to complex ecological networks, such as this mammal-fungi-plant interaction, can cause loss of biodiversity. For example, it is logical to assume that reduced spore dispersal via loss of mammal abundance and diversity would reduce gene flow among truffle-like populations, resulting in undocumented impacts on truffle-like fungal community structure and potential species extinctions. Loss of truffle-like species diversity may in turn alter mycorrhizal communities, potentially impacting fungi-plant interactions.

Mammals are also thought to play a pivotal role in plant-mycorrhiza symbioses, and by extension, plant productivity, diversity and ecosystem health (Maser et al. 1978; Malajczuk et al. 1987; Johnson 1996; Vernes 2007). Such a hypothesis assumes that truffle-like taxa are important components of functioning mycorrhizal communities and that the higher the proportion of truffle-like taxa within the overall mycorrhizal community, either in terms of relative abundance or diversity, the higher the potential influence that mammalian spore dispersal has on the structuring of mycorrhizal and plant communities. Yet, these assumed linkages remain largely untested.

To understand the strength of the relationship between mycorrhizal communities and mycophagous mammals, first we must understand how important truffle-like taxa are to functioning mycorrhizal communities. Few studies have identified the fruiting habits of the various components of the Australian mycorrhizal fungal community (or presented data with enough resolution that this can be inferred post hoc; Online Resource 1, Table S1). In three different studies of ectomycorrhizal (ECM) fungal sporocarps in Australia (Reddell et al. 1999; Lu et al. 1999; Adams et al. 2006), between 18 and 27% of taxa found were truffle-like (reported as hypogeous). However, these three surveys are difficult to compare in terms of the richness of hypogeous versus epigeous species because the same methodology was not used for both groups.

It is also important to make the distinction between ECM and arbuscular mycorrhizal (AM) communities as the relative diversity of truffle-like species are quite different in these groups as are their distribution, ecology and interactions with host plants. AM fungi associate with > 80% of global plant diversity and occur in almost every ecosystem where plants are present, while ECM associate with a much smaller proportion (Brundrett 2009). However, ECM trees can dominate forests in terms of biomass (Reddell et al. 1999). The diversity of truffle-like (sporocarpic) AM fungi is much lower compared to ECM fungi; at least two AM genera contain truffle-like species (*Glomus* and *Acaulospora*; Goto and Maia 2005) while thousands of species of ECM truffle-like fungi are within Basidiomycota, Ascomycota and Zygomycota (Bougher and Lebel 2001; Trappe et al. 2009). Therefore, any influence that mammals may have on mycorrhizal communities will likely depend on the differences between these groups.

To our knowledge, only one study has quantified the ECM community on natural forest plant host roots and compared this to mycophagous mammalian diets. Izzo et al. (2005) sampled roots from subtropical North America and found at least 21% of taxa were truffle-like and between 25 and 40% of ECM dry root biomass were truffle-like taxa (reported as hypogeous). However, this study was limited to Sanger sequencing of DNA samples from mammalian scats and consequently detected only three truffle-like taxa in mammalian diets. Modern high-throughput sequencing technologies that amplify and sequence DNA from complex communities provide the necessary

resolution to examine fungal communities from environmental samples like roots, scats and soil (Lindahl et al. 2013).

Our aim was to address the knowledge gap in the structure and relative proportion of mycorrhizal communities, particularly the truffle-like communities that are interacting with mycophagous mammals. Using high-throughput sequencing (Illumina MiSeq), we compared the mycorrhizal fungal communities (concentrating on ECM fungi) across three sample types including plant roots, soil and scats from mycophagous mammals. We compared our results to reference ITS2 sequences of truffle-like morpho-species collected and characterised from an extensive survey undertaken at one of our sampling sites (Abell-Davis 2008). We collected roots, soil and mammalian scat samples within the habitat of a fungal specialist (the northern bettong; *Bettongia tropica*) and other mycophagous mammals as this provides a baseline measurement of the interaction between root-associating mycorrhizal fungi and mycophagous mammals where limited known loss of mammal diversity has occurred.

Methods

Field sampling

Sampling was carried out on the Lamb Range in North Queensland in Australia at Davies Creek National Park (17°1'23.28"S, 145°34'55.71"E; elevation 600–730 m) in the late dry season (November to December) in 2014. Additional sampling was carried out at two locations in the early wet season (February to March): Danbulla National Park near Tinaroo Dam (17°9'50.30"S, 145°32'11.56"E; elevation 630–780 m) and Davies Creek in 2015. Six plots (12 × 20 m) at each site were established at least 500 m apart around the animal trapping grid (as used in Nuske et al. 2018) for the collection of soil and root samples. The dominant and putatively ectomycorrhizal tree species were *Eucalyptus crebra*, *E. tindaliae*, *E. mediocris*, *Corymbia intermedia*, *Allocasuarina littoralis*, *Al. torulosa* and *Acacia flavescens*.

Soil was collected from 40 cores (0–10 cm deep, 5 cm diameter) per plot. The corers were carefully cleaned with 70% ethanol between plots. Half of the plot (6 × 20 m) was also used to collect putative ECM root-associated taxa by raking the top 10 cm of soil for 60 person-minutes and collecting fine roots. Because the soil was often too compacted and contained large rocks, we were not able to trace fine roots back to the potential host plants. Instead, we collected all fine root material found. When possible, grass roots were eliminated from individual samples by tracing back to the grass plant. Preliminary work also suggested that a higher volume of roots was collected with this more targeted approach compared to sieving roots from soil cores. A high volume of root tip material was necessary to obtain three DNA extraction subsamples per plot (3 × 0.25 g

of wet weight root tips, see below). Soil and roots were frozen (-4°C) within 24 h of collection and placed into -20°C as soon as possible (up to 4 days).

Fungal diets of mammals were examined by collecting scat samples from trapped individuals. Trapping and animal handling protocols were as outlined in Nuske et al. (2018). Briefly, medium-sized and small mammals trapped for four consecutive nights each, at two locations and two seasons. We identified mammals according to Van Dyck et al. (2013). We marked each individual by either removing a small patch of hair with scissors at the base of the tail or microchipping (*B. tropica* only; Minichips, Micro Products Australia, Canning Vale, WA or ISO FDX-B Microchips, OzMicrochips, Peakhurst, NSW). Scats were collected from the bottom of Elliot traps or from plastic placed under each cage trap. All traps and plastic were initially cleaned with 70% ethanol and then re-cleaned if an animal was caught. Scats were stored on ice or in a portable fridge (4°C) in the field and transferred to -20°C as soon as possible (within 4 days). Each animal was handled according to James Cook University animal ethical guidelines (Approved ethics application A2044).

Laboratory

Roots were cleaned of excess soil in reverse osmosis water. Fine roots were examined under a dissecting microscope. For each cluster of fine roots collected within a plot, the same volume of root tips of each mycorrhizal morphotype were placed into three subsamples (0.25 g) for DNA extraction. We did not attempt to verify the colonisation of each ectomycorrhizal morphotype to maintain efficient processing.

Therefore, we consider the mycorrhizal community from root samples to be ‘root-associated’. These taxa are more likely to represent functioning mycorrhizal fungi (i.e. those taxa colonised and interacting with plants) compared to that sequenced from soil samples. For soil samples, each of the 40 soil cores per plot were homogenised and pooled. Then three subsamples of fine powdered soil were taken per plot for DNA extraction (0.25 g). We only used scats from the first capture of an individual per trapping session. Each bolus of scat was broken in half and a small sample of faecal material removed from the inside with sterile forceps. Samples were homogenised and 0.25 g was taken for DNA extraction.

DNA extraction, PCR, sequencing protocols and bioinformatics were processed as in Nuske et al. (2018). Briefly, DNA was extracted using PowerLyser PowerSoil DNA Isolation kit (Mo Bio, Carlsbad, CA USA) following manufacturer’s instructions, except that the samples were lysed using a Qiagen Tissue Lyser for 2×30 s at 30 Hz, swapping the position of the samples between runs. DNA was amplified with ITS3-Mix1-5 (5'-CTAGACTCGTCANCGATGAAGAACGYRG-3') and barcoded ITS4ngs (5'-TCCTSCGCTTATTG ATATGC-3') primers (Tedersoo et al. 2014). The primers were tagged with 10–11 base unique molecular identifiers (MIDs) to later distinguish samples with sequencing runs (Online Resource 1, Table S1). We used negative (for DNA extraction and PCR) and positive controls (PCR) throughout the experiment. Normalised amplicons were subjected to ligation of Illumina adaptors using the TruSeq DNA PCR-free HT Sample Prep kit (Illumina Inc., San Diego, CA, USA). All samples were sequenced using Illumina MiSeq 2×300 paired-end technology. Raw Illumina data is deposited in Sequence Read Archive (SRA; bioproject SRP150847).

Fig. 1 Rarefied accumulation curve $\pm 95\%$ confidence intervals for **a** mycorrhizal OTUs and **b** truffle (truffle-like/sequestate) OTUs. Orange lines are soil samples ($n = 18$, each containing three pooled samples), light blue lines are root samples ($n = 18$, each containing three pooled samples) and dark blue lines are scat samples ($n = 61$)

Bioinformatics were performed using PipeCraft (v1.0; Anslan et al. 2017) as per Nuske et al. (2018). Representative sequences for each operational taxonomic unit (OTU, clustered with CD-HIT at 97% similarity threshold, v4.6; Li and Godzik 2006) were chosen using the mothur ‘abundance’ method and compared against UNITE (v7.0), GenBank ITS and our local truffle-like fungal database (Nuske et al. 2018) to obtain taxonomic affiliation using BLASTn (Camacho et al. 2009). Taxonomic groups were assigned to functional categories using FUNGuild (v1.0; Nguyen et al. 2015a). All Glomeromycota taxa were assigned as arbuscular mycorrhizal.

Statistics

OTU subsetting and statistics were performed using the ‘phyloseq’ package (McMurdie and Holmes 2013) in R (R Core Team 2012). Altogether, six samples were removed from further analyses, because these comprised < 500 filtered sequences. The fungal dataset was examined at three broad levels: at the whole OTU community level, only examining the mycorrhizal OTUs (8.4% of all taxa) and only examining truffle-like taxa (9.3% of mycorrhizal taxa). The mycorrhizal subset of the data included only taxa that were assigned as ‘highly probable’ and ‘probable’ from the FUNGuild output (Nguyen et al. 2015b). Functional guilds are assigned to ECM status by FUNGuild based on genera, with the exception of Russulaceae. Russulaceae is also one of the most OTU-rich families in this dataset; therefore, when comparing mycorrhizal taxa (e.g. as relative richness of mycorrhizal families), Russulaceae are disproportionally over-represented. To make sure ECM taxa were evenly represented at the *genus* level, ECM OTUs assigned to family level, Russulaceae, were excluded from analyses (note: ECM OTUs assigned to genus and species level within Russulaceae, for example *Russula*, were retained). Identified truffle-like taxa are listed in Online Resource 2.

The three subsamples per soil or root sample were pooled computationally by mean sequence abundance per OTU as this likely gives the best estimate of richness (Song et al. 2015). To estimate the accumulation of OTUs per sample, we created rarefaction OTU accumulation curves for each sample type with mycorrhizal and truffle OTU data using the ‘ggrare’ function (richness.R, phyloseq extensions; <https://github.com/mahendra-mariadassou/phyloseq-extended> with added ggplot2 graphics).

We compared mycorrhizal communities between sample types by tabulating the number of OTUs that were shared and unshared (using ‘limma’ and ‘venneuler’ packages in R). We ranked OTUs according to their relative abundance per sample (based on sequence count) and considered the ‘dominant’ portion of the community to be the highest relative abundance that collectively accounted for > 90% of the

relative abundance. The dominant portion accounted for approximately 25% of the taxa present.

Results

Richness

Soil samples were more OTU-rich than root samples, and both were more OTU-rich than scat samples (6689, 3226 and 2932 OTUs, respectively). There were a total of 9358 filtered OTUs across all samples. Most were not assigned to a functional guild (7508 OTUs) by FUNGuild. Of those that were, most were symbiotrophic (805 OTUs, including 344 ECM OTUs and 428 AM OTUs) followed by saprotrophic (754 OTUs). For mycorrhizal OTUs, at equivalent read count, the OTU richness of scats was ~60% of the root-associating taxa (Fig. 1a). We detected a higher truffle-like OTU richness in scat samples than in root or soil samples (Fig. 1b).

Fig. 2 Relative OTU richness of mycorrhizal families in different samples (soil, roots and scats). Only taxa representing greater than 1% of the total OTU richness are shown for clarity

Both soil and root samples had AM fungi (Glomeromycota, Glomeraceae and Gigasporaceae) as the most OTU-rich mycorrhizal taxa. However, they made up < 13% of the taxa richness in the mammalian scat samples (Fig. 2, including Diversisporaceae) and constituted < 0.01% of the relative abundance for all samples (Fig. 3). The truffle-like family Mesophelliaceae was the most OTU-rich and highest relative abundance mycorrhizal taxon in mammalian scat samples (Figs. 2 and 3).

Overall, samples had 21.2% of ECM OTUs as truffle-like taxa (73 OTUs) and 54.1% as taxa with ambiguous fruiting habit (186 OTUs). Scat samples had a higher proportion of OTUs matching truffle-like fungi than root or soil samples (30–90%, depending on site and season, compared to 17–23% for root samples and 7–8% for soil samples). Scat samples from the fungal specialist (*B. tropica*) had a higher proportion of ECM truffle-like OTUs (30–90%) than all other mammal species with generalist diets combined (18–73%).

Relative abundance

Truffle-like and secotioid taxa (e.g. Cortinariaceae and Hysterangiaceae) constituted higher proportions of the dominant mycorrhizal root-associating communities compared to soil communities (Table 1; Fig. 3). Soil mycorrhizal communities were dominated by taxa with mixed fruiting habits (e.g. truffle-like taxa and mushroom taxa and *Russula* and *Cortinarius* that could represent truffle-like, secotioid or mushroom taxa; Table 1 and Fig. 3). These patterns of relative

abundance were consistent across the two sites and two seasons we measured (Table 1). Of the dominant mycorrhizal OTUs associating with root samples, three out of four genera were truffle-like from Hysterangiales (*Hysterangium*, *Mesophellia* and *Chondrogaster*; Table 1). Four truffle-like OTUs were shared between dominant root taxa and mammalian diets and five dominant ECM OTUs were shared between soil and scat samples (Table 1). Within taxa, comparisons showed that *Cortinarius* was more OTU-rich and more relatively abundant in roots and soil compared to scat samples, whereas *Malajczukia* and *Mesophellia* (Mesophelliaceae) were more OTU-rich and abundant in scat samples compared to soil and roots (Table 1; Fig. 2). *Hysterangium* (Hysterangiaceae) and *Chondrogaster* (Mesophelliaceae) were relatively more abundant in roots.

Shared taxa between samples

The percentage of shared taxa between fungal specialist diets and ECM communities on roots (36.6%) and in soil (28%) was slightly higher than that for fungal generalist diets (26 and 14.7%, respectively; Fig. 4). The percentage of shared truffle-like OTUs from roots and soil and fungal specialist diets was much higher (87.5 and 78.8%, respectively; Fig. 4). In contrast, just over half of the truffle-like taxa from root and soil samples overlapped with fungal generalist diets (52–53%). Arbuscular mycorrhizal (AM) OTU richness was highest from soil samples and AM communities from roots did not overlap significantly with mammalian scat samples (< 1.8%; Fig. 4).

Fig. 3 Relative abundance of mycorrhizal families in different samples (soil, roots and scats) split by mycorrhizal type (AM = arbuscular mycorrhizal) and fruiting habit (*n* = fruiting habit other than truffle-like

taxa, *n/y* = unknown fruiting habit, *y* = truffle-like/sequestrate taxa). Black bars are overlapping boundary lines representing OTUs with a relative abundance too small to display

Table 1 Mycorrhizal OTUs that make up the dominant proportion (cumulatively 90% of sequence abundance) of samples per site (DC = Davies Creek, TD = Tinaroo Dam), sample type and season, with number of replicates (*n*), total OTUs per sample group (total), percent relative abundance (RA), accession number, e-value, percentage similarity with database sequence (ID), percentage overlap with reference sequence (Cov) and OTU sequence length (SL). Taxa in boldface are truffle-like/sequestrate and underlined are secotoid or higher taxa that include

truffle-like/sequestrate or secotoid taxa. When fruiting habit is listed at genus level, it applies to the whole genus. Mycorrhizal status (Myc) is either ectomycorrhizal (ECM), ericoid mycorrhizal (ErM) or arbuscular mycorrhizal (AM). The mammalian specialist scats are from *Bettongia tropica*. The mammalian generalist scats are from *Isoodon macrourus*, *I. obesulus*, *Melomys* sp., *Trichosurus vulpecula*, *Uromys caudimaculatus* and *Zyzomys argurus*

Site	Sample	Season	n	Total	RA	Fungal taxa	Myc	Accession	e-value	ID	Cov	SL
DC	roots	dry	6 ^b	84	64.4	<i>Hysterangium aggregatum</i>	ECM	KY697566-7	2.21E-133	100	97.9	333
					17.2	<u><i>Cortinarius globuliformis</i></u>	ECM	AF325582	2.50E-141	99	100	350
					5.7	<i>Hysterangium aggregatum</i>	ECM	KY697566-7	2.16E-129	98	97.9	335
					5.3	<u><i>Cortinarius globuliformis</i></u>	ECM	AF325582	3.24E-142	99	100	349
					44.1	<u><i>Cortinarius</i></u>	ECM	FR731477	1.53E-143	100	99.7	350
	roots	wet	6 ^b	102	12.5	<i>Hysterangium</i> ^c	ECM	KC222660	1.00E-144	94	100	340
					11.8	<i>Hysterangium</i> cf <i>gardneri</i> ^a	ECM	KY697590	3.96E-139	100	99.7	340
					11.4	<i>Hysterangium aggregatum</i>	ECM	KY697566-7	2.21E-133	100	97.9	333
					7.1	<i>Mesophellia oleifera</i> ^c	ECM	KY697602-3	4.15E-177	100	100	425
					3.9	<u><i>Cortinarius</i></u>	ECM	KJ421051	4.44E-112	91	100	358
TD	roots	wet	6 ^b	99	53.1	<u><i>Cortinarius</i></u>	ECM	FR731477	1.53E-143	100	99.7	350
					12.5	<i>Mesophellia</i> ^c	ECM	GQ981511	2.00E-111	91	98	298
					12.4	<u><i>Cortinarius</i></u>	ECM	KF732610	5.34E-105	89	100	353
					9.3	<i>Mesophellia oleifera</i> ^c	ECM	KY697602-3	4.86E-166	97	100	420
					4.4	<i>Chondrogaster spB/spF</i> ^a	ECM	KY697582-5	4.75E-151	100	100	366
DC	specialist diet	dry	7	51	31.0	<i>Mesophellia</i> ^{c,d}	ECM	GQ981511	2.00E-111	91	98	298
					19.5	<i>Malajcukia ingrattissima</i> ^d	ECM	KY697598	6.60E-156	100	100	377
					19.3	<i>Mesophellia oleifera</i> ^c	ECM	KY697602-3	4.15E-177	100	100	425
					17.1	<i>Mesophellia glauca</i>	ECM	GQ981510	9.13E-147	98	99	376
					5.8	<i>Mesophellia glauca</i>	ECM	GQ981511	1.00E-162	97	99	354
	specialist diet	wet	6	24	34.7	<u><i>Russula</i></u> ^d	ECM	LC006943	3.99E-138	91	100	426
					32.9	<i>Mesophellia</i> ^{c,d}	ECM	GQ981511	2.00E-111	91	98	298
					17.5	<u><i>Russula</i></u>	ECM	UDB016041	6.92E-129	93	100	376
					7.0	<u><i>Cortinarius</i></u>	ECM	FJ157098	8.08E-118	92	100	371
					26.4	<i>Malajcukia ingrattissima</i> ^d	ECM	KY697598	6.60E-156	100	100	377
TD	specialist diet	wet	16	70	16.6	<i>Mesophellia</i> ^{c,d}	ECM	GQ981511	2.00E-111	91	98	298
					15.2	<i>Mesophellia</i>	ECM	GQ981511	2.00E-110	91	99	295
					12.8	<i>Mesophellia glauca</i>	ECM	GQ981510	9.13E-147	98	99	376
					7.2	<u><i>Russula</i></u> ^d	ECM	LC008293	3.95E-138	91	100	422
					6.1	<i>Scleroderma spB/spC</i> ^{a,d}	ECM	KY697606-7	3.41E-146	100	100	355
	generalist diets	dry	11	27	3.4	<i>Mesophellia oleifera</i> ^c	ECM	KY697602-3	4.86E-166	97	100	420
					3.0	<i>Mesophellia oleifera</i> ^c	ECM	KY697602-3	4.15E-177	100	100	425
					68.8	<i>Malajcukia ingrattissima</i> ^d	ECM	KY697598	6.60E-156	100	100	377
					18.7	<i>Mesophellia</i> ^{c,d}	ECM	GQ981511	2.00E-111	91	98	298
					9.6	<i>Mesophellia</i>	ECM	GQ981511	2.00E-110	91	99	295
DC	generalist diets	wet	8	29	35.3	<i>Mesophellia</i>	ECM	GQ981511	2.00E-110	91	99	295
					30.0	<u><i>Russula</i></u> ^d	ECM	LC006943	3.99E-138	91	100	426
					16.8	<u><i>Russula</i></u>	ECM	UDB016041	6.92E-129	93	100	376
					11.5	<i>Lactarius rufus</i>	ECM	KT165272	4.86E-170	100	100	409
					71.4	<i>Rhizopogon pseudoroseolus</i>	ECM	AJ810040	2.84E-168	100	100	405
TD	generalist diets	wet	8	24	9.2	<i>Mesophellia</i> ^{c,d}	ECM	GQ981511	2.00E-111	91	98	298
					7.9	<i>Malajcukia ingrattissima</i> ^d	ECM	KY697598	6.60E-156	100	100	377
					2.6	<i>Hysterangium</i> ^c	ECM	KC222660	1.00E-144	94	100	341

Table 1 (continued)

Site	Sample	Season	n	Total	RA	Fungal taxa	Myc	Accession	e-value	ID	Cov	SL
DC	soil	dry	6 ^b	196	22.4	<i>Russula</i>	ECM	UDB016041	1.08E-122	92	100	370
					19.9	<i>Russula</i> ^d	ECM	LC006943	3.99E-138	91	100	426
					11.5	<i>Russula</i>	ECM	UDB016041	4.05E-127	92	100	373
					6.4	<i>Russula anthracina</i>	ECM	UDB011194	5.00E-151	97	100	382
					4.9	<i>Amanita</i>	ECM	KP071067	2.94E-126	94	100	356
					3.6	<i>Inocybe</i>	ECM	FJ904133	1.39E-100	88	100	344
					3.5	<i>Hysterangium aggregatum</i>	ECM	KY697566-7	2.21E-133	100	97.9	333
					3.1	<i>Russula</i>	ECM	EU019930.1	1.63E-104	93	80.4	382
					2.7	<i>Inocybe</i>	ECM	JQ085932	1.01E-91	87	100	346
					2.3	<i>Mesophellia</i> ^d	ECM	GQ981511	2.00E-111	91	98	298
					2.1	<i>Lactarius</i>	ECM	HQ318282	2.43E-129	94	100	368
					1.3	<i>Inocybe</i>	ECM	JX178624	2.40E-98	86	100	359
					1.3	<i>Auritella serpentinocystis</i>	ECM	KJ729858	3.63E-150	100	100	364
					1.2	<i>Cortinarius globuliformis</i>	ECM	AF325582	2.50E-141	99	100	350
					0.9	<i>Russula</i>	ECM	UDB016041	2.40E-125	92	100	373
					0.8	<i>Hysterangium aggregatum</i>	ECM	KY697566-7	2.60E-122	97	97.9	330
					0.8	<i>Russula</i>	ECM	AB509981	3.49E-138	94	99.7	380
					0.7	<i>Cortinarius</i>	ECM	KR011131	2.92E-122	93	100	363
					0.6	<i>Cortinarius globuliformis</i>	ECM	AF325582	3.24E-142	99	100	349
	soil	wet	6 ^b	251	24.8	<i>Russula</i>	ECM	UDB016041	4.05E-127	92	100	373
					13.7	<i>Cortinarius</i>	ECM	GU233352	1.41E-96	88	100	357
					9.6	<i>Russula</i> ^d	ECM	LC006943	3.99E-138	91	100	426
					8.0	<i>Auritella chamaecephala</i>	ECM	KT378201	9.00E-138	97	100	358
					7.5	<i>Inocybe</i>	ECM	JQ085932	1.01E-91	87	100	346
					4.7	<i>Cortinarius</i>	ECM	KJ421051	4.44E-112	91	100	358
					4.5	<i>Malajcukia ingrattissima</i> ^d	ECM	KY697598	6.60E-156	100	100	377
					2.5	<i>Russula</i>	ECM	UDB016041	1.08E-122	92	100	370
					2.3	<i>Lactarius</i>	ECM	HQ318282	2.43E-129	94	100	368
					1.8	<i>Mesophellia</i> ^d	ECM	GQ981511	2.00E-111	91	98	298
					1.5	<i>Amanita</i>	ECM	JF899547	4.64E-112	89	100	371
					1.1	<i>Cantharellus</i>	ECM	AB509732	8.03E-106	85	99.7	398
					0.9	<i>Russula</i>	ECM	UDB016041	3.12E-126	92	100	373
					0.8	<i>Auritella serpentinocystis</i>	ECM	KJ729858	3.63E-150	100	100	364
					0.7	<i>Russula</i>	ECM	KM373243	3.51E-134	93	100	391
					0.7	<i>Hysterangium aggregatum</i>	ECM	KY697566-7	2.21E-133	100	97.9	333
					0.6	<i>Scleroderma spB/spC</i> ^{a,d}	ECM	KY697606-7	3.41E-146	100	100	355
					0.5	<i>Oidiodendron</i>	ErM ^c	AF062808.1	1.51E-105	95	100	291
					0.5	<i>Glomeromycetes</i>	AM	JF276264	4.84E-128	96	100	348
					0.4	<i>Auritella</i>	ECM	KT378201	4.51E-116	92	100	354
					0.4	<i>Glomerales</i>	AM	AY394681	3.63E-76	81	100	380
					0.3	<i>Glomerales</i>	AM	HE794042	2.00E-117	93	100	338
					0.3	<i>Glomeraceae</i>	AM	KM226647	1.20E-115	97	88.6	343
					0.3	<i>Inocybe</i>	ECM	JX178624	2.40E-98	86	100	359
					0.3	<i>Glomerales</i>	AM	KP235575	1.87E-101	90	97.5	354
					0.3	<i>Oidiodendron</i>	ErM ^c	KX640607	3.00E-131	96	99.7	289
					0.2	<i>Russula</i>	ECM	UDB016041	8.38E-122	91	99.5	373
					0.2	<i>Glomerales</i>	AM	JX276895	4.56E-124	95	100	340
					0.2	<i>Glomerales</i>	AM	KM226647	3.30E-80	86	88.6	343
					0.2	<i>Glomerales</i>	AM	AY394681	1.72E-81	82	100	374

Table 1 (continued)

Site	Sample	Season	n	Total	RA	Fungal taxa	Myc	Accession	e-value	ID	Cov	SL
TD	soil	wet	6 ^b	289	12.2	<i>Cortinarius globuliformis</i>	ECM	AF325582	6.96E-141	99	100	351
					11.5	<i>Cortinarius globuliformis</i>	ECM	AF325582	3.24E-142	99	100	349
					9.2	<i>Inocybe alienospora</i>	ECM	KP171105	1.88E-140	99	100	343
					6.9	<i>Cortinarius</i>	ECM	FR731477	1.53E-143	100	99.7	350
					6.5	<i>Amanita egregia</i>	ECM	KP012748	2.82E-134	100	100	328
					5.6	<i>Lactarius</i>	ECM	AB509713	4.60E-112	90	99.7	369
					5.0	<i>Inocybe</i>	ECM	KP308804	6.00E-135	89	94	359
					4.8	<i>Inocybe</i>	ECM	KP308804	3.05E-99	87	100	352
					4.4	<i>Lactarius eucalypti</i>	ECM	UDB002671	1.70E-162	96	100	420
					3.2	<i>Clavulina</i>	ECM	JQ724058	3.49E-103	85	100	383
					3.1	<i>Austroboletus subvirens</i>	ECM	KP242209	5.04E-155	100	100	375
					2.8	<i>Zelleromyces spE</i>	ECM	KY697617-9	8.92E-153	96	100	399
					2.6	<i>Lactifluus</i>	ECM	KM282287	1.36E-127	95	100	351
					2.1	<i>Inocybe</i>	ECM	AM882711	2.72E-99	90	100	321
					2.0	<i>Russula</i>	ECM	UDB016041	1.08E-122	92	100	370
					1.5	<i>Scleroderma spB/spC^{a,d}</i>	ECM	KY697606-7	3.41E-146	100	100	355
					1.4	<i>Russula^d</i>	ECM	LC006943	3.99E-138	91	100	426
					1.1	<i>Pisolithus croceorrhizus</i>	ECM	JN847473	8.64E-157	100	100	379
					0.8	<i>Inocybe violaceocaulis</i>	ECM	KP641643	4.75E-151	100	100	366
					0.8	<i>Amanita</i>	ECM	GU222312	3.39E-111	92	94.1	356
					0.6	<i>Pisolithus croceorrhizus</i>	ECM	JN847473	6.64E-156	99	100	379
					0.5	<i>Cortinarius globuliformis</i>	ECM	AF325582	3.25E-142	99	100	350
					0.5	<i>Amanita</i>	ECM	AB015702	8.29E-95	87	100	355
					0.5	<i>Glomerales</i>	AM	JN195694	5.02E-132	96	100	350
					0.5	<i>Mesophellia</i>	ECM	GQ981511	8.00E-116	92	97	301
					0.4	<i>Russula^d</i>	ECM	LC008293	3.95E-138	91	100	422

^a Indicates taxa that are indistinguishable from ITS2 sequences (within 3% similarity) from morphological groups identified in Abell-Davis (2008), Online Resource 2

^b Includes three subsamples per sample pooled computationally

^c Indicates OTUs that are shared between root and scat samples

^d Indicates OTUs that are shared between scat and soil samples

^e The ericaceous shrub, *Melichrus urceolatus*, was present at low abundance at this site

Almost all Hysterangiaceae, Mesophelliaceae and Tuberaceae truffle and truffle-like taxa sequenced from roots and soil were also recovered from mammalian scats (Fig. 5). Hysterangiaceae and Mesophelliaceae made up between 28 and 71% of the ECM sequence abundance of root samples (depending on site and season; Table 1). Of the families sequenced from root samples, over half (59%) of Russulaceae, 20% of Inocybaceae and 7% of Cortinariaceae were also in scat samples (Fig. 5).

Discussion

The hypothesis that mammal communities are important for plant-mycorrhizal relationships, and indirectly are also contributing to the health of mycorrhizal host trees and nutrient

cycling (Johnson 1996) assumes that mammal-dispersed truffle-like fungi are an important part of mycorrhizal communities. Our data support this assumption, at least in north-east Australian woodlands with a diverse mammalian community including the fungal specialist, *B. tropica*. Dominant components of the root-associating mycorrhizal community were ECM truffle-like taxa dispersed by mammals. In another study of tropical ECM communities, truffle-like taxa (Hysterangiales; *Hysterangium* and *Nothocastoreum*) were also found in the dominant portion as sporocarps and on roots (Reddell et al. 1999). This indicates that mammals can potentially have a substantial influence on the functioning ECM community. Additionally, the fungal specialist, *B. tropica*, has previously found to consume a higher diversity and more unique truffle-like taxa than the combined diets of fungal generalists in the same community (Nuske et al. 2018). Indeed,

most truffle-like taxa associating with roots were within the fungal specialists' diet. By dispersing a high diversity of ECM inocula, these mammals, particularly fungal specialists, appear to indirectly contribute to plant productivity and nutrient cycling in these ecosystems.

The hypothesis linking mycophagous mammals to plant health and ecosystem functioning also implies that if the mammal community were to be altered, then the inoculum available for new colonising roots is also altered (over time, potentially lowering truffle-like fungal diversity). If dispersal of these taxa were reduced via changes to the mammal community, then the ECM community is likely to experience major shifts, with unknown consequences for plant health and nutrient cycling. Within Australia, altered ECM communities and decreases in ECM colonisation rates have previously been associated with Eucalypt dieback and decreased crown health (Scott et al. 2012; Ishaq et al. 2013; Horton et al. 2013). Additionally, Australia has experienced high rates of mammal extinction and decline (Short and Smith 1994; Woinarski et al. 2015). Combined with our results, these observations raise concerns of major alterations of landscape-level ecosystem function, underscoring the need for further research. Firstly, further studies are needed to confirm whether this ECM community structure is typical for

Australian woodlands. Secondly, future studies should utilise areas where fungal specialists have recently gone extinct or a reduction of mammal diversity has occurred and compare to areas with higher mammal diversity to measure any changes in ECM communities. Thirdly, studies are needed to investigate the functional redundancy of ECM taxa between truffle-like and epigeous taxa for aspects that interact with plant health and nutrient cycling.

Many mycophagy studies have found spores of sporocarpic AM fungi (truffle-like) in mammalian diets, mostly *Glomus* spp. (Janos et al. 1995; Vernes and Dunn 2009; Nuske et al. 2017). Indeed, these spores have been shown to be viable by inoculating bioassay seedlings with scats containing AM spores (McGee and Baczocha 1994; Reddell et al. 1997). However, our data from an Australian sclerophyll forest show that mammalian diets do not overlap significantly with AM fungi associating with roots or the general soil environment. This indicates that, at least in terms of species diversity, mammal dispersal of AM spores does not have a significant effect on the structure of AM communities in this system. This does not discount other affects mammals may indirectly have on AM communities through physical disturbances or altering plant communities (Gehring and Whitham 1994; Gehring et al.

Fig. 4 Venn diagrams; the size of a circle represents the relative OTU richness of each sample type (soil, roots or scats) within each subsample of data (ectomycorrhizal = ECM, arbuscular mycorrhizal = AM, sequestrate/truffle-like = truffles and all OTUs). Overlapping areas represent the proportion of OTUs shared between sample types, whereas the non-overlapping areas represent OTUs unique to specified substrate. The total number of OTUs is 9358, 428 for AM fungi, 344 for

ECM fungi and 116 for sequestrate/truffle-like fungi. Specialist scats are from fungal specialist *Bettongia tropica*, and generalist scats are from *Isoodon macrourus*, *I. obesulus*, *Melomys* sp., *Trichosurus vulpecula* (ECM only) and *Zyzomys argurus*. Note: for AM fungi, scats and roots shared 1.8% of taxa and scats and soil shared 0.8% of taxa and this is not shown because there were no taxa shared by all samples

Fig. 5 Venn diagrams; the size of a circle represents the relative OTU richness of each sample type (soil, roots or scats) within each family of fungi (Hysterangiaceae [26 OTUs], Mesophelliaceae [59 OTUs] and Tuberaceae [15 OTUs] contain only truffle and truffle-like/sequestate species; Cortinariaceae [128 OTUs], Russulaceae [159 OTUs] and

Inocybaceae [132 OTUs] contain truffle-like and/or secotioid species as well as mushroom species). Overlapping areas represent the proportion of OTUs shared between sample types, whereas the non-overlapping areas represent OTUs unique to specified substrate

2002). Other mycophagy studies focusing on AM fungi need to consider the whole community in the soil and on plant roots in order to place appropriate emphasis on these dispersal events for the whole AM fungal community. Nevertheless, mammal dispersal may significantly affect the population structure of sporocarpic AM fungi such as *Glomus* spp.

Previous ECM community studies in Australia classified between 3 and 27% of taxa as truffle-like (Online Resource 1, Table S1). We report a percentage of truffle-like taxa within this range (21%). We also found that truffle-like taxa comprise dominant portions of the community. *Russula* and *Cortinarius* were the most OTU-rich taxa and were included in the relatively abundant groups from our sequencing data and in other ECM surveys in Australia (Online Resource 1, Table S1). However, as many OTUs matching these genera did not match known species, there was not enough taxonomic resolution to discern fruiting habit as these genera contain truffle-like species and mushroom species (e.g. Peintner et al. 2001; Lebel and Tonkin 2007). This limits our capacity to draw conclusions about how truffle-like fungi form part of ECM diversity, and ultimately the overall influence of mycophagous mammals on the ECM community. It also emphasises the need for further targeted truffle-like fungal surveys and taxonomic work on these

groups, coupled with continuous updating of online sequence databases.

While we took precautions in this study by removing OTUs present in negative and positive controls, contaminant DNA could still be present and errors in OTU assignment to samples can occur via tag-switching (Carlsen et al. 2012; Nguyen et al. 2015a). For this reason, we consider the proportion of overlapping taxa between sample types to be estimates. Also, taxa we observe at low relative abundances may be indistinguishable from contamination. Amplicon sequencing data are considered ‘semi-quantitative’ in that relative abundances of sequences within rather than between taxa can be more meaningful as PCR procedures may selectively amplify certain taxa more than others (among other reasons) (Amend et al. 2010). Nevertheless, Nguyen et al. (2015a) argue that relative abundances of taxa may still have ecological value, provided the sequencing errors are appropriately handled and recognised. While we cannot verify whether dominant truffle-like taxa observed were selectively amplified, comparisons within truffle-like taxa (e.g. *Hysterangium*) show that they have a higher abundance in root-associated communities. OTUs matching truffle-like taxa *Malajczukia/Mesophellia*, which have a

high relative abundance in scats, are also present in root-associating communities. As our results were consistent across two sites and seasons, we consider our assessment of the ECM community structure sufficiently accurate to be confident in the conclusion that truffle-like taxa, and their mammalian dispersers, are important to ECM communities in this system.

Unexpectedly, some OTUs matched truffle sequences of non-Australian taxa (*Tuber* sp. and *Rhizopogon pseudorozeolus*). No *Tuber* species are known to occur natively in Australia, or have associations with native flora (Bonito et al. 2013). Introduced *Tuber* species associate with introduced trees (mainly on *Quercus* and *Corylus*) in temperate regions of Australia (Linde and Selmes 2012; Thomas 2014) and are also known to associate with Pinaceae (Bonito et al. 2013). *Rhizopogon* species also associate with Pinaceae and other non-native Australian trees (Ivory and Munga 1983; Tedersoo et al. 2007) and have been recorded in Australian pine plantations (Bell and Adams 2004). Incidentally, there is a plantation of *Pinus caribaea* ca. 10 km from one of the study sites (Tinaroo Dam) (Applegate and Nicholson 1988). *Rhizopogon* was found in highest abundance in *U. caudimaculatus* and *Per. nasuta* scats at Tinaroo Dam (Nuske et al. 2018); both mammal species have been known to have home ranges within this distance (Scott et al. 1999; Streatfeild 2009). Therefore, it is possible that the OTUs matching *R. pseudorozeolus* or *Tuber* sp. resulted from native mammals consuming a non-native, introduced ectomycorrhizae associated with local *Pinus* plantations. Alternatively, they may have resulted from a contamination from laboratory processing. Nevertheless, the five OTUs identified are unlikely to alter the main findings of this study.

Conclusion

Diverse ectomycorrhizal fungal communities are vital for healthy ecosystems because of their intimate interactions with plants and pivotal role in nutrient cycling. These fungi also provide food sources for many animals, including the endangered and threatened fungal specialists; species like *B. tropica*. Little is known about how disturbances to these ecosystems can change ECM fungal communities. Our data show that mammal dispersed truffle-like taxa can form a dominant proportion of root-associating ECM fungal communities. This suggests that changes in the mammal community could, over time, induce changes in functioning ECM fungal communities, which may in turn impact plant health and nutrient cycles on a large scale. Australia has already lost many mammal species and many species are in decline (Woinarski et al. 2015). Conservation of mammal diversity may not only be imperative for ecosystem function at higher trophic levels,

but also for maintaining fungal diversity and healthy mycorrhizal-plant relationships.

Acknowledgements Sincere thanks to Queensland Parks and Wildlife Service staff Rob Miller, Lana Little, Jack Cosgrove, Karl Goetze, Chris White, Julie Bunny and Miki Bradley, World Wildlife Fund Australia staff Jessica Koleček, James Cook University students and staff Tegan Whitehead, Stephanie Todd, Naomi Bowie and Elise Chatterton and many field volunteers for help with field work and collection of samples.

Funding sources This research was supported by an Australian Postgraduate Award, Wet Tropics Management Authority (Student Grant 916), Australasian Mycological Society Research Award (2014), North Queensland Wildlife Trust (2014) and Australian Government Caring for our Country 2 Target Area Grant 2013/14 (Project ID: TAG14-00542) *Bettongia tropica* population status, viability and impact of fire with project partners James Cook University, Queensland Parks and Wildlife Service, Department of Environment and Heritage Protection and World Wildlife Fund Australia (WWF-Australia).

Authors' contributions Nuske, Susan J: designed study; collected data; sequenced DNA; analysed data; wrote manuscript.

Anslan, Sten: performed bioinformatics; edited manuscript.

Tedersoo, Leho: contributed laboratory space, primers and reagents; provided advice on high-throughput sequencing; edited manuscript.

Congdon, Bradley C: provided advice on design of study and data analysis; edited manuscript.

Abell, Sandra E: provided advice on design of study and data analysis; edited manuscript.

Data availability - Raw Illumina data is deposited in Sequence Read Archive (SRA; bioproject SRP150847).

- Sanger data is available in GenBank (Accessions KY686200-KY686202, KY697566-KY697576, KY697578-KY697619; Online Resource 2).

Compliance with ethical standards

This research was performed under permits granted by the Queensland Government, EHP (WISP15227914, WITK14639614 and WITK11258712). All wildlife was handled according to JCU Animal Ethics Guidelines (approval number: A2044). The authors declare that they have no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Abell-Davis SE (2008) Tropical hypogeous fungal sporocarp distribution in time and space. Implications for an endangered specialist mycophagous marsupial, *Bettongia tropica*. PhD thesis. School of Marine and Tropical Biology. James Cook University
- Adams F, Reddell P, Webb M, Shipton W (2006) Arbuscular mycorrhizas and ectomycorrhizas on *Eucalyptus grandis* (Myrtaceae) trees and

- seedlings in native forests of tropical North-Eastern Australia. *Aust J Bot* 54:271–281
- Amend AS, Seifert KA, Bruns TD (2010) Quantifying microbial communities with 454 pyrosequencing: does read abundance count? *Mol Ecol* 19:5555–5565. <https://doi.org/10.1111/j.1365-294X.2010.04898.x>
- Anslan S, Bahram M, Hiiesalu I, Tedersoo T (2017) PipeCraft: flexible open-source toolkit for bioinformatics analysis of custom high-throughput amplicon sequencing data. *Mol Ecol Resour* May:online first 17:e234–e240. <https://doi.org/10.1111/1755-0998.12692>
- Applegate GB, Nicholson DI (1988) Caribbean pine in an agroforestry system on the Atherton tableland in north East Australia. *Agrofor Syst* 7:3–15. <https://doi.org/10.1007/BF01890466>
- Bell TL, Adams MA (2004) Ecophysiology of ectomycorrhizal fungi associated with *Pinus* spp. in low rainfall areas of Western Australia. *Plant Ecol* 171:35–52. <https://doi.org/10.1023/B:VEGE.0000029372.78102.9d>
- Bonito G, Smith ME, Nowak M, Healy RA, Guevara G, Cázares E, Kinoshita A, Nouhra ER, Domínguez LS, Tedersoo L, Murat C, Wang Y, Moreno BA, Pfister DH, Nara K, Zambonelli A, Trappe JM, Vilgalys R (2013) Historical biogeography and diversification of truffles in the Tuberaceae and their newly identified southern hemisphere sister lineage. *PLoS One* 8:e52765. <https://doi.org/10.1371/journal.pone.0052765>
- Bougher NL, Lebel T (2001) Truffle-like (truffle-like) fungi of Australia and New Zealand. *Aust Syst Bot* 14:439–484
- Brundrett MC (2009) Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. *Plant Soil* 320:37–77
- Camacho C, Coulouris G, Avagyan V et al (2009) BLAST plus: architecture and applications. *BMC Bioinformatics* 10:421. <https://doi.org/10.1186/1471-2105-10-421>
- Carlsen T, Bjørnsgaard Aas A, Lindner D et al (2012) Don't make a mista(g)ke: is tag switching an overlooked source of error in amplicon pyrosequencing studies? *Fungal Ecol* 5:747–749. <https://doi.org/10.1016/j.funeco.2012.06.003>
- Claridge AW, May TW (1994) Mycophagy among Australian mammals. *Aust J Ecol* 19:251–275
- Gehring CA, Whitham TG (1994) Interactions between aboveground herbivores and the mycorrhizal mutualists of plants. *Trends Ecol Evol* 9:251–255
- Gehring CA, Wolf JE, Theimer TC (2002) Terrestrial vertebrates promote arbuscular mycorrhizal fungal diversity and inoculum potential in a rain forest soil. *Ecol Lett* 5:540–548. <https://doi.org/10.1046/j.1461-0248.2002.00353.x>
- Goto BT, Maia LC (2005) Sporocarpic species of arbuscular mycorrhizal fungi (Glomeromycota), with a new report from Brazil. *Acta Bot Bras* 19:633–637. <https://doi.org/10.1590/S0102-33062005000300025>
- Horton BM, Glen M, Davidson NJ, Ratkowski D, Close DC, Wardlaw TJ, Mohammed C (2013) Temperate eucalypt forest decline is linked to altered ectomycorrhizal communities mediated by soil chemistry. *For Ecol Manag* 302:329–337. <https://doi.org/10.1016/j.foreco.2013.04.006>
- Ishaq L, Barber PA, Hardy GESJ, Calver M, Dell B (2013) Seedling mycorrhizal type and soil chemistry are related to canopy condition of *Eucalyptus gomphocephala*. *Mycorrhiza* 23:359–371. <https://doi.org/10.1007/s00572-012-0476-5>
- Ivory MH, Munga FM (1983) Growth and survival of container-grown *Pinus caribaea* infected with various ectomycorrhizal fungi. *Plant Soil* 71:339–344
- Izzo AD, Meyer M, Trappe JM (2005) Hypogeous ectomycorrhizal fungal species on roots and in small mammal diet in a mixed-conifer forest. *For Sci* 51:243–254
- Janos DP, Sahley CT, Emmons LH (1995) Rodent dispersal of vesicular-arbuscular mycorrhizal fungi in Amazonian Peru. *Ecology* 76:1852–1858
- Johnson CN (1996) Interactions between mammals and ectomycorrhizal fungi. *Trends Ecol Evol* 11:503–507
- Lebel T, Tonkin JE (2007) Australasian species of Macowanites are truffle-like species of Russula (Russulaceae, Basidiomycota). *Aust Syst Bot* 20:355–381. <https://doi.org/10.1071/SB07007>
- Li W, Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* 22:1658–1659. <https://doi.org/10.1093/bioinformatics/btl158>
- Lindahl BD, Nilsson RH, Tedersoo L, Abarenkov K, Carlsen T, Kjeller R, Kõljalg U, Pennanen T, Rosendahl S, Stenlid J, Kauserud H (2013) Fungal community analysis by high-throughput sequencing of amplified markers—a user's guide. *New Phytol* 199:288–299
- Linde CC, Selmes H (2012) Genetic diversity and mating type distribution of *Tuber melanosporum* and their significance to truffle cultivation in artificially planted truffières in Australia. *Appl Environ Microbiol* 78:6534–6539. <https://doi.org/10.1128/AEM.01558-12>
- Lu X, Malajczuk N, Brundrett M, Dell B (1999) Fruiting of putative ectomycorrhizal fungi under blue gum (*Eucalyptus globulus*) plantations of different ages in Western Australia. *Mycorrhiza* 8:255–261. <https://doi.org/10.1007/s005720050242>
- Malajczuk N, Trappe JM, Molina R (1987) Interrelationships among some ectomycorrhizal trees, hypogeous fungi and small mammals: Western Australian and northwestern American parallels. *Aust J Ecol* 12:53–55
- Maser C, Nussbaum RA, Trappe JM (1978) Fungal-small mammal interrelationships with emphasis on Oregon coniferous forests. *Ecology* 59:799–809. <https://doi.org/10.2307/1938784>
- McGee PA, Baczocha N (1994) Sporocarpic Endogonales and Glomales in the scats of *Rattus* and *Perameles*. *Mycol Res* 98:246–249
- McMurdie PJ, Holmes S (2013) Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 8:e61217. <https://doi.org/10.1371/journal.pone.0061217>
- Nguyen NH, Smith D, Peay K, Kennedy P (2015a) Parsing ecological signal from noise in next generation amplicon sequencing. *New Phytol* 205:1389–1393
- Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy PG (2015b) FUNGuild: an open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecol* 20:1–8. <https://doi.org/10.1016/j.funeco.2015.06.006>
- Nuske SJ, Anslan S, Tedersoo L, Bonner MTL, Congdon BC, Abell SE (2018) The endangered northern bettong, *Bettongia tropica*, performs a unique and potentially irreplaceable dispersal function for ectomycorrhizal truffle fungi. *Mol Ecol* 27:4960–4971. <https://doi.org/10.1111/mec.14916>
- Nuske SJ, Vernes K, May TW, Claridge AW, Congdon BC, Krockenberger A, Abell SE (2017) Data on the fungal species consumed by mammal species in Australia. *Data Br* 12:251–260. <https://doi.org/10.1016/j.dib.2017.03.053>
- Peintner U, Bougher NL, Castellano MA, Moncalvo JM, Moser MM, Trappe JM, Vilgalys R (2001) Multiple origins of truffle-like fungi related to *Cortinarius* (Cortinariaceae). *Am J Bot* 88:2168–2179
- R Core Team (2012) R: A Language and Environment for Statistical Computing
- Reddell P, Gordon V, Hopkins MS (1999) Ectomycorrhizas in *Eucalyptus tetrodonta* and *E. miniata* forest communities in tropical northern Australia and their role in the rehabilitation of these forests following mining. *Aust J Bot* 47:881–907. <https://doi.org/10.1071/bt97126>
- Reddell P, Spain AV, Hopkins M (1997) Dispersal of spores of mycorrhizal fungi in scats of native mammals in tropical forests of northeastern Australia. *Biotropica* 29:184–192

- Scott LK, Hume ID, Dickman CR (1999) Ecology and population biology of long-nosed bandicoots (*Perameles nasuta*) at north head, Sydney harbour National Park. *Wildl Res* 26:805–821
- Scott PM, Shearer BL, Barber PA, Hardy GESJ (2012) Relationships between the crown health, fine root and ectomycorrhizae density of declining *Eucalyptus gomphocephala*. *Australas Plant Pathol* 42:121–131. <https://doi.org/10.1007/s13313-012-0152-4>
- Short J, Smith A (1994) Mammal decline and recovery in Australia. *J Mammal* 75:288–297
- Song Z, Schlatter D, Kennedy P, Kinkel LL, Kistler HC, Nguyen N, Bates ST (2015) Effort versus reward: preparing samples for fungal community characterization in high-throughput sequencing surveys of soils. *PLoS One* 10:e0127234. <https://doi.org/10.1371/journal.pone.0127234>
- Streatfeild C (2009) The effects of habitat fragmentation on the demography and population genetic structure of *Uromys caudimaculatus*. PhD thesis. School of Natural Resource Sciences. Queensland University of Technology
- Tedersoo L, Bahram M, Põlme S et al (2014) Global diversity and geography of soil fungi. *Science* (80-) 346:125668–1–1256688–10. <https://doi.org/10.1126/science.aaa1185>
- Tedersoo L, Suvi T, Beaver K, Kõljalg U (2007) Ectomycorrhizal fungi of the Seychelles: diversity patterns and host shifts from the native *Vateriopsis seychellarum* (Dipterocarpaceae) and *Intsia bijuga* (Caesalpiniaceae) to the introduced *Eucalyptus robusta* (Myrtaceae), but not *Pinus caribea* (Pinaceae). *New Phytol* 175: 321–333. <https://doi.org/10.1111/j.1469-8137.2007.02104.x>
- Thomas PW (2014) An analysis of the climatic parameters needed for *Tuber melanosporum* cultivation incorporating data from six continents. *Mycosphere* 5:137–142. <https://doi.org/10.5943/mycosphere/5/1/5>
- Trappe JM, Molina R, Luoma DL, et al (2009) Diversity, ecology, and conservation of truffle Fungi in forests of the Pacific northwest. United States Department of Agriculture, Forest Service, Pacific northwest Research Station. Portland, Oregon
- Van Dyck S, Gynther I, Baker A (eds) (2013) Field companion to the mammals of Australia. New Holland Publishers, Sydney
- Vernes K (2007) Are diverse mammal communities important for maintaining plant-fungal associations and ecosystem health? *Australas Plant Conserv* 15:16–18
- Vernes K, Dunn L (2009) Mammal mycophagy and fungal spore dispersal across a steep environmental gradient in eastern Australia. *Austral Ecol* 34:69–76. <https://doi.org/10.1111/j.1442-9993.2008.01883.x>
- Woinarski JCZ, Burbidge AA, Harrison PL (2015) Ongoing unraveling of a continental fauna: decline and extinction of Australian mammals since European settlement. *Proc Natl Acad Sci U S A* 112: 4531–4540. <https://doi.org/10.1073/pnas.1417301112>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.