Correction to: Biochemical applications of *Kigelia africana* (Lam.) Benth. fruit extracts in diabetes mellitus

Oladapo F. Fagbohun¹ · Olumayowa V. Oriyomi² · Mukaila B. Adekola³ · Titus A. M. Msagati⁴

Published online: 16 November 2020

© Springer-Verlag London Ltd., part of Springer Nature 2020

Correction to: Comparative Clinical Pathology

<https://doi.org/10.1007/s00580-020-03179-9>

In the article, Biochemical applications of *Kigelia africana* (Lam.) Benth. fruit extracts in diabetes mellitus, the following errata were determined:

- 1 The instrument mentioned was incorrectly described as a GC-qTOFMS throughout the manuscript. The correct descriptor should have been a GC-TOFMS.
- 2 In the **instrumentation and analytical method** section, the following text better describes the GC-TOFMS experiments.

Corrected text:

A Pegasus 4D GCxGC-TOFMS was used in 1D (GC-TOFMS) mode. The unit is composed of a 7890B Gas Chromatograph (Agilent Technologies, Santa Clara, CA), a Pegasus IV TOFMS (LECO Corporation, St Joseph, MI, USA) and an MPS/2 autosampler (Gerstel GMBH, Mulheim Am Ruhr, Germany). 1 μ L of the

sample was injected using a 10 μ L syringe into a Split/Splitless inlet held at 250 °C operating in Splitless mode. A Rxi-5SilMS column (30 m $\times$ 0.25 mm ID $\times$ 0.25 μ m DF, Restek Corporation, Bellefonte, PA, USA) was used for all separation purposes, with a column flow of 1.4 mL/min and an oven temperature program as follows: 90 °C, hold 1 min, ramped at 20 °C to 200 °C, ramped at 15 °C to 300 °C, hold for 8 min for a total runtime of 20 min. The transfer line temperature was held at 250 °C and the ion source operated at 70 eV. The acquisition mass range for the whole run was from 50 to 500 m/z with a solvent delay of 1 min. The mass spectra generated by TOF/MS were compared with those of standard mass spectra obtained from the National Institute of Standards and Technology Library (NIST) version 14.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00580-020-03179-9>

✉ Oladapo F. Fagbohun
oladapo.fagbohun@tech-u.edu.ng

¹ Department of Biomedical Engineering, First Technical University, Ibadan, Oyo State, Nigeria

² Institute of Ecology and Environmental Studies, Obafemi Awolowo University (OAU), Ile-Ife, Osun State, Nigeria

³ Department of Environmental Management and Toxicology, Federal University of Agriculture, Abeokuta, Ogun State, Nigeria

⁴ Nanotechnology and Water Sustainability Research Unit, College of Science Engineering and Technology, University of South Africa (UNISA), Florida Park, Johannesburg, South Africa