

Ring current and auroral electrojets in connection with interplanetary medium parameters during magnetic storm

Y. I. Feldstein, A. E. Levitin, S. A. Golyshev, L. A. Dremukhina, U. B. Vestchezerova, T. E. Valchuk, A. Grafe

142092 Troitsk, Moscow Region, Izmiran, Russia

(2) Georesearch Center, Telegrafenberg, D-14482 Potsdam, Germany

Received: 25 May 1993/Revised: 9 February 1994/Accepted: 11 February 1994

Abstract. The relationship between the auroral electrojet indices (AE) and the ring current magnetic field (DR) was investigated by observations obtained during the magnetic storm on 1-3 April 1973. During the storm main phase the DR development is accompanied by a shift of the auroral electrojets toward the equator. As a result, the standard AE indices calculated on the basis of data from auroral observatories was substantially lower than the real values (AE'). To determine AE' during the course of a storm main phase data from subauroral magnetic observatories should be used. It is shown that the intensity of the indices (AE') which take into account the shift of the electrojets is increased substantially relative to the standard indices during the storm main phase. AE' values are closely correlated with geoeffective solar wind parameters. A high correlation was obtained between AE' and the energy flux into the ring current during the storm main phase. Analysis of magnetic field variations during intervals with intense southward IMF components demonstrates a decrease of the saturation effect of auroral electrojet currents if subauroral stations magnetic field variations are taken into account. This applies both to case studies and statistical data. The dynamics of the electrojets in connection with the development of the ring current and of magnetospheric substorms can be described by the presence (absence) of saturation for minimum (maximum) AE index values during a 1-h interval. The ring current magnetic field asymmetry (ASY) was calculated as the difference between the maximum and minimum field values along a parallel of latitude at low latitudes. The ASY value is closely correlated with geoeffective solar wind parameters and simultaneously is a more sensitive indicator of IMF B_z variations than the symmetric ring current. ASY increases (decreases) faster during the main phase (the recovery phase) than DR . The magnetic field decay at low latitudes in the recovery phase occurs faster in the afternoon sector than at dusk.

Correspondence to: Y. Feldstein

Article not available online

Last change: October 3, 1997

helpdesk.link@springer.de

© Springer Berlin Heidelberg 1994