Correction to: Axial loading during MRI reveals insufficient effect of percutaneous interspinous implants (Aperius™ PerCLID™) on spinal canal area

Hrafnhildur Hjaltadottir¹ · Hanna Hebelka^{1,2} · Caroline Molinder¹ · Helena Brisby^{2,3} · Adad Baranto^{2,3}

Published online: 1 November 2019

© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: European Spine Journal

<https://doi.org/10.1007/s00586-019-06159-y>

Unfortunately, the ninth reference was incorrectly published in the original publication. The complete correct reference is given below.

Meyer B, Baranto A, Schils F, Collignon F, Zoega B, Tan L, LeHuec JC, NICE Trial Study Group (2018) Percutaneous interspinous spacer vs decompression in patients with neurogenic claudication: an alternative in selected patients? *Neurosurgery* 82(5):621–629. <https://doi.org/10.1093/neuros/nyx326>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00586-019-06159-y>.

✉ Adad Baranto
adad.baranto@vgregion.se

¹ Department of Radiology, Sahlgrenska University Hospital, Gothenburg, Sweden

² Sahlgrenska University Hospital, Gothenburg, Sweden

³ Department of Orthopaedics, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Sahlgrenska University Hospital, 413 45 Gothenburg, Sweden