Correction to: Deterioration of sagittal spinal alignment with age originates from the pelvis not the lumbar spine: a 4-year longitudinal cohort study

Shin Oe¹ · Yu Yamato¹ · Tomohiko Hasegawa² · Go Yoshida² · Sho Kobayashi³ · Tatsuya Yasuda² · Tomohiro Banno² · Hideyuki Arima² · Yuki Mihara² · Hiroki Ushirozako² · Tomohiro Yamada² · Koichiro Ide² · Yuh Watanabe² · Daisuke Togawa⁴ · Haruo Niwa⁵ · Yukihiro Matsuyama²

Published online: 3 June 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: European Spine Journal

<https://doi.org/10.1007/s00586-020-06431-6>

Unfortunately, Fig. 3 was incorrectly published in the original publication. The complete correct Fig. 3 is given below.

The original article has been updated.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00586-020-06431-6>.

✉ Shin Oe
mecersior@gmail.com

- ¹ Division of Geriatric Musculoskeletal Health, Department of Orthopedic Surgery, Hamamatsu University School of Medicine, 1-20-1 Handayama Higashi-ku, Hamamatsu, Shizuoka 431-3192, Japan
- ² Department of Orthopedic Surgery, Hamamatsu University School of Medicine, Hamamatsu, Japan
- ³ Department of Orthopedic Surgery, Hamamatsu Medical Center, Hamamatsu, Japan
- ⁴ Department of Orthopedic Surgery, Kindai University Nara Hospital, Ikoma, Japan
- ⁵ Department of General Medicine, Toei Hospital, Toei Town, Japan

Fig. 3 The change in spinal alignment from 2014 to 2018 in males. The PT increased significantly in the eighth decade, with no other significant change in alignment parameters noted over the 4-year period of observation