



Correction to: Ultrasound-guided erector spinae plane blocks for pain management after open lumbar laminectomy

Jesse W. Stewart¹ · Douglas Dickson² · Michael Van Hal² · Lemuelson Aryeetey² · Mary Sunna³ · Cedar Schulz³ · John C. Alexander¹ · Irina Gasanova¹ · Girish P. Joshi¹

Published online: 28 November 2023
© Springer-Verlag GmbH Germany, part of Springer Nature 2023

Correction to: European Spine Journal

<https://doi.org/10.1007/s00586-023-07881-4>

The original article has been corrected.

Table 3 data have been incorrectly published in the original publication. The complete correct Table 3 is given below.

Table 3 Numerical rating scores over time in patients undergoing open lumbar laminectomy with or without erector spinae plane block (ESPB) as a component of multimodal analgesic care

	No block (<i>n</i> =25)	ESPB (<i>n</i> =25)	<i>p</i> -value*
Preoperative, mean (SD)	4.3 (2.3) [3.3, 5.3]	4.4 (2.6) [3.3, 5.5]	0.73
PACU, mean (SD) [95% CI]	5.1 (3.7) [3.6, 6.6]	2.8 (2.2) [1.9, 3.8]	0.01
24 h, mean (SD)	5.5 (1.4) [4.9, 6.0]	3.0 (1.3) [2.4, 3.5]	<0.001
48 h, mean (SD)	4.8 (1.7) [4.0, 5.5]	3.5 (1.7) [2.9, 4.2]	0.01
72 h, mean (SD)	4.4 (1.4) [3.8, 4.9]	3.7 (2.0) [2.9, 4.6]	0.128

Data are expressed as mean (standard deviation [SD]), 95% confidence interval [CI]

*Significant if $P < 0.05$

PACU, post-anesthesia care unit

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00586-023-07881-4>.

✉ Jesse W. Stewart
jesse.stewart@utsouthwestern.edu

¹ Department of Anesthesiology and Pain Management, UT Southwestern Medical Center, 5323 Harry Hines Blvd, Dallas, TX 75390-9068, USA

² Department of Orthopedic Surgery, UT Southwestern Medical Center, Dallas, TX, USA

³ Parkland Health and Hospital System, Dallas, TX, USA