Notice of redundant publication

Massimo Porta · Guido Pozza · Renato Lauro

Published online: 27 February 2014
© Springer-Verlag Italia 2014

After publishing the paper, ‘A nomogram to estimate the proportion of patients at hemoglobin A1c target <7 % with noninsulin antidiabetic drugs in type 2 diabetes: a systematic review of 137 randomized controlled trials with 39,845 patients’ by Esposito et al. [1], it was brought to our journal’s attention that the article contained some redundancies with an earlier article published in *Diabetes, Obesity and Metabolism* entitled, ‘Proportion of patients at HbA1c target <7 % with eight classes of antidiabetic drugs in type 2 diabetes: systematic review of 218 randomized controlled trials with 78,945 patients’ by Esposito et al. [2].

The later article, by Esposito et al. [1], neither mentioned nor referenced the earlier published work.

References

1. Esposito E, Chiodini P, Bellastella G, Maiorino MI, Giugliano D (2012) A nomogram to estimate the proportion of patients at haemoglobin A1c target <7% with noninsulin antidiabetic drugs in type 2 diabetes: a systematic review of 137 randomized controlled trials with 39 845 patients. *Acta Diabetol*. Published online 28 Jan 2012. doi:[10.1007/s00592-012-0370-9](https://doi.org/10.1007/s00592-012-0370-9)
2. Esposito E, Chiodini P, Ceriello A, Giugliano D (2012) Proportion of patients at HbA1c target <7% with eight classes of antidiabetic drugs in type 2 diabetes: systematic review of 218 randomized controlled trials with 78 945 patients. *Diabetes Obes Metab* 14:228–233

This comment refers to the article available
at doi:[10.1007/s00592-012-0370-9](https://doi.org/10.1007/s00592-012-0370-9).

M. Porta (✉)
Department of Medical Sciences, University of Turin,
Turin, Italy
e-mail: massimo.porta@unito.it

G. Pozza
San Raffaele Hospital, Milan, Italy

R. Lauro
University of Rome Tor Vergata, Rome, Italy