Correction to: The ZJU index is a powerful index for identifying NAFLD in the general Chinese population

Linman Li¹ · Wenyi You² · Wei Ren³

Published online: 26 September 2019
© Springer-Verlag Italia S.r.l., part of Springer Nature 2019

Correction to: Acta Diabetol (2017) 54:905–911
<https://doi.org/10.1007/s00592-017-1024-8>

In paragraph Index calculation formulae, Point 2 should read as follows:

$VAI = [WC / (39.68 + (1.88 \times BMI))] \times (TG / 1.03) \times (1.31 / HDL)$ for males; $[WC / (36.58 + (1.89 \times BMI))] \times (TG / 0.81) \times (1.52 / HDL)$ for females [21].

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00592-017-1024-8>.

✉ Linman Li
1360896542@qq.com

- ¹ Department of Physical Examination, The First Affiliated Hospital of Chongqing Medical University, Chongqing 400016, People's Republic of China
- ² Infectious Diseases Department, The People's Hospital of Yubei, District of Chongqing City, Chongqing 401120, People's Republic of China
- ³ Department of Endocrinology, The First Affiliated Hospital of Chongqing Medical University, Chongqing 400016, People's Republic of China