Erratum to: An analysis of genotype–phenotype correlations and survival outcomes in patients with primary hyperparathyroidism caused by multiple endocrine neoplasia type 1: the experience at a single institution

Kiyomi Horiuchi · Takahiro Okamoto ·
Masatoshi Iihara · Toshihiko Tsukada

Published online: 27 June 2013
© Springer Japan 2013

Erratum to: Surg Today DOI 10.1007/s00595-012-0354-y

The original version of this article unfortunately contained the following errors:

In the “Introduction” section, in the 2nd paragraph, the 1st sentence should read:

The locus of the gene responsible for MEN1 was assigned to the long arm of chromosome 11 in 1988 [3].

In the “Methods” section, in the paragraph headed “*MEN1* gene”, the reference citation numbers should be [22] and [23], and consequently the following entries should be added to the reference list:

22. Shimizu S, Tsukada T, Futami H, Ui K, Kameya T, Kawanaka M, et al. Germline mutations of the *MEN1* gene in Japanese kindred with multiple endocrine neoplasia type 1. *Jpn J Cancer Res.* 1997;88:1029–32.

23. Kikuchi M, Ohkura N, Yamaguchi K, Takao Obara T, Tsukada T. Gene dose mapping delineated boundaries of a large germline deletion responsible for multiple endocrine neoplasia type 1. *Cancer Lett.* 2004;208:81–8.

The online version of the original article can be found under doi:[10.1007/s00595-012-0354-y](https://doi.org/10.1007/s00595-012-0354-y).

K. Horiuchi (✉) · T. Okamoto · M. Iihara
Department of Endocrine Surgery, Tokyo Women's
Medical University, 8-1 Kawada-cho, Shinjuku-ku,
Tokyo 162-8666, Japan
e-mail: horiuchi.kiyomi@twmu.ac.jp

T. Tsukada
Tumor Endocrinology Project, National Cancer Center Research
Institute, Tokyo, Japan