

## Erratum to: Surgical risk model for acute diffuse peritonitis based on a Japanese nationwide database: an initial report on the surgical and 30-day mortality

Tohru Nakagoe · Hiroaki Miyata · Mitsukazu Gotoh · Takayuki Anazawa ·  
Hideo Baba · Wataru Kimura · Naohiro Tomita · Mitsuo Shimada ·  
Yuko Kitagawa · Kenichi Sugihara · Masaki Mori

Published online: 17 January 2015  
© Springer Japan 2015

**Erratum to: Surg Today**  
**DOI 10.1007/s00595-014-1026-x**

In Tables 3 and 5, there were errors in the platelet counts.

In both tables, the following corrections should be made:

<15,000/ $\mu$ L should be <150,000/ $\mu$ L

<12,000/ $\mu$ L should be <120,000/ $\mu$ L

<8,000/ $\mu$ L should be <80,000/ $\mu$ L

---

The online version of the original article can be found under  
doi:[10.1007/s00595-014-1026-x](https://doi.org/10.1007/s00595-014-1026-x).

---

T. Nakagoe · H. Miyata · M. Gotoh · H. Baba · W. Kimura ·  
N. Tomita · M. Shimada · Y. Kitagawa  
The Japanese Society of Gastroenterological Surgery,  
Database Committee, Tokyo, Japan

H. Miyata · M. Gotoh  
National Clinical Database (NCD), Tokyo, Japan

M. Gotoh (✉) · T. Anazawa  
Department of Regenerative Surgery, Fukushima Medical  
University, 1 Hikarigaoka, Fukushima 960-1295, Japan  
e-mail: [mgotoh@fmu.ac.jp](mailto:mgotoh@fmu.ac.jp)

K. Sugihara · M. Mori  
The Japanese Society of Gastroenterological Surgery,  
Tokyo, Japan