Correction to: Standard protocol devised by the Japanese Pediatric Impedance Working Group for combined multichannel intraluminal impedance-pH measurements in children

Suguru Fukahori¹ · Hisayoshi Kawahara² · Takanori Oyama³ · Takeshi Saito⁴ · Ryuichi Shimono⁵ · Aya Tanaka⁵ · Takuo Noda³ · Reiko Hatori⁶ · Junko Fujino⁷ · Minoru Yagi¹ · Japanese Pediatric Impedance Working Group (Japanese-PIG)

Published online: 12 November 2019
© Springer Nature Singapore Pte Ltd. 2019

Correction to: Surgery Today

<https://doi.org/10.1007/s00595-019-01833-6>

The authors would like to add the following sentence in Acknowledgements in the original publication of this paper: The fee of Open choice was paid by the research grant of each affiliation of Minoru Yagi and Hisayoshi Kawahara.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00595-019-01833-6>.

✉ Suguru Fukahori
s_fukahori@med.kurume-u.ac.jp

¹ Department of Pediatric Surgery, Kurume University School of Medicine, 67 Asahi-machi, Kurume, Fukuoka 830-0011, Japan

² Department of Pediatric Surgery, Hamamatsu University School of Medicine, Shizuoka, Japan

³ Department of Pediatric Surgery, Okayama University, Okayama, Japan

⁴ Department of Pediatric Surgery, Chiba University Graduate School of Medicine, Chiba, Japan

⁵ Department of Pediatric Surgery, Kagawa University, Kagawa, Japan

⁶ Department of Pediatrics, Gunma University Graduate School of Medicine, Gunma, Japan

⁷ Department of Pediatric Surgery, Saitama Medical Center, Dokkyo Medical University, Saitama, Japan