



Correction to: Safety and efficacy of preoperative chemotherapy followed by esophagectomy versus upfront surgery for resectable esophageal squamous cell carcinoma

Takanori Kuroguchi¹ · Michitaka Honda^{1,2} · Kotaro Yamashita¹ · Masaru Hayami¹ · Akihiko Okamura¹ · Yu Imamura¹ · Shinji Mine¹ · Masayuki Watanabe¹

Published online: 31 March 2020
© Springer Nature Singapore Pte Ltd. 2020

Correction to: *Surgery Today* (2019) 49:150–157
<https://doi.org/10.1007/s00595-018-1718-8>

In the original publication, in Abstract, the second sentence of Results has been incorrectly published as:

Complications occurred in 76 (68.5%) and 76 (64.4%) patients in NAC and US groups, respectively ($p=0.51$), and severe complications occurred in 17 (22.4%) and 30 (39.5%) patients, respectively ($p=0.057$).

The correct sentence is given below:

Complications occurred in 76 (68.5%) and 76 (64.4%) patients in NAC and US groups, respectively ($p=0.51$), and

severe complications occurred in 17 (22.4%) and 30 (25.4%) patients, respectively ($p=0.057$).

Under the section: “Surgical procedures and outcomes”, the sixth sentence has been incorrectly published as:

The incidences of severe complications of Clavien–Dindo classification grade IIIa or higher were 22.4% ($n=17$) in the NAC group and 39.5% ($n=30$) in the US group ($p=0.057$).

The correct sentence is given below:

The incidences of severe complications of Clavien–Dindo classification grade IIIa or higher were 15.3% ($n=17$) in the NAC group and 25.4% ($n=30$) in the US group ($p=0.057$).

The original article can be found online at <https://doi.org/10.1007/s00595-018-1718-8>.

✉ Masayuki Watanabe
sayuki.watanabe@jfcrr.or.jp

¹ Department of Gastroenterological Surgery, The Cancer Institute Hospital of Japanese Foundation for Cancer Research, 3-8-31 Ariak, Koto-ku, Tokyo 135-8550, Japan

² Department of Minimally Invasive Surgical and Medical Oncology, Fukushima Medical University, Fukushima, Japan