



Correction to: Clinical and oncological benefits of left hepatectomy for Bismuth type I/II perihilar cholangiocarcinoma

Yoshitsugu Nakanishi¹ · Satoshi Hirano¹ · Keisuke Okamura¹ · Takahiro Tsuchikawa¹ · Toru Nakamura¹ · Takehiro Noji¹ · Toshimichi Asano¹ · Aya Matsui¹ · Kimitaka Tanaka¹ · Soichi Murakami¹ · Yuma Ebihara¹ · Yo Kurashima¹ · Yusuke Watanabe¹ · Toshiaki Shichinohe¹

Published online: 18 November 2021

© The Author(s) under exclusive licence to Springer Nature Singapore Pte Ltd. 2021

Correction to: Surgery Today
<https://doi.org/10.1007/s00595-021-02401-7>

The original publication has been corrected.

In the original publication, Reference 12 was published incorrectly. The correct version of Reference 12 is given in this Correction as below:

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

12. Nakanishi Y, Tsuchikawa T, Okamura K, Nakamura T, Tamoto E, Noji T, et al. Risk factors for a high Comprehensive Complication Index score after major hepatectomy for biliary cancer: a study of 229 patients at a single institution. *HPB (Oxford)*. 2016;18:735–41.

The original article can be found online at <https://doi.org/10.1007/s00595-021-02401-7>.

✉ Yoshitsugu Nakanishi
y.nakanishi@mac.com

¹ Department of Gastroenterological Surgery II, Faculty of Medicine, Hokkaido University, N-15, W-7, Kita-Ku, Sapporo, Hokkaido 060-8638, Japan