Correction to: Quick detection of causative bacteria in cases of acute cholangitis and cholecystitis using a multichannel gene autoanalyzer

Ryutaro Watanabe^{1,2,3} · Koji Asai¹ · Makoto Kuroda² · Manabu Kujiraoka¹ · Tsuyoshi Sekizuka² · Miwa Katagiri¹ · Nanako Kakizaki¹ · Hodaka Moriyama¹ · Manabu Watanabe¹ · Yoshihisa Saida¹

Published online: 21 February 2022
© The Author(s) under exclusive licence to Springer Nature Singapore Pte Ltd. 2022

Correction to: Surgery Today (2021) 51:1938–1945
<https://doi.org/10.1007/s00595-021-02332-3>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

In the original publication, the column headings of Table 5 are incorrectly given as: “Bile culture positive”. The column headings should be: “Bile culture positive”. The corrected Table 5 is given in this Correction.

Table 5 Correlation between the maximum colony quantity of the strain isolated in the bile culture and the patient characteristics

Factors	Maximum colony quantity (CFU/mL)			<i>p</i>
	Bile culture positive $\geq 10^6$ (<i>n</i> = 31)	Bile culture positive $< 10^6$ (<i>n</i> = 25)	Bile culture negative (<i>n</i> = 52)	
Sex (M/F)	15/16	17/8	30/22	0.356
Age	69 (31–93)	76 (37–89)	67 (30–93)	0.091
Body temperature (°C)	38.2 (36.4–39.7)	37.9 (37.1–39.6)	37.6 (36.7–38.8)	< 0.001
White blood cell (/μl)	11,500 (5600–32,700)	11,700 (6200–25,900)	12,500 (4900–26,400)	0.644
C-reactive protein (mg/dL)	14.34 (0.04–28.52)	9.49 (0.05–39.48)	7.62 (0.07–34.00)	0.352

The original article can be found online at <https://doi.org/10.1007/s00595-021-02332-3>.

✉ Koji Asai
k-asai@mvb.biglobe.ne.jp

¹ Department of Surgery, Toho University Ohashi Medical Center, 2-22-36 Ohashi, Meguro-ku, Tokyo 153-8515, Japan

² Laboratory of Bacterial Genomics, Pathogen Genomics Center, National Institute of Infectious Diseases, Tokyo, Japan

³ Department of Clinical Oncology, Toho University Graduate School, Tokyo, Japan