



María Gómez-Rocha · Jai More · Kamil Serafin

Baryon Masses Estimate in Heavy Flavor QCD

An Effective Particle Approach to Hadron Spectra

Received: 16 April 2023 / Accepted: 9 May 2023 / Published online: 10 June 2023
© The Author(s) 2023

Abstract We apply the renormalization group procedure for effective particles (RGPEP) to the QCD eigenvalue problem for only heavy quarks. We derive the effective Hamiltonian that acts on the Fock space by solving the RGPEP equation up to second order in powers of the coupling constant. The eigenstates that contain three quarks and two or more gluons are eliminated by inserting a gluon-mass term in the component with one gluon and the eigenvalue problem for baryons is formulated. We estimate masses for bbb and ccc states and find that the results match the estimates obtained in lattice QCD and in quark models.

1 Introduction

In spite of many years of research, the issue of bound states in QCD remains to be a long-standing problem to which an exact solution is still unknown. The QCD Hamiltonian which defines the Schrödinger equation is full of complexities. The determination of its eigenvalues, which would lead to hadron masses and the corresponding wave functions is certainly not straightforward.

The main difficulty concerns the fact that in quantum field theory one needs to deal with an infinite number of degrees of freedom in the bound-state equation $\hat{H}|\psi\rangle = E|\psi\rangle$. For a baryon, which is the case discussed in this work, the eigenstate has the following structure in terms of Fock components

$$|\psi\rangle = |3Q\rangle + |3QG\rangle + |3QGG\rangle + \dots, \quad (1)$$

where we have denoted $|3Q\rangle \equiv |QQQ\rangle$; and there is no limit in the number of particles allowed.

In this context, the renormalization group procedure for effective particles (RGPEP) was formulated as a non-perturbative tool to construct bound states in quantum field theory. Nonetheless, any candidate for a basic physical theory requires an initial perturbative search for the set of interaction terms that provides the basis on which the full effective theory can be constructed in a series of successive approximations [1].

The RGPEP has its origin in the similarity renormalization group (SRG) for Hamiltonians [2,3] but, in addition, it introduces the concept of effective particles [1,4]. The renormalization-group approach allows one to consider particle interactions and phenomena at different energy scales. The key idea is that it is possible to express the initial Hamiltonian through a unitary similarity transformation in a scale-dependent operator basis, in such a way that for a certain scale, the number of non-negligible Fock components is small. The eigenstates depend on the renormalization-group parameter t too:

$$|\psi_t\rangle = |3Q_t\rangle + |3Q_tG_t\rangle + |3Q_tG_tG_t\rangle + \dots. \quad (2)$$

Presented by María Gómez-Rocha at the International Conference on the Structure of Baryons, November 7–11th, Universidad Pablo de Olavide, Sevilla, Spain

Universidad de Granada, Departamento de Física Atómica, Molecular y Nuclear and Instituto Carlos I de Física Teórica y Computacional, Granada, Spain · Department of Physics, Indian Institute of Technology Bombay, Powai, Mumbai 400076, India · Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China

If an infinite number of components can be neglected in Eq. (2), the bound-state equation is enormously simplified and one can attempt to seek numerical solutions to the equation.

The RGPEP has been applied and solved exactly in several simple theories [5–7]. But, for the complex case of QCD, only perturbative expressions of the Hamiltonian have been considered so far. Second-order calculations with the inclusion of a gluon-mass ansatz have allowed us to examine the effective potential between heavy quarks in mesons [8] and in baryons [9]. Third-order calculations have been employed to calculate the running coupling in the front-form Hamiltonian [4], and recently, a new regularization procedure that includes a canonical gluon mass has led to analogous results [10].

In this contribution, we focus on the study of triply heavy baryon spectra with equal quark masses, ccc and bbb . For interested readers, we refer to a detailed analysis of ccb and bbc states provided in [9].

The next section presents the most general step in the RGPEP approach. In Sect. 3 we derive the bound state equation for a system of three heavy quarks and provide the analytical result in Sect. 4. The numerical setup is presented in Sect. 5 and the corresponding results are commented in Sect. 6. Finally, Sect. 7 concludes the article.

2 Key Elements of the RGPEP

The starting point of this method is the Lagrangian density of the chosen theory. In this particular case, we choose QCD, \mathcal{L}_{QCD} . The classical Hamiltonian, H_{QCD} , can be derived using Noether's theorem to calculate the energy-momentum tensor.

We use the front-form of dynamics [11, 12]. In this form, four vectors are represented as $x^\mu = (x^+, x^-, x^\perp)$, where $x^+ = x^0 + x^3$, $x^- = x^0 - x^3$, $x^\perp = (x^1, x^2)$, and the scalar product in Minkowski space-time is given by $a \cdot b = \frac{1}{2}a^+b^- + \frac{1}{2}a^-b^+ - a^\perp b^\perp$.

The quantum Hamiltonian $\hat{H}_{\text{QCD}}^{\text{can}}$ is derived using canonical quantization with the initial conditions set on the hypersurface $x^+ = 0$ and in the light-cone gauge $A^+ = 0$. The Hamiltonian can be expressed by the “-” component of the four-momentum operator, \hat{P}^μ , which is the generator of space-time translations [12]:

$$\hat{H}_{\text{QCD}}^{\text{can}} = \hat{P}^- = \int dx^- d^2x^\perp : \hat{\mathcal{H}}_{x^+=0} : . \quad (3)$$

The dots on both sides of the Hamiltonian density, $\hat{\mathcal{H}}_{x^+=0}$, indicate normal ordering of creation and annihilation operators. In the sequel we use the notation considered in [8, 12, 13].

The canonical Hamiltonian needs regularization and counterterms. The regularized canonical Hamiltonian with counterterms is called *initial* Hamiltonian, since it provides the initial condition for solving the RGPEP equation.

The regularization is provided by inserting functions defined in Ref. [8] in every interaction vertex. Such functions depend on ultraviolet and small- x cutoffs, Δ and δ , respectively, which will be removed at the end of the calculation.

The RGPEP provides a means for the calculation of counterterms. It introduces effective particle operators related by a unitary transformation

$$q_s = \mathcal{U}_s q_0 \mathcal{M}_s^\dagger, \quad (4)$$

where s has units of length and plays the role of a renormalization group parameter. It is associated with the *size* of the effective particles. If q and q^\dagger are operators that annihilate or create pointlike particles in the Fock space, effective particle operators q_s and q_s^\dagger annihilate or create particles of size s . It is convenient to consider the scale parameter $\lambda = 1/s$ which has units of energy and the parameter $t = s^4$ which we have already used in Eq. (2).

The renormalization-group parameter labels a family of equivalent Hamiltonians that correspond to the same theory but expressed in terms of degrees of freedom that are differently defined. If $H_0 = \mathcal{H}_0(q_0)$ is the initial Hamiltonian, then the RGPEP demands that:

$$\mathcal{H}_0(q_0) = \mathcal{H}_t(q_t). \quad (5)$$

An effective Hamiltonian that satisfies this condition is a solution of the RGPEP equation:

$$\mathcal{H}'_t = [[\mathcal{H}_f, \mathcal{H}_{P_t}], \mathcal{H}_t], \quad (6)$$

where $\mathcal{H}_t = \mathcal{H}(q_0)$, and the prime on the effective Hamiltonian, \mathcal{H}_t , indicates differentiation with respect to t . The subscript f stands for *free* and refers to terms that do not depend on the coupling constant. Finally, \mathcal{H}_{P_t} is identical to \mathcal{H}_t but multiplied by a factor $\frac{1}{2} (\sum_i p_i^+)^2$, with i referring to all incoming and outgoing particles involved in a vertex [1].

The ease of solving Eq. (6) depends on the complexity of the initial Hamiltonian. Although there are theories for which it is possible to find exact solutions [5–7], the complexity of QCD forces us to use a perturbative expansion in powers of the coupling constant. Such form of a solution can be written as

$$\mathcal{H}_t = \mathcal{H}_0 + g\mathcal{H}_{t1} + g^2\mathcal{H}_{t2} + g^3\mathcal{H}_{t3} + g^4\mathcal{H}_{t4} + \dots \quad (7)$$

The numerical subscript 0, 1, 2, ... refers to the power of the coupling constant. Thus, $\mathcal{H}_0 \equiv \mathcal{H}_f$ is the 0th-order term which does not depend on the coupling constant g or on the renormalization-group parameter t . In the 2nd-order expansion one has

$$\mathcal{H}'_0 + g\mathcal{H}'_{t1} + g^2\mathcal{H}'_{t2} = [[\mathcal{H}_0, \mathcal{H}_0 + g\mathcal{H}_{P_t1} + g^2\mathcal{H}_{P_t2}], \mathcal{H}_0 + g\mathcal{H}_{t1} + g^2\mathcal{H}_{t2}], \quad (8)$$

which can be solved order by order:

$$\mathcal{H}'_0 = 0, \quad (9)$$

$$g\mathcal{H}'_{t1} = [[\mathcal{H}_0, g\mathcal{H}_{P_t1}], \mathcal{H}_0], \quad (10)$$

$$g^2\mathcal{H}'_{t2} = [[\mathcal{H}_0, g^2\mathcal{H}_{P_t2}], \mathcal{H}_0] + [[\mathcal{H}_0, g\mathcal{H}_{P_t1}], g\mathcal{H}_{t1}]. \quad (11)$$

Solving Eqs. (9)–(11) yields exponentials of products of t by differences of invariant masses squared. These functions play the role of form factors that appear at interaction vertices. The renormalized Hamiltonian is determined by the initial condition that at $t = 0$ it should equal the regularized canonical Hamiltonian plus counterterms. The counterterms should be such that every matrix element of the renormalized Hamiltonian is cutoff independent for $t > 0$, i.e. free of ultraviolet divergences. In this work, we restrict our calculation to second-order expansions.

Note that this perturbative expansion is made at the level of the RGPEP equation, not at the level of the Schrödinger equation.

3 Effective Hamiltonian and Bound-State Equation for Triply Heavy Baryons

The simplest possible systems that can be considered in QCD are heavy quarkonia and triply-heavy baryons. Thus, we simplify the picture by considering only heavy flavors, and neglecting light quarks. The eigenvalue problem simplifies enormously choosing the renormalization-group parameter in the following region

$$m_Q \gg \lambda \gg \Lambda_{\text{QCD}}, \quad (12)$$

where m_Q is the quark mass. The fact that λ is much larger than Λ_{QCD} allows one to keep only the first term in the Hamiltonian expansion in powers of the coupling constant [4]. The condition $m_Q \gg \lambda$, on the other hand, makes Fock sectors with extra quark-antiquark pairs strongly suppressed by RGPEP form factors and they can be neglected. However, sectors with more gluons cannot be neglected, since they are massless, and many of them can be produced without adding much to the invariant mass of a system. We cannot deal with infinitely many Fock sectors of gluons. To address this problem, we drop all the sectors with more than one gluon and account for them by introducing a gluon mass ansatz in the sector $Q_t \bar{Q}_t G_t$ for mesons and $Q_t Q_t Q_t G_t$ for baryons [8, 9]. Thus, our gluon mass ansatz accounts for all possible non-Abelian effects that we cannot take into account explicitly. Higher-order calculations should be able to replace such an ansatz by elements of the theory.

The triply-heavy baryon bound-state problem with two Fock sectors (i.e. in the second order in the RGPEP) and gluon mass ansatz is (cf. [8] and [9] for more details):

$$\left\{ \begin{bmatrix} (H_{t0} + \mu_t^2) & gH_{t1} \\ gH_{t1} & (H_{t0} + g^2H_{t2}) \end{bmatrix} - E \right\} \begin{bmatrix} |3Q_t G_t\rangle \\ |3Q_t\rangle \end{bmatrix} = 0, \quad (13)$$

where μ_t^2 is the gluon-mass operator, which acts on the $Q_t Q_t Q_t G_t$ sector. We assume that the mass ansatz depends on the relative motion of the gluon with respect to the quarks in that sector.

Since we consider only terms up to second order in powers of the coupling constant in the effective Hamiltonian, the approximate eigenvalue problem Eq. (13) can be reduced to the sector with no gluons [14]. Matrix elements after the reduction are

$$\begin{aligned} & \langle l | H_{\text{eff } t} | r \rangle \\ &= \langle l | \left[H_{t0} + g^2 H_{t2} + \frac{1}{2} g H_{t1} \left(\frac{1}{E_l - H_{t0} - \mu_t^2} + \frac{1}{E_r - H_{t0} - \mu_t^2} \right) g H_{t1} \right] | r \rangle. \end{aligned} \quad (14)$$

where left (l) and right (r) states are both in the $3Q_t$ sector, $H_f |l\rangle = E_l |l\rangle$ and $H_f |r\rangle = E_r |r\rangle$, where H_f is the *free* term of the Hamiltonian, that does not depend on the coupling constant.

4 Result: Coulomb and Harmonic-Oscillator Potentials

The effective front-form eigenvalue equation for baryons has the following structure

$$H_{\text{eff } t} |3Q_t\rangle = \frac{M^2 + P^{\perp 2}}{P^+} |3Q_t\rangle, \quad (15)$$

where the state $|3Q_t\rangle$ is defined as

$$|3Q_t\rangle = \int_{123} P^+ \tilde{\delta}_{P,123} \psi_t(123) \frac{\epsilon^{c_1 c_2 c_3}}{\sqrt{6}} b_{t1}^\dagger b_{t2}^\dagger b_{t3}^\dagger |0\rangle. \quad (16)$$

where the spin-momentum wave function, $\psi_t(123)$ is multiplied by the color factor $\epsilon^{c_1 c_2 c_3} / \sqrt{6}$. We have used the shortcut notation $\tilde{\delta}_{P,123} = 2(2\pi)^3 \delta^3(P - p_1 + p_2 - p_3)$ for the delta function of momentum conservation. Details of the structure of the effective Hamiltonian can be found in [9].

In the non-relativistic limit, the expressions of the interaction potentials and mass functions simplify enormously. To define this limit we introduce variables defined in [15,16],

$$K_{12}^\perp = \sqrt{\frac{x_1 + x_2}{6x_1 x_2}} \kappa_{12}^\perp, \quad K_{12}^z = \sqrt{\frac{x_1 + x_2}{6x_1 x_2}} \frac{x_1 - x_2}{x_1 + x_2} m_Q, \quad (17)$$

$$Q_{12}^\perp = \sqrt{\frac{2/9}{x_3(1-x_3)}} \kappa_3^\perp, \quad Q_{12}^z = \sqrt{\frac{2/9}{x_3(1-x_3)}} (3x_3 - 1) m_Q. \quad (18)$$

where κ_{12}^\perp is the relative transverse momentum of particle 1 with respect to particle 2, κ_3^\perp is the relative transverse momentum of particle 3 with respect to particles 1 and 2, and x_1, x_2, x_3 are longitudinal momentum fractions $x_i = p_i^+ / P^+$ of particles $i = 1, 2$, and 3, respectively. In the nonrelativistic approximation the eigenvalue equation can be written in the form

$$\begin{aligned} & \left[\frac{\mathbf{K}_{12}^2}{2\mu_{12}} + \frac{\mathbf{Q}_3^2}{2\mu_{3(12)}} - B + 3 \frac{\delta m_{1t}^2}{2m_Q} \right] \psi_t(123) \\ &+ \sum_{\sigma_1' \sigma_2'} \int \frac{d^3 K'_{12}}{(2\pi)^3} [f_{i12,1'2'} V_{C,BF}^{12} + W^{12}] \psi_t(1'2'3) \\ &+ \sum_{\sigma_2' \sigma_3'} \int \frac{d^3 K'_{23}}{(2\pi)^3} [f_{i23,2'3'} V_{C,BF}^{23} + W^{23}] \psi_t(12'3') \\ &+ \sum_{\sigma_3' \sigma_1'} \int \frac{d^3 K'_{31}}{(2\pi)^3} [f_{i31,3'1'} V_{C,BF}^{31} + W^{31}] \psi_t(1'23') = 0, \end{aligned} \quad (19)$$

where B is the binding energy; $V_{C,BF}^{ij} = V_{C,BF}(\mathbf{K}_{ij}, \mathbf{K}'_{ij})$ and $W^{ij} = W(\mathbf{K}_{ij} - \mathbf{K}'_{ij})$ are, respectively, the Coulomb term with Breit-Fermi (BF) corrections and the additional interaction resulting from the gluon mass

ansatz. $\mu_{12} = m_Q/2$, $\mu_{3(12)} = 2m_Q/3$ are the reduced masses. Both V and W are similar to the ones in the quarkonium case [8]:

$$V_{C,BF}(\mathbf{K}, \mathbf{K}') = -\frac{2}{3}g^2 \frac{1}{\Delta\mathbf{K}^2}(1 + BF), \quad (20)$$

$$W(\Delta\mathbf{K}) = -\frac{2}{3}g^2 \left[\frac{1}{(\Delta K^z)^2} - \frac{1}{\Delta\mathbf{K}^2} \right] \frac{\mu^2}{\mu^2 + \Delta\mathbf{K}^2} \exp \left[-2tm_Q^2 \frac{\Delta\mathbf{K}^4}{(\Delta K^z)^2} \right], \quad (21)$$

where $\Delta\mathbf{K} = \mathbf{K} - \mathbf{K}'$ and the RGPEP form factor is

$$f_{ij,i'j'} = \exp \left\{ -16t[\mathbf{K}_{ij}^2 - (\mathbf{K}'_{ij})^2]^2 \right\}. \quad (22)$$

We assume now that the mass ansatz μ^2 dominates $\Delta\mathbf{K}^2$ in the relevant integration range, then $\frac{\mu^2}{\mu^2 + \Delta\mathbf{K}^2} \approx 1$, and the wave function can be expanded in a Taylor series in such a way that the resulting potential that corrects the Coulomb term is a harmonic oscillator one, with oscillator frequencies [9]

$$\omega_{\text{baryon}} = \frac{\sqrt{3}}{2} \sqrt{\frac{\alpha}{18\sqrt{2\pi}}} \frac{\lambda^3}{m_Q^2}. \quad (23)$$

The result of quarkonia and triply heavy baryons differ by a factor $\sqrt{3}/2$, in such a way that $\omega_{\text{baryon}}^2/\omega_{\text{meson}}^2 = 3/4$.

5 Numerical Studies

To provide numerical results for heavy quarkonia and baryons, we need to estimate the values of α , m_c , and m_b . Although the RGPEP is defined to provide an exact effective Hamiltonian independent of the scale parameter λ , the second-order approximation results in a certain λ -dependence, which is negligible in a window of values of λ (cf. Figures in Refs. [17, 18]).

We assume that the scale parameter is proportional to the coupling constant and the quark mass, $\lambda \sim \sqrt{\alpha} m_Q$. In this way, if α is sufficiently small, the assumption satisfies our hierarchy of scales Eq. (12) and, for the quark-antiquark system $\lambda \gg k_B \sim \alpha\mu$, where k_B is the Bohr momentum and μ is the quark reduced mass. This ensures that the RGPEP form factors are approximately 1 and do not influence significantly the eigenvalue problem¹ [9]. Furthermore, the fact that λ is proportional to $\sqrt{\alpha}$ makes the resulting hadron binding energies proportional to α^2 , in analogy to QED [19]. In fact, the harmonic oscillator frequencies obtained from QCD are expected to be comparable in size with the strong-interaction Rydberg-like constant $R = \mu(4\alpha/3)^2/2$, since the low-mass quarkonium spectra can be characterized as intermediate between the Coulomb and the oscillator spectra [20].

Several simplifications are taken into account in this pilot application of our method. The numerical sketch we provide yields approximate results restricted to the low-mass hadron spectrum. In this numerical sketch, we estimate the Coulomb effects in first-order perturbation theory around the oscillator solution. Therefore, we consider only the diagonal matrix elements of the Coulomb potential in the basis of harmonic oscillator wave functions, since the effects of the non-diagonal ones are relatively small and do not change significantly the lowest-mass heavy-baryon spectrum. In particular, those effects are smaller than the effects due to spin-dependent interactions, which we neglect (yet we include the effects of the Pauli exclusion principle).

5.1 Adjustment of Parameters

The coupling constant α depends on the scale parameter λ in the following way [4]:

$$\alpha = \left[\beta_0 \log \left(\frac{\lambda^2}{\Lambda_{\text{QCD}}} \right) \right]^{-1} \quad (24)$$

Table 1 Masses in MeV for all bbb and ccc states up to the second excited band of the harmonic oscillator

| bbb -states | | | ccc -states | | |
|------------------|-------------|-------|------------------|-------------|------|
| J^P | Name | Mass | J^P | Name | Mass |
| 1/2 ⁺ | $B_{1/2^+}$ | 14885 | 1/2 ⁺ | $B_{1/2^+}$ | 5350 |
| | $C_{1/2^+}$ | 14896 | | $C_{1/2^+}$ | 5358 |
| 3/2 ⁺ | 0ω | 14347 | 3/2 ⁺ | 0ω | 4797 |
| | $A_{3/2^+}$ | 14832 | | $A_{3/2^+}$ | 5309 |
| | $C_{3/2^+}$ | 14896 | | $C_{3/2^+}$ | 5358 |
| | $D_{3/2^+}$ | 14917 | | $D_{3/2^+}$ | 5374 |
| 5/2 ⁺ | $C_{5/2^+}$ | 14896 | 5/2 ⁺ | $C_{5/2^+}$ | 5358 |
| | $D_{5/2^+}$ | 14917 | | $D_{5/2^+}$ | 5374 |
| 7/2 ⁺ | $C_{7/2}$ | 14896 | 7/2 ⁺ | $C_{7/2}$ | 5358 |
| 1/2 ⁻ | 1ω | 14645 | 1/2 ⁻ | 1ω | 5103 |
| 3/2 ⁻ | 1ω | 14645 | 3/2 ⁻ | 1ω | 5103 |

with $\beta_0 = (33 - 2n_f)/(12\pi)$. We take $n_f = 2$, for two heavy flavors, b and c , though the result does not change significantly for n_f in the range up to 5. The value of $\Lambda_{\text{QCD}} = 371$ MeV is imposed by the fact that $\alpha = 0.1181$ for $\lambda = M_Z = 91.1876$ GeV.

Quark masses, m_b and m_c , and the renormalization group parameter λ are determined by the fit of computed heavy quarkonia spectra to the known experimental ones. The numerical results of this fitting are given in Appendix A. Hence, our estimates of the baryon masses are predictions without any free parameters.

We would like to point out that our estimates are in a primitive stage of development. However, the obtained results in this crude approximation are in surprisingly good agreement with other long-standing and widely used approaches. The purpose of this preliminary study is to find out whether the oscillator terms that follow from the assumption of gluon mass are capable of reproducing a good approximation to the heavy hadron spectrum. This would motivate higher-order studies in our perturbative expansion Eq. (7), to provide a theoretical explanation of the gluon-mass generation.

Therefore, we ignore Breit-Fermi spin interactions and estimate Coulomb effects by evaluating the expectation values of the corresponding interaction terms in the oscillator eigenstates. In this paper we select the most remarkable results and present them in Table 1 and Figure 1.

6 Analysis of Results

The notation is the following. State 0ω is the ground state of the system, while state 1ω is the first (orbitally) excited state. States called A , B , C , and D in bbb and ccc refer to the second excitation of the harmonic oscillator with excitation energy 2ω (with $\omega \equiv \omega_{\text{baryon}}$) above the ground state. They are mixtures of radial and orbital excitations and their masses differ due to different expectation values of the Coulomb potential.

The values of masses obtained for bbb and ccc baryons agree well with model calculations [21–27] including quark-diquark [28] and hypercentral approximations [29,30], Regge phenomenology [31,32], bag models [33–35], pNRQCD [36], sum rules [37–40], Dyson-Schwinger approach [41] and lattice studies [42–48].

We remark the comparison of the ground state of ccc , for which different lattice approaches yield values between 4733 and 4796 MeV. Our result, 4797 MeV, differs by 29 MeV from the average result, 4768 MeV, which corresponds to 0.6%. In the case of the ground state of bbb , we obtain 14347 MeV, as compared with the lattice result of 14369 MeV, a difference of 23 MeV, 0.2%. Concerning mass splittings, we differ in about 10% with lattice results provided in [43] for bbb states, and in 20% with lattice results [45] for the case of ccc states. Analyses of results for states bbc and ccb are not presented in this document. The reader is invited to consult the detailed analysis provided in [9].

It is surprising that this preliminary approximation of our RGPEP method with no free parameters, after fitting quark masses and scale to heavy quarkonia spectra, produces similar splittings to those obtained from advanced calculations.

¹ Note that form factors are necessary in higher-order calculations since they regulate terms that otherwise would be divergent.




Fig. 1 Graphical representation of mass spectra for all bbb and ccc states up to second band of harmonic oscillators

7 Conclusion

The effective Hamiltonian for heavy quarkonia and triply-heavy baryons derived in the second-order of our RGPEP with gluon-mass ansatz leads to baryon mass spectra that are comparable with the expectation obtained from other approaches to physics of bbb and ccc .

The considered method is invariant under boosts and in principle appears capable of providing a relativistic description of hadrons in terms of a small number of effective constituents, with suitably adjusted size. Therefore, an extension to higher-order calculations appears worth doing. A fourth-order calculation is needed to verify if the introduced gluon-mass ansatz provides an adequate representation of the gluon dynamics in the presence of heavy quarks. Furthermore, such calculations are also needed in the study of spin splittings and rotational symmetry. As a remark, it should be pointed out that the ratio $\sqrt{8/6}$ of harmonic oscillator frequencies in heavy quarkonia and baryons is close to the ratio $\sqrt{8/5}$ obtained for u and d quarks in constituent models using the gluon condensate. This suggests studying if the RGPEP formalism can be applied also to light hadrons as built from constituent quarks and massive gluons. Even in the heavy-quarks case, the effective oscillator potential provides simple wave functions that can be used in relativistic processes involving heavy hadrons.

Acknowledgements We thank Stanisław D. Głazek for fruitful discussions and acknowledge financial support by the FEDER2020 funds, project ref. A-FQM-406-UGR20 and from MCIN/AEI/10.13039/501100011033, Project Ref. PID2020-114767GB-I00. Figures 1 and 2 are made using the open software JaxoDraw [49] distributed under the GNU General Public license. J. M. would like to thank the Department of Science and Technology (DST), Government of India, for financial support through Grant No. SR/WOS-A/PM-6/2019(G).

Author contribution M.G.R. and K.S. calculated the expressions for the mesons and baryon potentials analytically. K.S. did the calculation of masses and wave functions and provided numerical result. M. Gomez-Rocha prepared the figures, wrote the paper, and presented the contribution in the conference. J. M. contributed to discussions and checked some calculations. All authors reviewed the manuscript.

Funding Funding for open access publishing: Universidad de Granada/CBUA

Declarations

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Masses of Quarks and Other Parameters

Quark masses and scale parameters are calculated using a fit to the masses of $\Upsilon(1S)$, $\Upsilon(2S)$, and $\chi_{b1}(1P)$ for bottomonia and J/ψ , $\psi(2S)$ and $\chi_{c1}(1P)$ for charmonia. Results of the fit yield:

$$m_b = 4698 \text{ MeV} \quad \text{and} \quad \lambda_{b\bar{b}} = 4258 \text{ MeV}, \quad (25)$$

$$m_c = 1460 \text{ MeV} \quad \text{and} \quad \lambda_{c\bar{c}} = 1944 \text{ MeV}, \quad (26)$$

These values are associated, respectively, with

$$\alpha(\lambda_{b\bar{b}}) = 0.2664 \quad \text{and} \quad \omega_{b\bar{b}} = 268.8 \text{ MeV}, \quad (27)$$

$$\alpha(\lambda_{c\bar{c}}) = 0.3926 \quad \text{and} \quad \omega_{c\bar{c}} = 321.6 \text{ MeV}. \quad (28)$$

References

1. S.D. Glazek, Acta Phys. Polon. **B43**, 1843–1862 (2012)
2. Stanislaw D. Glazek, Kenneth G. Wilson, Phys. Rev. D **48**, 5863–5872 (1993)
3. S.D. Glazek, K.G. Wilson, Phys. Rev. D **49**, 4214–4218 (1994)
4. María Gómez-Rocha, Stanislaw D. Glazek, Phys. Rev. D **92**(6), 065005 (2015)
5. S.D. Glazek, Phys. Rev. D **85**, 125018 (2012)
6. S.D. Glazek, Phys. Rev. D **87**(12), 125032 (2013)
7. Stanislaw D. Glazek, Elementary example of exact effective-Hamiltonian computation. Phys. Rev. D **103**(1), 014021 (2021)
8. Stanislaw D. Glazek, María Gómez-Rocha, Jai More, Kamil Serafin, Renormalized quark–antiquark Hamiltonian induced by a gluon mass ansatz in heavy-flavor QCD. Phys. Lett. B **773**, 172–178 (2017)
9. Kamil Serafin, María Gómez-Rocha, Jai More, Stanislaw D. Glazek, Approximate Hamiltonian for baryons in heavy-flavor QCD. Eur. Phys. J. C **78**(11), 964 (2018)
10. Juan José Gálvez-Viruet and María Gómez-Rocha, Asymptotic freedom using a gluon mass as a regulator. EPJ Web Conf. **274**, 02006 (2022)
11. A.M. Paul, Dirac. Rev. Mod. Phys. **21**, 392–399 (1949)
12. Stanley J. Brodsky, Hans-Christian. Pauli, Stephen S. Pinsky, Phys. Rept. **301**, 299–486 (1998)
13. S.D. Glazek, Phys. Rev. D **69**, 065002 (2004)
14. G. Kenneth, Phys. Rev. D **2**, 1438 (1970)
15. D. Stanislaw, Acta Phys. Polon. **B42**, 1933–2010 (2011)
16. Arkadiusz P. Trawiński, Stanislaw D. Glazek, Stanley J. Brodsky, Guy F. de Téramond, Hans Günter, Dosch, Phys. Rev. D **90**(7), 074017 (2014)
17. Stanislaw D. Glazek, Kenneth G. Wilson, Asymptotic freedom and bound states in Hamiltonian dynamics. Phys. Rev. D **57**, 3558–3566 (1998)
18. S.D. Glazek, J. Mlynik, Phys. Rev. D **67**, 045001 (2003)
19. Kenneth G. Wilson, Timothy S. Walhout, Avaroth Harindranath, Wei-Min. Zhang, Robert J. Perry, Stanislaw D. Glazek, Phys. Rev. D **49**, 6720–6766 (1994)
20. E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane, Tung-Mow. Yan, Charmonium: The Model. Phys. Rev. D **17**, 3090 (1978) [Erratum: Phys.Rev.D 21, 313 (1980)]
21. J.D. Bjorken, Is the ccc a new deal for baryon spectroscopy? AIP Conf. Proc. **132**, 390–403 (1985)
22. Masayasu Tsuge, Toshiyuki Morii, Junya Morishita, Baryon Spectrum in the Potential Model With Relativistic Corrections. Mod. Phys. Lett. A **1**, 131 (1986) [Erratum: Mod. Phys. Lett.A2,283(1987)]
23. B. Silvestre-Brac, Spectrum and static properties of heavy baryons. Few Body Syst. **20**, 1–25 (1996)
24. Yu. Jia, Variational study of weakly coupled triply heavy baryons. JHEP **10**, 073 (2006)
25. A.P. Martynenko, Ground-state triply and doubly heavy baryons in a relativistic three-quark model. Phys. Lett. B **663**, 317–321 (2008)
26. W. Roberts, Muslema Pervin, heavy baryons in a quark model. Int. J. Mod. Phys. A **23**, 2817–2860 (2008)
27. J. Vijande, A. Valcarce, H. Garcilazo, Constituent-quark model description of triply heavy baryon nonperturbative lattice QCD data. Phys. Rev. D **91**(5), 054011 (2015)

28. Floriana Giannuzzi, Doubly heavy baryons in a Salpeter model with AdS/QCD inspired potential. *Phys. Rev. D* **79**, 094002 (2009)
29. Zahra Ghalehovi, Ali Akbar Rajabi, Si.-xue Qin, H Dirk Rischke, Ground-state masses and magnetic moments of heavy baryons. *Mod. Phys. Lett. A* **29**, 1450106 (2014)
30. Zalak Shah and Ajay Kumar Rai, Masses and Regge trajectories of triply heavy Ω_{ccc} and Ω_{bbb} baryons. *Eur. Phys. J. A* **53**(10), 195 (2017)
31. Ke-Wei. Wei, Bing Chen, Xin-Heng. Guo, Masses of doubly and triply charmed baryons. *Phys. Rev. D* **92**(7), 076008 (2015)
32. Ke-Wei. Wei, Bing Chen, Na. Liu, Qian-Qian. Wang, Xin-Heng. Guo, Spectroscopy of singly, doubly, and triply bottom baryons. *Phys. Rev. D* **95**(11), 116005 (2017)
33. W. Ponce, Heavy quarks in a spherical bag. *Phys. Rev. D* **19**, 2197 (1979)
34. P. Hasenfratz, R.R. Horgan, J. Kuti, J.M. Richard, Heavy baryon spectroscopy in the QCD bag model. *Phys. Lett.* **94B**, 401–404 (1980)
35. Andrius Bernotas, Vytautas Simonis, Heavy hadron spectroscopy and the bag model. *Lith. J. Phys.* **49**, 19–28 (2009)
36. Felipe J. Llanes-Estrada, Olga I. Pavlova, Richard Williams, A first estimate of triply heavy baryon masses from the pNRQCD perturbative static potential. *Eur. Phys. J. C* **72**, 2019 (2012)
37. Jian-Rong. Zhang, Ming-Qiu. Huang, Deciphering triply heavy baryons in terms of QCD sum rules. *Phys. Lett. B* **674**, 28–35 (2009)
38. Zhi-Gang. Wang, Analysis of the triply heavy baryon states with QCD sum rules. *Commun. Theor. Phys.* **58**, 723–731 (2012)
39. T.M. Aliev, K. Azizi, M. Savci, Masses and residues of the triply heavy spin-1/2 Baryons. *JHEP* **04**, 042 (2013)
40. T.M. Aliev, K. Azizi, M. Savci, Properties of triply heavy spin-3/2 baryons. *J. Phys.* **G41**, 065003 (2014)
41. Si-Xue. Qin, Craig D. Roberts, Sebastian M. Schmidt, Poincaré-covariant analysis of heavy-quark baryons. (2018)
42. Zachary S. Brown, William Detmold, Stefan Meinel, Kostas Orginos, Charmed bottom baryon spectroscopy from lattice QCD. *Phys. Rev. D* **90**(9), 094507 (2014)
43. Stefan Meinel, Excited-state spectroscopy of triply-bottom baryons from lattice QCD. *Phys. Rev. D* **85**, 114510 (2012)
44. Raul A. Briceno, Huey-Wen. Lin, Daniel R. Bolton, Charmed-Baryon Spectroscopy from Lattice QCD with $N_f = 2 + 1 + 1$ Flavors. *Phys. Rev. D* **86**, 094504 (2012)
45. M. Padmanath, R.G. Edwards, N. Mathur, M. Peardon, Spectroscopy of triply-charmed baryons from lattice. *Phys. Rev. D* **90**(7), 074504 (2014)
46. Y. Namekawa et al., Charmed baryons at the physical point in 2+1 flavor lattice QCD. *Phys. Rev. D* **87**(9), 094512 (2013)
47. C. Alexandrou, V. Drach, K. Jansen, C. Kallidonis, G. Koutsou, Baryon spectrum with $N_f = 2 + 1 + 1$ twisted mass fermions. *Phys. Rev. D* **90**(7), 074501 (2014)
48. K.U. Can, G. Erkol, M. Oka, T.T. Takahashi, Look inside charmed-strange baryons from lattice QCD. *Phys. Rev. D* **92**(11), 114515 (2015)
49. D. Binosi, L. Theußl, JaxoDraw: a graphical user interface for drawing Feynman diagrams. *Comput. Phys. Commun.* **161**(1–2), 76–86 (2004)