



## Correction to: Photoelectrochemical sensing of tannic acid based on the use of TiO<sub>2</sub> sensitized with 5-methylphenazinium methosulfate and carboxy-functionalized CdTe quantum dots

Fernanda Maria dos Reis Lima<sup>1</sup> · André da Silva Freires<sup>1</sup> · Neuma das Mercês Pereira<sup>2</sup> · Glaura Goulart Silva<sup>2</sup> · Cláudia Quintino da Rocha<sup>1</sup> · Flavio Santos Damos<sup>1</sup> · Rita de Cássia Silva Luz<sup>1</sup>

Published online: 29 November 2018  
© Springer-Verlag GmbH Austria, part of Springer Nature 2018

### Correction to: Microchimica Acta

<https://doi.org/10.1007/s00604-018-3047-4>

The published version of this article, unfortunately, contains an error in that Fig. 2 shows the same plots as Fig. 3. The correct Fig. 2 is shown below.

---

The online version of the original article can be found at <https://doi.org/10.1007/s00604-018-3047-4>

✉ Rita de Cássia Silva Luz  
rita.luz@ufma.br

<sup>1</sup> Department of Chemistry, Federal University of Maranhão, São Luís, MA 65080-805, Brazil

<sup>2</sup> Department of Chemistry, Federal University of Minas Gerais, Belo Horizonte, MG 31270-901, Brazil

**Fig. 2** Photocurrent response of PMS/Nf/FTO (a), CdTe QDs/Nf/FTO (b), TiO<sub>2</sub>-PMS/Nf/FTO (c) and TiO<sub>2</sub>-PMS/CdTe QDs/Nf/FTO (d) in the absence (red lines) and presence (black lines) of 700 μmol L<sup>-1</sup> TA. Experiments carried in 0.1 mol L<sup>-1</sup> PB (pH 7.0). E<sub>app</sub> = +0.4 V vs Ag/AgCl

