

## Erratum to: The IMPACT prognosis calculator used in patients with severe traumatic brain injury treated with an ICP-targeted therapy

Magnus Olivecrona · Lars-Owe D. Koskinen

Published online: 12 August 2012  
© Springer-Verlag 2012

**Erratum to: Acta Neurochirurgica**  
**DOI 10.1007/s00701-012-1351-z**

The article contains two errors.

The first is found in the *Conclusion* section of the Abstract: The second last sentence reads: “We conclude that we have to initiate treatment in all patients with blunt sTBI and an initial ICP > 10 mmHg.”

This sentence should read: “We conclude that we have to initiate treatment in all patients with blunt sTBI and an initial cerebral perfusion pressure (CPP) ≥ 10 mmHg.”

In analogy with this the second error is found in the Materials and methods section’s last sentence; which reads: “Patients with dilated, fixed pupils and/or GCS 3 were included provided that their first measured ICP was 10 mmHg or higher”.

This sentence should read: “Patients with dilated, fixed pupils and/or GCS 3 were included provided that their first measured CPP was 10 mmHg or higher”

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00701-012-1351-z>.

---

M. Olivecrona · L.-O. D. Koskinen  
Department of Pharmacology and Clinical Neuroscience,  
Neurosurgery, Umeå University,  
Umeå, Sweden

L.-O. D. Koskinen  
e-mail: [lars-owe.koskinen@neuro.umu.se](mailto:lars-owe.koskinen@neuro.umu.se)

M. Olivecrona (✉)  
Department of Neurosurgery, Umeå University,  
SE 901 85 Umeå, Sweden  
e-mail: [magnus.olivecrona@neuro.umu.se](mailto:magnus.olivecrona@neuro.umu.se)