Correction to: Revision and complication rates in adult shunt surgery: a single institution study

Nadia Mansoor^{1,2} · Ole Solheim^{1,3} · Oddrun A. Fredriksli^{1,3} · Sasha Gulati^{1,3}

Published online: 19 January 2021

© Springer-Verlag GmbH Austria, part of Springer Nature 2021

Acta Neurochirurgica (2020), Oct 31. Online ahead of print
<https://doi.org/10.1007/s00701-020-04526-z>

The odds ratio was given as 0.97 and 95 % CI: 0.9, 1.0. The correct odds ratio and 95 % CI is 0.3 and 0.0–5.1, respectively. The correct table 1 is shown below:

In the original table 1 there was an error in regard to odds ratio and the 95 % confidence interval for the ICH group (non-revision group versus revision group).

Table 1 Patient demographics

	All patients		Non-revision group		Revision group		p-value	OR	95 % CI
	n=227	%	n = 180	%	n = 47	%			
Age: mean, yrs (±std)	58.4 (15.6)		59.8 (14.9)		53.0 (16.9)		0.014¹		1.4, 12.3
Gender: Male	104	45.8	87	48.3	17	36.1	0.136	1.7	0.9, 3.2
Status (end of follow-up): Dead	71	31.3	62	34.4	9	19.1	0.044	0.45	0.2, 1.0
Smoker: Yes	71 ²	37.0	53 ³	35.6	18 ⁴	41.9	0.452	1.3	0.7, 2.6
Comorbidities									
Hypertension	74	32.6	63	35.0	11	23.4	0.131	0.57	0.3, 1.2
Diabetes	23	10.1	20	11.1	3	6.4	0.339	0.55	0.2, 1.9
COPD	26	11.5	22	12.2	4	8.5	0.477	0.67	0.2, 2.0
Cerebrovascular disease	22	9.7	16	8.9	6	12.8	0.424	1.5	0.6, 4.1
Cardiovascular disease	29	12.8	25	13.9	4	8.5	0.325	0.6	0.2, 1.7
Tumour or malignancy	34	15.0	29	16.1	5	10.6	0.349	0.62	0.2, 1.7
ASA > 2	73 ⁵	50.3	65 ⁶	55.1	8 ⁷	29.6	0.017	0.3	0.1, 0.9
Aetiology/Indication									
SAH	81	35.7	64	35.6	17	36.2	0.938	1.0	0.5, 2.0
NPH	55	24.2	50	27.8	5	10.6	0.015	0.3	0.1, 0.8
Tumour or malignancy	43	18.9	36	20.0	7	14.9	0.426	0.7	0.3, 1.7
Unspecified HC	11	4.8	6	3.3	5	10.6	0.038	3.5	1.0, 11.9
ICH	7	3.1	7	3.9	0	0	0.170	0.3	0.0, 5.1
Traumatic	8	3.5	6	3.3	2	4.3	0.760	1.3	0.6, 6.6
Meningitis	3	1.3	2	1.1	1	2.1	0.587	1.9	0.2, 21.8
Chiari I	7	3.1	3	1.7	4	8.5	0.016	5.5	1.2, 25.4
Other	6	2.6	3	1.7	3	6.4	0.073	3.0	0.7, 13.9
Pseudotumor cerebri	6	2.6	3	1.7	3	6.4	0.073	4.0	0.8, 20.6

¹ Mann-Whitney U = 3249, p = 0.014. ² n = 192. ³ n = 149. ⁴ n = 43. ⁵ n = 145. ⁶ n = 118. ⁷ n = 27.

OD: odds ratio, CI: confidence interval, COPD: chronic obstructive pulmonary disease, SAH: subarachnoid haemorrhage, NPH: normal pressure hydrocephalus, HC: hydrocephalus, ICH: intracerebral haemorrhage. Age both groups: median 61, min 18, max 89, range 71.

The online version of the original article can be found at <https://doi.org/10.1007/s00701-020-04526-z>

✉ Nadia Mansoor
 nadia.mauland.mansoor@stolav.no

² Nevroklinikken, St. Olavs Hospital, 7006 Trondheim, Norway

³ Department of Neuromedicine and Movement Science, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

¹ Department of Neurosurgery, St. Olavs University Hospital, Trondheim, Norway