Correction to: Evidence-based treatment of neurogenic orthostatic hypotension and related symptoms

Sabine Eschlböck¹ · Gregor Wenning¹ · Alessandra Fanciulli¹

Published online: 28 November 2017

© Springer-Verlag GmbH Austria, part of Springer Nature 2017

Correction to: J Neural Transm

<https://doi.org/10.1007/s00702-017-1791-y>

Unfortunately, Author has now noticed two mistakes in Fig. 1. Author has written “peripher” instead of “peripheral” and “Secondary” instead of “Secondary”.

The revised figure is placed in the following page.

The online version of the original article can be found under
<https://doi.org/10.1007/s00702-017-1791-y>.

✉ Alessandra Fanciulli
alessandra.fanciulli@i-med.ac.at

¹ Department of Neurology, Medical University of Innsbruck,
Anichstrasse 35, 6020 Innsbruck, Austria

Fig. 1 Primary and secondary causes of nOH according to the site of lesion—schematic representation. *AAG* autoimmune autonomic ganglionopathy, *AAN* autoimmune autonomic neuropathy, *ADLD* autosomal dominant leukodystrophy, *CRF* chronic renal failure, *DBH* deficiency dopamine- β -hydroxylase deficiency, *DM* diabetes mellitus,

FD familial dysautonomia (=hereditary sensory and autonomic neuropathy type III, Riley-Day syndrome), *LBD* Lewy-body dementia, *MS* multiple sclerosis, *MSA* multiple system atrophy, *nOH* neurogenic orthostatic hypotension, *PAF* pure autonomic failure, *PD* Parkinson's disease