



Correction to: An optimum initial manifold for improved skill and lead in long-range forecasting of monsoon variability

K. C. Gouda¹ · S. Joshi² · Nagaraj Bhat³

Published online: 2 April 2021

© Springer-Verlag GmbH Austria, part of Springer Nature 2021

Correction to: Theoretical and Applied Climatology <https://doi.org/10.1007/s00704-021-03589-x>

The original version of this article unfortunately contained a mistake. The presentation of Table 2 was incorrect. The corrected table is given below.

The original article has been corrected.

Table 2 Comparative evaluation of 9 initial manifolds (ensembles) with two test ensembles for 12 extreme (amplitude of anomaly in all-India seasonal rainfall > 5%) years. The number in the bracket for each year indicates the observed % of anomaly for the year. “y” indicates that the simulation and observation values of rainfall anomalies are in phase

Ensemble	1980 (12.5)	1982 (-9)	1986 (-7)	1987 (-13)	1988 (13)	1990 (6)	1994 (11)	1998 (7)	1999 (-9)	2000 (-7)	2002 (-18)	2003 (9)	No of year
1	y	y	y	y			y	y	y	y			8
2	y		y	y			y	y	y	y		y	8
3	y		y	y			y	y	y	y		y	8
4 (OIM)	y	y	y	y			y	y	y	y		y	9
5	y		y	y		y	y	y	y	y			8
6	y	y	y	y		y	y	y	y	y		y	10
7		y	y	y		y		y	y	y		y	8
8		y	y	y			y	y	y	y		y	8
9		y	y	y			y	y	y	y			7
CE	y	y	y	y	y		y	y		y	y	y	10
LE	y		y	y			y	y	y	y			7

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00704-021-03589-x>

✉ K. C. Gouda
kcgouda@csir4pi.in

¹ CSIR Fourth Paradigm Institute (Formerly Centre for Mathematical Modelling and Computer Simulation), Wind Tunnel Road, Bangalore 560 037, India

² Government Degree College, Kanda, Kumaun University, Nainital, Uttarakhand, India

³ Shri Madhwa Vadiraja Institute of Technology and Management, Vishwothama Nagar, Bantakal, Karnataka, India