

Erratum to: ω -Amidase: an underappreciated, but important enzyme in L-glutamine and L-asparagine metabolism; relevance to sulfur and nitrogen metabolism, tumor biology and hyperammonemic diseases

Arthur J. L. Cooper¹ · Yevgeniya I. Shurubor^{2,3} · Thambi Dorai⁴ · John T. Pinto¹ · Elena P. Isakova³ · Yulia I. Deryabina³ · Travis T. Denton⁵ · Boris F. Krasnikov^{1,3}

Published online: 27 October 2015
© Springer-Verlag Wien 2015

Erratum to: Amino Acids DOI 10.1007/s00726-015-2061-7

In the original publication of the article, Fig. 1 and Acknowledgments were wrongly published. The correct Fig. 1 and acknowledgments are given below.

Acknowledgments This study was supported by Russian Ministry of Science and Education (Grant No. 14.604.21.0116 awarded to BFK). The unique identification number of applied scientific research is RFMEFI60414X0116.

The online version of the original article can be found under doi:[10.1007/s00726-015-2061-7](https://doi.org/10.1007/s00726-015-2061-7).

✉ Arthur J. L. Cooper
Arthur_Cooper@nymc.edu

¹ Department of Biochemistry and Molecular Biology, New York Medical College, Valhalla, NY 10595, USA

² Department of Neurology, The Brain and Mind Research Institute, Weill Cornell Medical College, New York, NY 10021, USA

³ A.N. Bach Institute of Biochemistry, Russian Academy of Sciences, 33 Leninsky Prospekt, Moscow 119071, Russia

⁴ Department of Urology, New York Medical College, Valhalla, NY 10595, USA

⁵ Department of Pharmaceutical Sciences, College of Pharmacy, Washington State University, Spokane, WA 99210, USA




Fig. 1 Glutaminase II (a) and asparaginase II (b) pathways. Note the central importance of ω -amidase in hydrolyzing the α -keto acids derived from transamination of

both L-glutamine (i.e., α -ketoglutaramate, KGM) and L-asparagine (α -ketosuccinamate, KSM). *Gln Ts* glutamine transaminases, *Asn Ts* asparagine transaminases