

Erratum to: Pilot early intervention antenatal group program for pregnant women with anxiety and depression

Naomi Thomas · Angela Komiti · Fiona Judd

Published online: 4 September 2014
© Springer-Verlag Wien 2014

Erratum to: Arch Womens Ment Health DOI 10.1007/s00737-014-0447-2

Figure 1 in the original version of this article was incorrectly placed in the method section with the description of the EPDS rather than as a part of the results of the study. The paragraph relating to the findings depicted in Figure 1 was unfortunately deleted from the results section. All information shown in the figure was correct. The description of the results in Figure 1 appears below.




Figure 1 shows the change in mean EPDS depression scores from pre-treatment ($M=16.2$, $SD=5.0$), post-treatment ($M=10.5$, $SD=4.0$) and at two months postpartum follow up ($M=8.3$, $SD=4.2$). Figure 1 shows a significant change in depression scores between pretreatment and post-treatment ($p<.001$) and pre-treatment and two-months postpartum ($p<.001$). A significant difference was also observed between post-treatment (T2) and two month postpartum follow-up (T3) depression scores ($p=.006$).

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00737-014-0447-2>.

N. Thomas · A. Komiti · F. Judd (✉)
Centre for Women's Mental Health, Royal Women's Hospital,
Locked Bag 300, Parkville, VIC, Australia
e-mail: fionakj@unimelb.edu.au

A. Komiti · F. Judd
Department of Psychiatry, University of Melbourne, Parkville,
VIC 3052, Australia