Correction to: Osteoprotegerin/bone morphogenetic protein 2 combining with collagen sponges on tendon-bone healing in rabbits

Bing Wei¹ · Chao Wang¹ · Cheng Yan¹ · Bushun Tang¹ · Xiaofei Yu² · Hui Zhang³ · Lixia Tang³ · Qing Wang¹

Published online: 17 December 2021

© The Japanese Society Bone and Mineral Research 2021

Correction to:

Journal of Bone and Mineral Metabolism

(2020) 38:432–441

<https://doi.org/10.1007/s00774-019-01078-w>

In the original publication of the article, Fig. 4A and 4B are duplicate. The correct Fig. 4A is provided in this correction.

The original article can be found online at <https://doi.org/10.1007/s00774-019-01078-w>.

✉ Lixia Tang
tanglixia138@163.com

✉ Qing Wang
wangqinglijun@163.com

¹ Department of Orthopedics, The First People's Hospital of Yongkang, Jinhua 321300, People's Republic of China

² Department of Pathology, The First People's Hospital of Yongkang, Jinhua 321300, People's Republic of China

³ Department of General Diseases, The First People's Hospital of Yongkang, Jinhua 321300, People's Republic of China

In addition, the term “at 12 weeks” were incorrectly included in the captions of Fig. 4 and 5.

The correct captions should read as below.

Caption of Fig. 4: Tissue morphology post operation (Hematoxylin and Eosin, ×100). *T* tendon, *B* bone, *UFC* uncalcified fibrocartilage, *CFC* calcified fibrocartilage, *TM* Tide mark; A black arrow: Sharpey-like collagen fibers. **a** the blank control group; **b** the OPG/BMP-2; **c** the CS group; **d** the OPG/BMP-2/CS group.

Caption of Fig. 5: M–T staining (×100) shows fibrocartilage and collagen fiber formation around the tendon. *T* tendon, *B* bone, *FC* fibrocartilage, *CF* collagen fibers. **a** the blank control group; **b** the OPG/BMP-2; **c** the CS group; **d** the OPG/BMP-2/CS group.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.