



Correction: Movement control during one-leg standing is important for the bone mineral density maintenance or improvement

Takuya Umehara¹ · Akinori Kaneguchi¹ · Keita Watanabe² · Nobuhisa Katayama² · Daisuke Kuwahara³ · Ryo Kaneyashiki³ · Nobuhiro Kito¹ · Masayuki Kakehashi⁴

Published online: 27 August 2022

© The Japanese Society Bone and Mineral Research 2022

Correction:

Journal of Bone and Mineral Metabolism

<https://doi.org/10.1007/s00774-022-01348-0>

In the original publication of the article, under the results section, the following section heading “Factors influencing BMD maintenance or improvement at 6 months (after propensity score matching)” was missing and included in this correction.

The original article has been updated.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00774-022-01348-0>.

✉ Takuya Umehara
start.ume0421@gmail.com

- ¹ Department of Rehabilitation, Faculty of Rehabilitation, Hiroshima International University, Kurose-Gakuendai 555-36, Higashi-Hiroshima, Hiroshima, Japan
- ² Department of Rehabilitation, Kure Kyosai Hospital, Nishichuo 2-3-28, Kure, Hiroshima, Japan
- ³ Department of Rehabilitation, Saiseikai Kure Hospital, Sanjo 2-1-13, Kure, Hiroshima, Japan
- ⁴ Department of Health Informatics, Graduate School of Biomedical and Health Sciences, Hiroshima University, Kasumi 1-2-3, Hiroshima Minami-ku, Hiroshima, Hiroshima, Japan