

## Erratum to: The tetraheme cytochrome from *Shewanella oneidensis* MR-1 shows thermodynamic bias for functional specificity of the hemes

Bruno M. Fonseca · Ivo H. Saraiva · Catarina M. Paquete · Claudio M. Soares · Isabel Pacheco · Carlos A. Salgueiro · Ricardo O. Louro

Published online: 26 January 2011  
© SBIC 2011

**Erratum to: J Biol Inorg Chem (2009) 14:375–385**  
DOI 10.1007/s00775-008-0455-7

The original publication did not report the final calculated values in Table 2. This only has a significant consequence

for reported geometry of the axial ligands of heme IV and changes one of the 16 panels of Fig. 1.

The updated versions of Table 2 and Fig. 1 are given below.

**Table 2**

	Heme I		Heme II		Heme III		Heme IV	
	NMR	X-ray	NMR	X-ray	NMR	X-ray	NMR	X-ray
$\theta$ (°)	−13.0 (0.4)	−10.4	−46 (2)	−72.9	−28.3 (0.4)	−33.8	−8.5 (0.5)	−0.9
$\beta$ (°)	0	18.2	71	65.7	0	6.3	53	56.8
$\Delta E$ (kJ/mol)	6.0 (0.2)		1.32 (0.05)		6.3 (0.2)		3.5 (0.1)	

The online version of the original article can be found under doi:10.1007/s00775-008-0455-7.

B. M. Fonseca · I. H. Saraiva · C. M. Paquete ·  
C. M. Soares · I. Pacheco · R. O. Louro (✉)  
Instituto de Tecnologia Química e Biológica,  
Universidade Nova de Lisboa,  
Av. da Republica (EAN),  
2780-157 Oeiras, Portugal  
e-mail: louro@itqb.unl.pt

C. A. Salgueiro  
Requimte-CQFB, Departamento de Química,  
Faculdade de Ciências e Tecnologia,  
Universidade Nova de Lisboa,  
Quinta da Torre,  
2829-516 Caparica, Portugal

Fig. 1

