Correction to: A copper(II)-binding triazole derivative with ionophore properties is active against *Candida* spp.

A. Gaspar-Cordeiro¹ · S. da Silva¹ · M. Aguiar² · C. Rodrigues-Pousada¹ · H. Haas² · L. M. P. Lima¹ · C. Pimentel¹

Published online: 29 May 2021

© Society for Biological Inorganic Chemistry (SBIC) 2021

Correction to:

JBIC Journal of Biological Inorganic Chemistry
(2020) 25:1117–1128
<https://doi.org/10.1007/s00775-020-01828-6>

The original version of this article was published with two errors in materials and methods and on Table 2. Instead of referring to MIC 50, it is referred as MIC 90.

The corrected version is given below.

Antifungal susceptibility testing in broth microdilution assays

Minimal inhibitory concentration (MIC) of *Candida* sp. and *Aspergillus fumigatus* was determined by conducting broth microdilution assays in accordance to the CLSI (Clinical Laboratory and Standards Institute) standard methods M27-A3 and M38 for yeasts and filamentous fungi, respectively [1, 2]. Drug concentrations ranged from 125 to 0.24 μ M

and CuCl_2 concentrations from 5 to 0.01 mM. Growth in RPMI medium was recorded after 48 h at 30 °C (*Candida albicans*) or 37 °C (*Candida glabrata* and *Aspergillus fumigatus*), by measuring OD600 (*Candida* sp.) or using phase contrast microscopy (*Aspergillus fumigatus*). For the latter, an inverted Leica DM IL LED microscope (Leica Microsystems) was used and images were captured using an AxioCam MR3 camera (Carl Zeiss GmbH). The growth condition without drug was used as the normalization condition, after background (RPMI medium) subtraction. The MIC was defined as the concentrations of the drug between which the relative ODs fall below 50% of the no-drug condition. At least, three independent assays were performed for each chemical and strain.

The original article can be found online at <https://doi.org/10.1007/s00775-020-01828-6>.

✉ L. M. P. Lima
llima@itqb.unl.pt

✉ C. Pimentel
pimentel@itqb.unl.pt

¹ Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, Av. República, 2780-157 Oeiras, Portugal

² Division of Molecular Biology/Biocenter, Innsbruck Medical University, Innrain 80-82, 6020 Innsbruck, Austria

Table 2 In vitro susceptibility of *Candida glabrata*, *Candida albicans* and *Aspergillus fumigatus* (wild-type and $\Delta crpA$ mutant strains) to **4**, **Cu.4**, CuCl_2 , **5**, **Cu.5** and fluconazole (FLC)

	MIC (μM)						
	4	Cu.4	CuCl_2	5	Cu.5	1	FLC
<i>C. glabrata</i>	31.3–15.6	62.5–31.3	162.5–81.3	> 125	125–62.5	> 125	105–52.5
<i>C. albicans</i>	62.5–31.3	62.5–31.3	1300–650	> 125	> 125	> 125	3.3–1.6
<i>A. fumigatus</i>	> 125	> 125	5000–2500	> 125	> 125	> 125	> 416
<i>A. fumigatus</i> $\Delta crpA$	> 125	78–39	313–156	> 125	> 125	> 125	> 416

References

1. Institute CLSICaLS (2017) Reference method for broth dilution antifungal susceptibility testing of filamentous fungi. CLSI standard M38, 3rd edn
2. Institute CLSICaLS (2017) Reference method for broth dilution antifungal susceptibility testing of yeasts. CLSI standard M27, 4th edn

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.