

Erratum to: Transverse dental arch relationship at 9 and 12 years in children with unilateral cleft lip and palate treated with infant orthopedics: a randomized clinical trial (DUTHCLEFT)

R. L. M. Noverraz¹ · M. A. Disse¹ · E. M. Ongkosuwito² · A. M. Kuijpers-Jagtman² · C. Prahl¹

Published online: 16 August 2015
© Springer-Verlag Berlin Heidelberg 2015

Erratum to: Clinical Oral Investigations DOI 10.1007/s00784-015-1451-2

The original version of the above article contained a mistake due to incorrect alignment of entries in Tables 11 and 14 during production. The entries are presented correctly below.

Table 11 Evaluation of the total arch constriction in pairs

	Center	Center	Mean	95 % CI		P
				Lower	Upper	
9 years	1	2	1.63	-2.09	5.34	0.842
	1	3	-1.71	-5.61	2.18	0.836
	2	3	-3.34	-6.73	.05	0.054
12 years	1	2	4.85	0.53	9.17	0.023*
	1	3	-4.39	-4.02	4.95	1.000
	2	3	0.47	-8.33	-0.44	0.025*

Mean mean total arch constriction score (points), 95% CI 95 % confidence interval of the difference

* $p < 0.05$

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00784-015-1451-2>.

✉ C. Prahl
c.prahl@acta.nl

¹ Department of Orthodontics, Academic Centre for Dentistry Amsterdam (ACTA), Gustav Mahlerlaan 3004, 1081 LA Amsterdam, The Netherlands

² Department of Orthodontics and Craniofacial Biology and Cleft Palate Craniofacial Centre, Radboud University Medical Centre, Nijmegen, The Netherlands

Table 14 Evaluation of age of hard palate closure with alveolar bone grafting in pairs

Center	Center	Mean diff	95 % CI		P
			Lower	Upper	
1	2	-1.55	-2.43	0.68	0.000***
1	3	-0.99	-1.88	-0.11	0.023
2	3	0.56	-0.20	1.32	0.218

Mean diff mean difference in age of hard palate closure with alveolar bone grafting (years), 95% CI 95 % confidence interval of the difference

*** $p < 0.001$