

Erratum to: Early emotional and behavioral difficulties and adult educational attainment: an 18-year follow-up of the TEMPO study

Ariella Zbar¹ · Pamela J. Surkan² · Eric Fombonne³ · Maria Melchior⁴

Published online: 25 January 2017
© Springer-Verlag Berlin Heidelberg 2017

**Erratum to: Eur Child Adolesc Psychiatry (2016)
25:1141–1143
DOI 10.1007/s00787-016-0858-x**

In the original publication, the first column of Table 1 presented some erroneous numbers under the low educational attainment heading. The corrected Table 1 is given below.

The online version of the original article can be found under doi:[10.1007/s00787-016-0858-x](https://doi.org/10.1007/s00787-016-0858-x).

✉ Maria Melchior
maria.melchior@inserm.fr

- ¹ Public Health and Preventive Medicine, Queen's University, Kingston, ON, Canada
- ² Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA
- ³ Department of Psychiatry, Oregon Health and Science University, Portland, OR, USA
- ⁴ Sorbonne Universités, UPMC Univ Paris 06, INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique (IPLESP UMRS 1136), 75012 Paris, France

Table 1 Socio-demographic and youth internalizing and externalizing symptoms in relation to low educational attainment in early adulthood: TEMPO study, 1991–2009 (bivariate and multivariate logistic regression models, OR, 95% CI)

| | Low educational attainment <i>N</i> , % | | Odds ratio | |
|---------------------------------|---|----------------------|--|--|
| | No <i>N</i> = 536 | Yes <i>N</i> = 77 | Crude (95% CI) <i>p</i> | Adjusted (95% CI) <i>p</i> |
| Age >30 years in 2009 | 229 (42.7) | 53 (40.8) | 0.92 (0.62–1.36) <i>p</i> = 0.67 | – |
| Male | 203 (15.5) | 64 (49.2) | 1.59 (1.08–2.34) <i>p</i> = 0.02 | 1.30 (0.86–1.98) <i>p</i> = 0.21 |
| Childhood academic difficulties | 125 (23.3) | 59 (45.4) | 2.73 (1.83–4.07) <i>p</i> < 0.0001 | 2.40 (1.57–3.66) <i>p</i> < 0.0001 |
| Family income <1982€ per month | 63 (18.8) | 30 (23.1) | 2.24 (1.38–3.64) <i>p</i> = 0.001 | 2.15 (1.29–3.58) <i>p</i> = 0.003 |
| Externalizing symptoms (1991) | 54 (10.1) | 21 (16.4) | 1.75 (1.01–3.01) <i>p</i> = 0.05 | 1.34 (0.74–2.39) <i>p</i> = 0.33 |
| Internalizing symptoms (1991) | 64 (11.9) | 26 (20.0) | 1.85 (1.15–3.06) <i>p</i> = 0.002 | 1.75 (1.00–3.02) <i>p</i> = 0.05 |
| Externalizing symptoms (1999) | 54 (10.1) | 17 (13.2) | 1.35 (0.76–2.42) <i>p</i> = 0.31 | – |
| Internalizing symptoms (1999) | 88 (16.5) | 15 (11.5) | 0.66 (0.37–1.19) <i>p</i> = 0.17 | 0.54 (0.29–1.02) <i>p</i> = 0.06 |