

Genomic analysis of *Acidianus hospitalis* W1 a host for studying crenarchaeal virus and plasmid life cycles

Xiao-Yan You · Chao Liu · Sheng-Yue Wang · Cheng-Ying Jiang · Shiraz A. Shah · David Prangishvili · Qunxin She · Shuang-Jiang Liu · Roger A. Garrett

Received: 4 March 2011 / Accepted: 26 April 2011 / Published online: 24 May 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract The *Acidianus hospitalis* W1 genome consists of a minimally sized chromosome of about 2.13 Mb and a conjugative plasmid pAH1 and it is a host for the model filamentous lipothrixvirus AFV1. The chromosome carries three putative replication origins in conserved genomic regions and two large regions where non-essential genes are clustered. Within these variable regions, a few orphan *orfB* and other elements of the IS200/607/605 family are concentrated with a novel class of MITE-like repeat elements. There are also 26 highly diverse *vapBC* antitoxin–toxin gene pairs proposed to facilitate maintenance of local chromosomal regions and to minimise the impact of environmental

stress. Complex and partially defective CRISPR/Cas/Cmr immune systems are present and interspersed with five *vapBC* gene pairs. Remnants of integrated viral genomes and plasmids are located at five intron-less tRNA genes and several non-coding RNA genes are predicted that are conserved in other *Sulfolobus* genomes. The putative metabolic pathways for sulphur metabolism show some significant differences from those proposed for other *Acidianus* and *Sulfolobus* species. The small and relatively stable genome of *A. hospitalis* W1 renders it a promising candidate for developing the first *Acidianus* genetic systems.

Keywords Toxin–antitoxin VapBC · CRISPR · Sulphur metabolism · *OrfB* element · MITE

Communicated by L. Huang.

X.-Y. You and C. Liu contributed equally to this work.

X.-Y. You · C.-Y. Jiang · S.-J. Liu (✉)
State Key Laboratory of Microbial Resources and Center for Environmental Microbiology, Institute of Microbiology, Chinese Academy of Sciences, Bei-Chen-Xi-Lu No. 1 Chao-Yang District, Beijing 100101, People's Republic of China
e-mail: liusj@sun.im.ac.cn

C. Liu · S. A. Shah · Q. She · R. A. Garrett (✉)
Archaea Centre, Department of Biology, Copenhagen University, Ole Maaløes Vej 5, 2200 N Copenhagen, Denmark
e-mail: garrett@bio.ku.dk

S.-Y. Wang
Shanghai-MOST Key Laboratory of Health and Disease Genomics, Chinese National Human Genome Center, Shanghai, People's Republic of China

D. Prangishvili
Molecular Biology of the Gene in Extremophiles Unit, Institut Pasteur, rue Dr Roux 25, 75724 Paris Cedex, France

Introduction

The *Acidianus* genus consists of acidothermophiles which grow optimally and slowly in the temperature range 65–95°C and at pH 2–4 and belongs to the order Sulfolobales. *Acidianus* species are chemolithoautotrophic and facultatively anaerobic and are generally versatile physiologically. Depending on the culturing conditions, they can either reduce S⁰ to H₂S, catalysed by a sulphur reductase and hydrogenase, or oxidise S⁰ to H₂SO₄ utilising the sulphur oxygenase-reductase holoenzyme (Kletzin 1992, 2007). In contrast to several *Sulfolobus* species, the genomic properties of an *Acidianus* species have not been analysed. The Sulfolobales have been a rich source of genetic elements, including novel conjugative plasmids (Prangishvili et al. 1998; Greve et al. 2004) and several exceptional and diverse viruses many of which have now been classified into eight new viral families (Rachel et al. 2002; Prangishvili et al. 2006; Lawrence et al. 2009).

Acidianus hospitalis W1 is the first *Acidianus* strain to be isolated carrying a conjugative plasmid pAH1 which is a member of the plasmid family predicted to generate an archaea-specific conjugative apparatus (Greve et al. 2004; Basta et al. 2009). These plasmids are also integrative elements and in an encapsulated state have been implicated in facilitating chromosomal DNA conjugation for some *Sulfolobus* species (Chen et al. 2005b). *A. hospitalis* is also a viable host for the model *Acidianus* alpha lipothrixvirus AFV1, a filamentous virus carrying exceptional claw-like structures at its termini which is currently the subject of detailed structural studies (Bettstetter et al. 2003; Goulet et al. 2009). Infection of *A. hospitalis* with AFV1 was shown to lead to a loss of the plasmid pAH1 and this contrasts with observations in bacteria where endogenous plasmids tend to determine the fate of an incoming phage (Basta et al. 2009).

In order to study further the metabolic capability of an *Acidianus* species and to examine the molecular mechanisms involved in virus–plasmid–host interactions, it was important to sequence and annotate the *A. hospitalis* genome. To date, most genomic studies of the Sulfolobales have concentrated on *Sulfolobus* species that have revealed relatively large genomes generally exhibiting high levels of transposable and integrated genetic elements, as well as considerable genetic diversity (Guo et al. 2011). Analysis of the *A. hospitalis* genome revealed a minimally sized chromosome that appeared relatively stable with few transposable elements and no evidence of recent integration events, apart from the reversible integration of pAH1 into a tRNA^{Arg} gene (Basta et al. 2009). Potentially, therefore, *A. hospitalis* W1 could provide a suitable host for developing genetic systems for the *Acidianus* genus.

Materials and methods

Genome sequencing and gap closure

Genomic DNA of *A. hospitalis* was sequenced using a Roche 454 Genome Sequencer FLX instrument (Titanium) with an average 19-fold coverage. All useful reads were initially assembled into seven contigs (>500 bp) using the Newbler assembler software (<http://www.454.com/>). Gaps were closed by a Multiplex PCR strategy and PCR products were gel purified and sequenced using an ABI3730 DNA sequencer. Raw sequence data were assembled into contigs using phred/phrap/consed software and the final consensus quality for each base was above 30 (<http://www.phrap.org>).

Sequence analysis and gene annotation

Initially, ORFs were predicted using the programmes Glimmer and FgeneSB and protein function predictions

were obtained from the following searches: (1) homology searches in the GenBank (<http://www.ncbi.nlm.nih.gov/>) and UniProt protein (<http://www.ebi.ac.uk/uniprot/>) databases, (2) function assignment searches in the *Sulfolobus* database (<http://www.Sulfolobus.org/>), and (3) domain or motif searches in the local CDD database (<http://www.ncbi.nlm.nih.gov/cdd/>), the InterPro and the Pfam databases. The KEGG database (<http://www.genome.jp/kegg/>) was used to reconstruct metabolic pathways in silico. Membrane proteins were predicted by Phobius, TMHMM and ConPred II programmes. Secretory proteins were divided into two groups; those with a signal peptide were predicted using the SignalP 3.0 (<http://www.cbs.dtu.dk/services/SignalP/>) and non-classical secretory proteins, lacking a signal peptide, were predicted by the SecretomeP 2.0 programme (<http://www.cbs.dtu.dk/services/SecretomeP/>). Transporters were predicted by searching the TCDB database (<http://www.tcdp.org>) using BLASTP with *E* values lower than 1e-05. Insertion sequence (IS) elements and transposases were identified by BLASTN searches against the IS Finder database (<http://www-is.biotoul.fr/>). The MITE-like elements were detected using the programme LUNA (Brügger K, unpublished). Potential frameshifts were checked by sequencing after manual annotation and any remaining frameshifts were considered to be authentic. tRNA genes and their introns were identified using tRNAScan-SE (Lowe and Eddy 1997). All annotations were manually curated using Artemis software (Rutherford et al. 2000). Start codons for single genes and first genes of *Sulfolobus* operons were generally located 25–30 bp downstream from the archaeal hexameric TATA-like box. Only genes within operons were preceded by Shine–Dalgarno motifs, where GGUG dominated (Torarinsson et al. 2005). Where alternative start codons occur, a selection was made on the basis of experimental data when available or on its location relative to a putative promoter and/or Shine–Dalgarno motif. The genome sequence accession number at Genbank/EMBL is CP002535.

Results

Genomic properties

The *A. hospitalis* genome consists of a circular chromosome of 2,137,654 bp and a circular conjugative plasmid pAH1 of 28,644 bp. The chromosome has a GC content of 34.2% and carries 2,389 predicted open reading frames (ORFs), of which about half are assigned putative functions with many of the conserved hypothetical proteins being archaea-specific or specific to the Sulfolobales. About 320 of the encoded proteins are putative membrane proteins and a further 182 are predicted to be secretory proteins.

The plasmid sequence is identical to that of the conjugative plasmid pAH1 isolated earlier from the *A. hospitalis* strain W1, except that it is 4 bp shorter (Basta et al. 2009).

Comparison of the *A. hospitalis* genome with those of other members of the Sulfolobales provided no evidence of extensive conservation of gene synteny, in contrast to that observed for large regions of several *Sulfolobus* genomes (Guo et al. 2011), and consistent with *A. hospitalis* being relatively distant phylogenetically from these strains (Basta et al. 2009). Nevertheless, the genome carries two major regions that are predicted to be relatively labile. They extend approximately from positions 75,000–444,500 and from 1,300,000–1,870,000 and carry most of the transposable elements, all of the CRISPR loci and *cas* and *cmr* family genes, most of the *vapBC* toxin–antitoxin gene pairs, and many genes involved in transport-related functions and metabolism, as well as a degenerate fuselloviral genome (Fig. 1). These two regions lack genes essential for informational processes including DNA replication, transcription and translation and they appear to constitute sites where non-essential genes are collected, interchanged, exchanged intercellularly and where genetic innovation may occur, similarly to a single variable region observed in several *Sulfolobus* genomes (Guo et al. 2011).

Three origins of chromosomal replication, demonstrated experimentally for *Sulfolobus* species (Robinson et al. 2004; Lundgren et al. 2004), were also predicted to occur in the *Acidianus* genome. The Y component of a Z curve analysis (Zhang and Zhang 2003) revealed two major peaks corresponding to the *cdc6-3* gene (Ahos0001), and the *whiP/ctd1* gene (Ahos1370) and a broader peak coinciding with the *cdc6-1* gene (Ahos0780) (Fig. 1), where the three genes encode putative replication initiators (Robinson and Bell 2007). The sequences of the *cdc6* genes and *whiP* gene are quite conserved relative to the *S. solfataricus* and

S. islandicus genomes, as is the synteny of the flanking genes except for the region immediately downstream from *cdc6-3*.

Integrated genetic elements

Integration of genetic elements, generally fuselloviruses or conjugative plasmids at tRNA genes, occurs commonly for genomes of the Sulfolobales (She et al. 1998; Guo et al. 2011). Most integration events occur via a reversible archaea-specific mechanism whereby the integrase gene partitions into two sections which border the integrated element and the N-terminal-encoding region carrying the *intN* sequence overlaps with the tRNA gene (Muskhelishvili et al. 1993). Elements that become encaptured within the chromosome subsequently degenerate and are gradually lost, but will nevertheless leave a trace because the *intN* fragment overlapping the tRNA gene is generally retained (She et al. 1998) (Table 1).

Earlier plasmid pAH1 was sequenced and shown to integrate reversibly into a tRNA^{Arg} gene (Basta et al. 2009). Genome sequencing of *A. hospitalis* revealed that a low fraction of reads matched to the junctions of the integrated plasmid whilst the majority matched the unpartitioned integrase gene of pAH1, consistent with both integrated and free forms being present in the culture. The integration site of pAH1 was located at genome positions 1,075,876–1,075,946 bp within the gene of tRNA^{Arg} [TCG] (Table 1). In addition, the chromosome carries remnants of integrated elements adjoining another five intron-less tRNA genes, each consisting of a few genes or pseudogenes (Table 1). Three derive from fuselloviruses, one from a pDL10-like plasmid of the pRN family of cryptic plasmids (Kletzin et al. 1999) and another originates from an unknown element (Table 1). Whether these all derive from single integration events remains unclear




Fig. 1 The Y component of a Z curve plot for the *A. hospitalis* chromosome showing the three putative replication origins. The positions of the *cdc6-3* gene (origin 2), *cdc6-1* gene (origin 3) and the *whiP/ctd1* gene (origin 1) are indicated as well as locations of the

ribosomal RNA genes, the CRISPR-based systems, transposable elements of the IS200/605/607 family, and *vapBC* antitoxin–toxin gene pairs

Table 1 Integration events at tRNA genes showing the numbers of residual integrated genes

tRNA	Intron	Ahos W1
Arg–TCG	No	pAH1
Pro–TGG	No	<i>intN</i> fragment
Glu–CTC	No	0986a–0988 fusellovirus
Arg–TCT	No	1232–1238 unknown element
Cys–GCA	No	1550–1558 plasmid pDL10
Leu–GAG	No	2147–2151 fusellovirus ASV1
1604–1609 kb (no tRNA)	–	1778–1786 fusellovirus SSV

ASV1 *Acidianus* spindle-shaped virus, SSV *Sulfolobus* spindle-shaped virus

because, in principle, successive integrations can occur at a given tRNA gene (Redder et al. 2009). An additional 8–10 genes and pseudogenes, most of which are fusellovirus-related, are clustered distantly from a tRNA gene and they may have become displaced from one of the three tRNA-integrated elements.

Transposable elements

The *A. hospitalis* genome carries five IS elements belonging to the IS200/607 family, only three of which carry intact transposase genes, and there are 11 copies of orphan *orfB* elements of the IS605 family, 10 of which carry intact *orfB* genes. None of these elements carry inverted terminal repeats and they all appear to be transposed by “cut-and-paste” mechanisms, with the *orfB* elements, at least, transposing via circular single stranded intermediates and inserting after TTAC sequences (Filée et al. 2007; Ton-Hoang et al. 2010).

Sulfolobus genomes generally carry IS elements from a wide variety of families most of which carry inverted terminal repeats and are mobilised by “copy-and-paste” mechanisms, and tend to be lost by gradual degeneration and not by deletion (Blount and Grogan 2005; Redder and Garrett 2006). None of these IS element classes were detected in the *A. hospitalis* genome and this suggests that the genome has rarely, if ever, taken up any of these IS element classes.

A new class of MITE-like elements

Although none of the MITE elements that are common to other *Sulfolobus* genomes were detected (Redder et al.

2001; Guo et al. 2011), the *A. hospitalis* genome carries 10 copies of a repeat sequence resembling a MITE-like element (Fig. 3). At one end, it carries a short open reading frame corresponding in amino acid sequence to the downstream end of an OrfB protein (Fig. 3). The conserved terminal sequence and the internal similarity to the *orfB* element suggests that it could be a transposable element. This supposition is reinforced by the presence of 10 full copies in the genome (and a few degenerate copies), and also by the presence of multiple copies in some *Sulfolobus* and other crenarchaeal genomes (unpublished data).

Non-coding RNAs

Many untranslated RNAs have been characterised experimentally for different *Sulfolobus* species using a variety of techniques including probing cellular RNA extracts for K-turn-binding motifs and generating cDNA libraries of total cellular RNA extracts, as well as numerous antisense RNAs (Tang et al. 2005; Omer et al. 2006; Wurtzel et al. 2010). Most of these RNAs were characterised for partial sequence and nucleotide length, and several were detected by more than one experimental approach. Based on the genome sequence comparisons and gene contexts, 23 putative conserved non-coding RNAs were annotated in the *A. hospitalis* genome. Genes for 12 C/D box RNAs were localised of which 7 were predicted to modify rRNAs, 2 to target tRNAs and a further 2 to modify unknown RNAs. In addition, a single copy of a gene for an H/ACA box RNA was located which together with aPus7 should generate pseudouridine-35 in *Sulfolobus* pre-tRNA^{Tyr} transcripts (Muller et al. 2009). However, in *A. hospitalis*, the aPus7 gene (Ahos0631) is degenerate. A further 10 genes were assigned to encode RNAs of unknown function. The relatively high conservation of sequence and gene synteny for these RNAs between *Sulfolobus* and *Acidianus* species underlines their potential functional importance.

Reading frame shifts and mRNA intron splicing

Examples of translational reading frame shifts yielding single polypeptides have been demonstrated experimentally for *S. solfataricus* P2 (Cobucci-Ponzano et al. 2010). For two of these, a transketolase (Ahos1219/1218) and a putative *O*-sialoglycoprotein endopeptidase (Ahos0695/0696), the *A. hospitalis* genes overlap in a similar way, and are likely to undergo translational frame shifts. Moreover, transcripts of the intron-carrying *cbf5* gene (Ahos0734/0735) are likely to undergo splicing at the mRNA level by the archaeal splicing enzyme complex (Ahos0689/0798/1417) as has been demonstrated experimentally for different crenarchaea (Yokobori et al. 2009).

Metabolic pathways

Genome analyses indicate the presence of versatile metabolic pathways in *A. hospitalis*. They suggest that it can grow autotrophically by fixing CO₂ or heterotrophically using yeast extract, as has been demonstrated experimentally (Basta et al. 2009). Genome analyses also revealed genes encoding sugar transporters and glycosidases suggesting that *A. hospitalis* can assimilate carbohydrates, such as starch, glucose, mannose and galactose. Moreover, enzymes are encoded that are implicated in energy generation from oxidising elemental sulphur, hydrogen sulphides and other reduced inorganic sulphide compounds, but not ferrous ions. However, no hydrogenase genes were detected suggesting that *A. hospitalis* cannot use H₂ as electron donor for growth.

Enzymes were identified for a complete TCA cycle that is important for generating different intermediates for the biosynthesis of many cellular components, as well as producing reduced electron carriers, such as NAD(P)H, reduced ferredoxin (Fd_R) and FADH₂. Formation of acetyl-CoA from pyruvate and the formation of succinyl-CoA from 2-oxoglutarate were predicted to be catalysed, respectively, by pyruvate ferredoxin oxidoreductase (Ahos 1949-1952) and 2-oxoglutarate ferredoxin oxidoreductase (Ahos0089/0090/0300/0301). Moreover, both enzymes were predicted to use ferredoxin instead of NAD⁺ as a cofactor.


Genes encoding enzymes involved in pathways for fixing atmosphere N₂, or reducing nitrate and nitrite, as nitrogen sources were absent, as observed for other *Acidianus* species, and the genome analyses suggest that ammonium is an exclusive source of nitrogen that is

assimilated via formation of carbamoyl phosphate, glutamine and glutamate. Genes encoding putative carbamoyl phosphate synthetase (Ahos1106/1107), glutamine synthetase (Ahos0460, Ahos1272, Ahos2233) and glutamate dehydrogenase (Ahos0494) are present.

Sulphur metabolism

A. hospitalis encodes several enzymes involved in sulphur metabolism, including the oxidation and reduction of sulphur, the thiosulphate–tetrathionate cycle which generates sulphate, and the participation of sulphur in electron transport. However, genes for some sulphur metabolism enzymes, including sulphite-acceptor oxidoreductase, adenosine phosphosulphate reductase, sulphate adenylyl transferase and adenylylsulphate phosphate adenylyltransferase were not found which suggested that *A. hospitalis* has some pathways differing from those of other *Acidianus* and *Sulfolobus* species (Kletzin 2007). Therefore, based on the gene annotations, a model is presented for the proposed sulphur oxidation and reduction pathways in *A. hospitalis* (Fig. 2). Extracellular H₂S is oxidised by a secretory-type sulphide:quinone oxidoreductase (Ahos0513) and flavocytochrome *c* sulphide dehydrogenase (Ahos0188) to produce a surface layer of sulphur on the outer cell membrane. Elemental sulphur is then transported into the cell by putative-SH radical transporter(s) using an unknown mechanism. Subsequently, sulphur is oxidised by sulphur oxygenase-reductase (Ahos0131) to yield sulphite, thio-sulphate and hydrogen sulphide. Sulphite and elemental sulphur convert spontaneously and non-enzymatically to thiosulphate and elemental sulphur and, consistent with this mechanism, no candidate gene encoding sulphite:acceptor

Fig. 2 Model of pathways for oxidation and reduction of sulphur in *A. hospitalis* indicating the predicted functions of genes in the *A. hospitalis* genome and corresponding gene numbers are given for each step. The following abbreviations are used: *OM* outer membrane, *IM* inner membrane, *SQR* sulphide:quinone oxidoreductase, *Fcc* flavocytochrome *c* sulphide dehydrogenase, *SOR* sulphur oxygenase-reductase, *TetH* tetrathionate hydrolase, *TQO* thiosulphate–quinone oxidoreductase; *SulP* sulphate transporter permease, *QH₂* quinol pool



oxidoreductase was identified in the *A. hospitalis* genome. Thiosulphate enters the putative thiosulphate/tetrathionate cycle and is finally oxidised to sulphate. The enzymes involved in this cycle were all annotated: thiosulphate:quinone oxidoreductase (Ahos0112-0113 and Ahos0238-0239) and tetrathionate hydrolase (Ahos1670). H₂S is either oxidised by the sulphide:quinone oxidoreductase (Ahos1014) in the cytoplasm with quinone-cytochrome as electron acceptor or it reacts with tetrathionate spontaneously under the high temperature growth conditions. Finally, sulphate generated from sulphur oxidation is effluxed from the cell by a putative sulphate transport permease (Ahos1256). Electrons generated from sulphur oxidation enter the electron transport chain via quinone. Terminal quinol oxidase receives electrons from quinone and transfers them to O₂ coupled with ATP generation. Some electrons may be transmitted to the NADH complex to produce NADH for use in other pathways.

Transporters and proteolytic enzymes

Twenty-eight gene products were predicted to be involved in the transport of amino acids, oligopeptide/dipeptides and ammonium. Of these, 19 are implicated in amino acid transport, including 5 amino acid transporters (Ahos0100/0163/0197/0986/1721), three amino acid permeases (Ahos0328/0439/1725) and 11 amino acid permease-like proteins (Ahos0272/0276/0958/1040/1086/1868/1891/1907/1953/2065/2251) of unknown specificity for amino acid uptake. Genes encoding an ammonium transporter (Ahos1467) and two oligopeptide/dipeptide ABC transporter gene clusters (Ahos0337-0342 and Ahos0170-0175) are present. In addition, 21 genes were predicted to encode proteolytic enzymes, including 20 peptidases. Of these, four are endopeptidases (Ahos0428/0516/0695-6/0800), three are aminopeptidases (Ahos0013/0588/1941), two are pepsins (Ahos1929/2087) and one is a carboxypeptidase (Ahos0991). Five of the proteolytic enzymes are predicted to be membrane-bound and are designated secretory proteins. These results suggest that *A. hospitalis*, like *Acidianus brierleyi* (Seeger et al. 1986), *Acidianus tengchongensis* (He and Li 2004) and *Acidianus manzaensis* (Yoshida et al. 2006), can grow on organic compounds, such as yeast extract, peptone, tryptone and casamino acids.

Toxin–antitoxin systems

VapBC complexes constitute the main family of antitoxin–toxins that are encoded by members of the Sulfolobales (Pandey and Gerdes 2005; Guo et al. 2011), and they occur mainly in variable genomic regions where they may undergo loss or gain events (Guo et al. 2011). The *A. hospitalis* genome carries 26 *vapBC* gene pairs that are

concentrated in the genomic regions 350–410 and 1,374–1,912 kb with a single *vapC*-like gene lying in an operon (Fig. 1). The VapB antitoxins, in contrast to VapC toxins, could be classified into three families of transcriptional regulators, AbrB, CcdA/CopG and DUF217 (Fig. 4a), whilst no subclassification was observed for the VapC proteins (Fig. 4b). Tree building based on the sequence alignments demonstrated that the sequences of these antitoxins and toxins are highly diverse, with sequence identities between them rarely exceeding 30%, as indicated by all the proteins exhibiting long branches (Fig. 4). This result contrasted with the finding that VapBC complexes with closely similar sequences are commonly found when comparing different genomes of the Sulfolobales. For example, 11 of the 26 VapBC protein pairs have closely similar homologs encoded in at least 7 of the 13 available *Sulfolobus* genomes (Fig. 4b). This indicates that there is likely to be a selection against the uptake of closely similar *vapBC* gene pairs in a given genome, despite the abundance of such gene pairs in the environment.

The *A. hospitalis* genome also encodes six copies of RelE-related toxin proteins, in common with other *Sulfolobus* genomes (Pandey and Gerdes 2005, unpublished results). At least three of the *relE* genes occur in integrated regions carrying degenerated conjugative plasmids, and they show sequence similarity to proteins encoded in *Sulfolobus* conjugative plasmids pKEF9 (ORF69b), pING1 (ORF98) and pL085 (gene no. 3195) (Greve et al. 2004; Stedman et al. 2000; Reno et al. 2009). However, none of the putative toxin genes are linked physically to antitoxin *relB* genes and their function remains unknown.

Diverse CRISPR-based immune systems

The CRISPR-based immune systems of *A. hospitalis* can be classified into two main types based on analyses of their Cas1 protein, leader and repeat sequences (Shah et al. 2009; Lillestøl et al. 2009). In total, there are six CRISPR loci, carrying 129 spacer-repeat units none of which are identical (Fig. 5). The first three loci in the genome (Ahos-53, -13 and -9a) are physically linked by cassettes of *cmr* and *cas* family genes, each of which contains a *vapBC* antitoxin–toxin gene pair, and they constitute a family II CRISPR/Cas system (Fig. 5a). The last two CRISPR loci (Ahos-9b and 5) are coupled into a typical family I paired CRISPR/Cas module (Fig. 5b) and there is a *vapBC* gene pair immediately upstream. Preceding the latter CRISPR/Cas module, there is a single unclassified locus (Ahos-40) that lacks both *cas* genes and a leader region (Fig. 5c) (Shah and Garrett 2011).

We analysed the degree to which CRISPR spacers exhibited sequence matches to the many diverse genetic elements available from *Acidianus* and *Sulfolobus* species

using an earlier approach examining nucleotide and translated sequences of the spacers (Shah et al. 2009; Lillestøl et al. 2009). Relatively few significant sequence matches were found and most of these were to conjugative plasmids, with a few matches to members of five different viral families (Fig. 5).

Discussion

At about 2.1 Mbp, the genome of *A. hospitalis* is much smaller than other sequenced genomes of members of the Sulfolobales. Although this partly reflects the presence of low levels of transposable elements and few genes deriving from integrated elements, it also results from a lower diversity of metabolic and transporter genes (Guo et al. 2011). The Z curve analysis suggests that the chromosome carries three replication origins as for *Sulfolobus* species (Fig. 1), although in contrast to the sequenced strains of *S. solfataricus* and *S. islandicus*, the *whiP/cdt1* and *cdc6-2* genes are widely separated.

Although no systematic analysis has been performed experimentally on the metabolic capacity of *A. hospitalis*, genome analyses revealed that *A. hospitalis* possesses the capacity to assimilate a broad range of organic compounds, including different amino acids and proteolytic products, which is similar to some other *Acidianus* and *Sulfolobus* species (Seegerer et al. 1986; Grogan 1989; He et al. 2004; Yoshida et al. 2006; Plumb et al. 2007). The analyses also support that *A. hospitalis* can assimilate various carbohydrates, similarly to several *Sulfolobus* species (Grogan 1989) but in contrast to some *Acidianus* species (Yoshida et al. 2006; Plumb et al. 2007).

A. hospitalis, like other *Acidianus* and *Sulfolobus* species, obtains energy for growth mainly via oxidation of reduced inorganic sulphuric components (RISCs), and the enzymes involved were predicted from the genome analyses (Fig. 2). A sulphur oxygenase-reductase was identified showing amino acid sequence similarity to other *Acidianus* and *Sulfolobus* SORs of 67–99%, and we inferred that it is important for elemental sulphur oxidation and reduction, as occurs in both *Acidianus* and *Sulfolobus* species (Kletzin 1989, 1992; Sun et al. 2003; Chen et al. 2005a). One product of sulphur oxygenase-reductase catalysis is sulphite. Owing to the apparent lack of the four enzymes, sulphite-acceptor oxidoreductase, adenosine phosphosulphate reductase, sulphate adenylyl transferase and adenylylsulphate phosphate adenylyltransferase, *A. hospitalis* must have adopted a strategy for sulphite oxidation that differs from the currently known pathway (Kletzin 2007). Here, we propose that sulphite is channelled to thiosulphate in *A. hospitalis* via a spontaneous reaction with elemental sulphur, but this remains to be

tested experimentally. Some *Acidianus* species, such as *A. manzaensis* (Yoshida et al. 2006) and *A. sulfidivorans* (Plumb et al. 2007) grow chemolithoautotrophically with oxidation of molecular hydrogen, but this cannot occur in *A. hospitalis* because it apparently lacks an encoded hydrogen dehydrogenase.

Transposable elements include a few IS200/607 elements and several orphan *orfB* elements which all belong to the IS200/605/607 family. They lack inverted terminal repeats and are mobilised by “cut-and-paste” mechanisms (Filée et al. 2007; Ton-Hoang et al. 2010). No representatives of other transposable element families were found, common to other *Sulfolobus* genomes, which carry inverted terminal repeats and are mobilised by “copy-and-paste” mechanisms (Blount and Grogan 2005; Redder and Garrett 2006). It remains uncertain whether the OrfB protein is responsible for transposition of the *orfB* elements or whether they are mobilised *in trans* by the TnpA transposase encoded by the IS200/607 elements (Filée et al. 2007; Guo et al. 2011). The IS200/607 and *orfB* elements have been detected in *Sulfolobus* conjugative plasmids and *orfB* elements also occur in a few viruses of the Sulfolobales including four copies in the *Acidianus* two-tailed bicaudavirus ATV (She et al. 1998; Greve et al. 2004; Prangishvili et al. 2006). Thus, they are likely to be transmitted intercellularly, and enter chromosomes, via such genetic elements.

MITEs are common in *Sulfolobus* species and have been predicted to be mobilised by transposases encoded in different IS element families (Redder et al. 2001). The novel MITE-like elements in the *A. hospitalis* genome (Fig. 3) may derive from *orfB* elements and be mobilised by a similar mechanism but at present we can provide no evidence for their mobility. In this respect, they may be similar to other *Sulfolobus* MITEs which show a low level of transpositional activity (Redder and Garrett 2006). This is consistent with the hypothesis that MITEs drive the evolutionary diversification of their mobilising transposases to the point that they are no longer recognised which leads to their immobilisation and subsequent degeneration (Feschotte and Pritham 2007).

All of the integrated elements, except one, could be identified as originating from fuselloviruses or a pDL10-like member of the pRN family of cryptic plasmids (Kletzin et al. 1999), and the conjugative plasmid pAH1 was already shown to reversibly integrate at a tRNA^{Arg} [TCG] gene (Basta et al. 2009). None of these events occurred within any of the 15 tRNA genes carrying introns and this observation is consistent with the hypothesis that archaeal introns protect tRNA genes against integration events (Guo et al. 2011).

VapBC constitutes the predominant antitoxin–toxin family found amongst the Sulfolobales and the *A. hospitalis* genome carries 26 *vapBC* gene pairs, more than occur




Fig. 3 Alignment of 10 MITE-like repeat elements present in the genome of *A. hospitalis*. The shaded area denotes to a small open reading frame corresponding to the downstream part of the OrfB found within transposable *orfB* elements




Fig. 4 VapBC trees. Phylogenetic trees for **a** VapB antitoxins and **b** VapC toxins. They demonstrate that VapBs, despite their high sequence diversity, can be classified into three main families AbrB, CcdA/CopG and DUF217, whereas the VapCs are highly diverse in their sequences but cannot be classified into major subgroups. The Aho gene numbers are given for each protein. Moreover, the class of the VapB corresponding to each VapC is given in **b**. The degree of conservation of the VapC proteins in the available 13 *Sulfolobus*

genomes is indicated in **b** where 0 indicates that it is absent from all the genomes whilst 13 indicates that it is present in all. The antitoxin corresponding to VapC-0183 is not annotated in the genome because it lacks a start codon but it is included in the figure. The VapC-like protein (Aho0712) is part of the operon with a translation-related protein and lacks a VapB. The Aho1664/1663 pair are variant ORFs where both VapB and VapC are longer than usual and the VapB does not cluster with the families in **a**

in more rapidly growing *Sulfolobus* species (Pandey and Gerdes 2005; Guo et al. 2011). Moreover, the groups of VapB and VapC proteins are highly diverse in sequence (Fig. 4). Antitoxin–toxins were originally shown to enhance plasmid maintenance as a consequence of the growth of plasmid-free cells being preferentially inhibited by free toxins which are inherently more stable than the antitoxins (Gerdes 2000). By analogy with this mechanism,

it was proposed that chromosomally encoded toxins may facilitate maintenance of local DNA regions where *vapBC* gene pairs are located that might otherwise be prone to loss (Magnuson 2007; Van Melder 2010). This hypothesis is consistent with the observation that most of the *A. hospitalis* *vapBC* gene pairs lie within the two variable genomic regions where DNA regions are exchanged (Fig. 1). Moreover, it receives strong support from both the high




Fig. 5 Schematic representations of the CRISPR loci of *A. hospitalis*. **a** Family II CRISPR module carrying three CRISPR loci and Cmr and Cas family gene cassettes which are both interrupted by, or bordered by, four *vapBC* gene pairs (orange). **b** Paired family I CRISPR/Cas system flanked by one *vapBC* gene pair, and **c**. an unclassified CRISPR locus lacking a leader region and adjacent *cas* genes. *csm1* is a homolog of *cmr2*, *csm2* is a homolog of *cmr5* and *csm3* is a

homolog of *cmr4*. The light blue genes each carry two short RAMP motifs. **a–c** Structures of the individual CRISPR loci are shown together with the leader region (L) where each triangle represents a spacer-repeat unit. Significant spacer matches to sequenced viruses and plasmids are colour coded: red rudivirus, orange lipothrixvirus, yellow fusellovirus, green bicaudivirus, turquoise turreted icosahedral virus, blue conjugative plasmid and violet cryptic plasmid

diversity, and the uniqueness of all the VapC proteins encoded within the *A. hospitalis* chromosome (Fig. 4b), because any similar VapBC complexes would compensate for the loss of one another, thereby undermining any DNA maintenance activity.

In slowly growing organisms, from nutrient poor environments, multiple toxins are also assumed to be involved in stress response and/or quality control (Gerdes 2000; Pandey and Gerdes 2005). Involvement in stress response entails that the more stable toxins inhibit growth and allow the host to lie in a dormant state during the period of environmental stress (Gerdes 2000). However, there may also be a negative effect on host growth due to the continuous presence of low levels of free toxin (Wilbur et al. 2005). Thus, the presence of many *vapBC* gene pairs in *A. hospitalis* could reflect a compromise between the ability to survive different environmental stresses and maintaining an adequate growth rate under normal conditions. This would be also consistent with the presence of three families of VapB proteins and high sequence diversity of the VapC proteins, since functionally overlapping systems would be redundant for stress responses and they would confer an unnecessary burden on host growth. The proposed dual roles of maintenance of local chromosomal DNA regions and providing resistance to stress and are not mutually exclusive.

Although the mechanism of action of VapC toxins remains unknown (Arcus et al. 2011), in *A. hospitalis*, a single *vapC*-like gene (Ahos0712) is directly coupled to genes encoding proteins involved in transcription and initiator tRNA binding to the ribosome, and this gene cassette

is highly conserved in gene content and sequence in other *Sulfolobus* genomes (Guo et al. 2011). This suggests that this VapC protein, at least, may also regulate or inhibit translational initiation by binding at the ribosomal A-site, as demonstrated recently for a RelE type toxin (Neubauer et al. 2009). A similar inactivation mechanism would be plausible for the VapC toxins, if one assumes that expression of the individual VapBC complexes is stimulated by either the requirement to maintain different local regions of chromosomal DNA or different environmental stresses.

Despite the complexity of the CRISPR-based immune systems present in the genome, they appear to be, at best, only partially functional. Thus, the family II CRISPR/Cas system is coupled with an archaeal family D Cmr module in *A. hospitalis*, but is apparently defective, retaining only its putative RNA, but not DNA, targeting function. The system lacks the group 2 *cas* genes (*cas3*, *cas5*, *csa2*, *csa5*, *csaX*) which encode proteins implicated in targeting and inactivating foreign DNA elements (Fig. 5). However, the *cas* group 1 genes (*cas1*, *cas2*, *cas4*, *csa1*), putatively involved in integrating new spacers from invading DNA elements are present, and the Cmr module implicated in RNA targeting are also present (Garrett et al. 2011; Shah et al. 2011). The family I system exhibits small CRISPR loci, with intact leader regions and group 2 *cas* genes. However, the *cas2* gene in the group 1 *cas* gene cassette is truncated, having incurred a point mutation which produces a premature stop codon. Thus, this system has apparently lost the ability to integrate new spacers. This suggests that neither CRISPR-based system is fully functional, despite

their apparent complexity. The presence of five *vapBC* gene pairs located either within the *cmr* and *cas* gene cassettes of the family II CRISPR/Cas module, or immediately upstream from the modules of both families, may reflect that they help to maintain these gene cassettes on the chromosome (see above).

Although a range of genetic systems have been developed for *Sulfolobus* species, at present no genetic systems are available for the *Acidianus* genus and *A. hospitalis* provides a promising candidate for such studies. It has a minimal size and the relative stability of its chromosome suggests that it is likely to generate stable deletion mutants. This, combined with its ability to host different plasmids and viruses provides a promising starting point for developing a genetic system.

Acknowledgments We thank Mery Pina and Tamara Basta for help with the DNA preparation. The work was supported by the National Nature Science Foundation of China (30621005) and the Ministry of Science and Technology (2010CB630903), and by the Danish Natural Science Research Council (Grant no. 272-08-0391) and Danish National Research Foundation.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Arcus VL, McKenzie JL, Robson J, Cook GM (2011) The PIN-domain ribonucleases and the prokaryotic VapBC toxin–antitoxin array. *Prot Engin Design Select* 24:33–40
- Basta T, Smyth J, Forterre P, Prangishvili D, Peng X (2009) Novel archaeal plasmid pAH1 and its interactions with the lipothrix-virus AFV1. *Mol Microbiol* 71:23–34
- Bettstetter M, Peng X, Garrett RA, Prangishvili D (2003) AFV1, a novel virus infecting hyperthermophilic archaea of the genus *Acidianus*. *Virology* 315:68–79
- Blount ZD, Grogan DW (2005) New insertion sequences of *Sulfolobus*: functional properties and implications for genome evolution in hyperthermophilic archaea. *Mol Microbiol* 55:312–325
- Chen Z-W, Jiang C-Y, She Q, Liu S-J, Zhou P-J (2005a) Key role of cysteine residues in catalysis and subcellular localization of sulfur oxygenase reductase of *Acidianus tengchongensis*. *Appl Environ Microbiol* 71:621–628
- Chen L, Brügger K, Skovgaard M, Redder P, She Q, Torarinsson E, Greve B, Awayez M, Zibat A, Klenk HP, Garrett RA (2005b) The genome of *Sulfolobus acidocaldarius*, a model organism of the Crenarchaeota. *J Bacteriol* 187:4992–4999
- Cobucci-Ponzano B, Guzzini L, Benelli D, Londei P, Perrodou E, Lecompte O, Tran D, Sun J, Wei J, Mathur EJ, Rossi M, Moracci M (2010) Functional characterisation and high-throughput proteomic analysis of interrupted genes in the archaeon *Sulfolobus solfataricus*. *J Proteome Res* 9:2496–2507
- Feschotte C, Pritham EJ (2007) DNA transposons and the evolution of eukaryotic genomes. *Annu Rev Genet* 41:331–368
- Filée J, Siguier P, Chandler M (2007) Insertion sequence diversity in archaea. *Microbiol Mol Biol Revs* 71:121–157
- Garrett RA, Shah SA, Vestergaard G, Deng L, Gudbergdottir S, Kenchappa CS, Erdmann S, She Q (2011) CRISPR-based immune systems of the Sulfolobales—complexity and diversity. *Biochem Soc Trans* 39:51–57
- Gerdes K (2000) Toxin-antitoxin modules may regulate dynthsis of macromolecules during nutritional stress. *J Bacteriol* 182:561–572
- Goulet A, Blangy S, Redder P, Prangishvili D, Felisberto-Rodrigues C, Forterre P, Campanacci V, Cambillau C (2009) *Acidianus* filamentous virus 1 coat proteins display a helical fold spanning the filamentous archaeal viruses lineage. *Proc Natl Acad Sci USA* 106:21155–21160
- Greve B, Jensen S, Brügger K, Zillig W, Garrett RA (2004) Genomic comparison of archaeal conjugative plasmids from *Sulfolobus*. *Archaea* 1:231–239
- Grogan DW (1989) Phenotypic characterization of the archaeobacterial genus *Sulfolobus*: comparison of five wild-type strains. *J Bacteriol* 171:6710–6719
- Guo L, Brügger K, Liu C, Shah SA, Zheng H, Zhu Y, Wang S, Lillestøl RK, Chen L, Frank J, Prangishvili D, Paulin L, She Q, Huang L, Garrett RA (2011) Genome analyses of Icelandic strains of *Sulfolobus islandicus*: model organisms for genetic and virus-host interaction studies. *J Bacteriol* 193:1672–1680
- He Z-G, Zhong H, Li Y (2004) *Acidianus tengchongensis* sp. nov., a new species of acidothermophilic archaeon isolated from an acidothermal spring. *Curr Microbiol* 48:156–193
- Kletzin A (1989) Coupled enzymatic production of sulfite, thiosulfate, and hydrogen sulfide from sulfur: purification and properties of a sulfur oxygenase reductase from the facultatively anaerobic archaeobacterium *Desulfurolobus ambivalens*. *J Bacteriol* 171:1638–1643
- Kletzin A (1992) Molecular characterization of the *sor* gene, which encodes the sulfur oxygenase/reductase of the thermoacidophilic Archaeum *Desulfurolobus ambivalens*. *J Bacteriol* 174:5854–5859
- Kletzin A (2007) Oxidation of sulfur and inorganic sulfur compounds in *Acidianus ambivalens*. In: Dahl C, Friedrich CG (eds) *Microbial sulfur metabolism*. Springer, Heidelberg, pp 184–199
- Kletzin A, Lieke A, Urich T, Charlebois RL, Sensen CW (1999) Molecular analysis of pDL10 from *Acidianus ambivalens* reveals a family of related plasmids from extremely thermophilic and acidophilic archaea. *Genetics* 152:1307–1314
- Lawrence CM, Menon S, Eilers BJ, Bothner B, Khayat R, Douglas T, Young MJ (2009) Structural and functional studies of archaeal viruses. *J Biol Chem* 284:12599–12603
- Lillestøl RK, Shah SA, Brügger K, Redder P, Phan H, Christiansen J, Garrett RA (2009) CRISPR families of the crenarchaeal genus *Sulfolobus*: bidirectional transcription and dynamic properties. *Mol Microbiol* 72:259–272
- Lowe TM, Eddy SR (1997) tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25:955–964
- Lundgren M, Andersson A, Chen L, Nilsson P, Bernander R (2004) Three replication origins in *Sulfolobus* species: synchronous initiation of chromosome replication and asynchronous termination. *Proc Natl Acad Sci USA* 101:7046–7051
- Magnuson RD (2007) Hypothetical functions of toxin–antitoxin systems. *J Bacteriol* 189:6089–6092
- Melderer LV (2010) Toxin-antitoxin systems: why so many, what for? *Curr Opin Microbiol* 13:781–785
- Muller S, Urban A, Hecker A, Leclerc A, Branlant C, Motorin Y (2009) Deficiency of the tRNA^{Tyr}:Ψ35-synthase aPus7 in archaea of the Sulfolobales order might be rescued by the

- H/ACA sRNA-guided machinery. *Nucleic Acids Res* 37:1308–1322
- Muskhlishvili G, Palm P, Zillig W (1993) SSV1-encoded site-specific recombination system in *Sulfolobus shibatae*. *Mol Gen Genet* 273:334–342
- Neubauer C, Gao YG, Andersen KR, Dunham CM, Kelley AC, Hentschel J, Gerdes K, Ramakrishnan V, Brodersen DE (2009) The structural basis for mRNA recognition and cleavage by the ribosome-dependent endonuclease RelE. *Cell* 139:1084–1095
- Omer AD, Zago M, Chang A, Dennis PP (2006) Probing the structure and function of an archaeal C/D-box methylation guide sRNA. *RNA* 12:1708–1720
- Pandey DP, Gerdes K (2005) Toxin-antitoxin loci are highly abundant in free-living but lost from host-associated prokaryotes. *Nucleic Acids Res* 33:966–976
- Plumb JJ, Haddad CM, Gibson JAE, Franzmann PD (2007) *Acidianus sulfidivorans* sp. nov., an extremely acidophilic, thermophilic archaeon isolated from a solfatara on Lihir Island, Papua New Guinea, and amendment of the genus description. *Int J Syst Evol Microbiol* 57:1418–1423
- Prangishvili D, Albers SV, Holz I, Arnold HP, Stedman K, Klein T, Singh H, Hiort J, Schweier A, Kristjansson JK, Zillig W (1998) Conjugation in archaea: frequent occurrence of conjugative plasmids in *Sulfolobus*. *Plasmid* 40:190–202
- Prangishvili D, Forterre P, Garrett RA (2006) Viruses of the Archaea: a unifying view. *Nat Rev Microbiol* 4:837–848
- Rachel R, Bettstetter M, Hedlund BP, Häring M, Kessler A, Stetter KO, Prangishvili D (2002) *Arch Virol* 147:2419–2429
- Redder P, Garrett RA (2006) Mutations and rearrangements in the genome of *Sulfolobus solfataricus* P2. *J Bacteriol* 188:4198–4206
- Redder P, She Q, Garrett RA (2001) Non-autonomous elements in the crenarchaeon *Sulfolobus solfataricus*. *J Mol Biol* 306:1–6
- Redder P, Peng X, Brügger K, Shah SA, Roesch F, Greve B, She Q, Schleper C, Forterre P, Garrett RA, Prangishvili D (2009) Four newly isolated fuselloviruses from extreme geothermal environments reveal unusual morphologies and a possible intervirial recombination mechanism. *Environ Microbiol* 11:2849–2862
- Reno ML, Held NL, Fields CJ, Burke PV, Whitaker RJ (2009) *Sulfolobus islandicus* pan-genome. *Proc Natl Acad Sci USA* 106:8605–8610
- Robinson NP, Bell SD (2007) Extrachromosomal element capture and the evolution of multiple replication origins in archaeal chromosomes. *Proc Natl Acad Sci USA* 104:5806–5811
- Robinson NP, Dionne I, Lundgren M, Marsh VL, Bernander R, Bell SD (2004) Identification of two origins of replication in the single chromosome of the archaeon *Sulfolobus solfataricus*. *Cell* 116:25–38
- Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B (2000) Artemis: sequence visualization and annotation. *Bioinformatics* 16:944–945
- Segerer A, Neuner A, Kristjansson JK, Stetter KO (1986) *Acidianus infernus* gen. nov., sp. nov., and *Acidianus brierleyi* comb. nov.: facultatively aerobic, extremely acidophilic thermophilic sulfur-metabolizing archaeobacteria. *Int J Syst Bacteriol* 36:559–564
- Shah SA, Garrett RA (2011) CRISPR/Cas and Cmr modules, mobility and evolution of adaptive immune systems. *Res Microbiol* 162:27–38
- Shah SA, Hansen NR, Garrett RA (2009) Distributions of CRISPR spacer matches in viruses and plasmids of crenarchaeal acidothermophiles and implications for their inhibitory mechanism. *Trans Biochem Soc* 37:23–28
- Shah SA, Vestergaard G, Garrett RA (2011) CRISPR/Cas and CRISPR/Cmr immune systems of archaea. In: Marchfelder A, Hess W (eds) *Regulatory RNAs in prokaryotes*. Springer, Berlin
- She Q, Phan H, Garrett RA, Albers S-V, Stedman KM, Zillig W (1998) Genetic profile of pNOB8 from *Sulfolobus*: the first conjugative plasmid from an archaeon. *Extremophiles* 2:417–425
- Stedman KM, She Q, Phan H, Holz I, Singh H, Prangishvili D, Garrett RA, Zillig W (2000) The pING family of conjugative plasmids from the extremely thermophilic archaeon *Sulfolobus islandicus*: insights into recombination and conjugation in Crenarchaeota. *J Bacteriol* 182:7014–7020
- Sun CW, Chen ZW, He ZG, Zhou PJ, Liu SJ (2003) Purification and properties of the sulphur oxygenase/reductase from the acidothermophilic archaeon, *Acidianus* strain S5. *Extremophiles* 7:131–134
- Tang TH, Polacek N, Zywicki M, Huber H, Brügger K, Garrett R, Bachelier JP, Hüttenhofer A (2005) Identification of novel non-coding RNAs as potential antisense regulators in the archaeon *Sulfolobus solfataricus*. *Mol Microbiol* 55:469–481
- Ton-Hoang B, Pasternak C, Siguier P, Guynet C, Hickman AB, Dyda F, Sommer S, Chandler M (2010) Single-stranded DNA transposition is coupled to host replication. *Cell* 142:398–408
- Torarinsson E, Klenk H-P, Garrett RA (2005) Divergent transcriptional and translational signals in Archaea. *Environ Microbiol* 7:47–54
- Wilbur JS, Chivers PT, Mattison K, Potter L, Brennan RG, So M (2005) *Neisseria gonorrhoeae* FitA interacts with FitB to bind DNA through its ribbon-helix-helix motif. *Biochemistry* 44:12515–12524
- Wurtzel O, Sapra R, Chen F, Zhu ZY, Simmons BA, Sorek R (2010) A single-base resolution map of an archaeal transcriptome. *Genome Res* 20:133–141
- Yokobori S, Itoh T, Yoshinari S, Nomura N, Sako Y, Yamagishi A, Oshima T, Kita K, Watanabe Y (2009) Gain and loss of an intron in a protein-coding gene in Archaea: the case of an archaeal RNA pseudouridine synthase gene. *BMC Evol Biol* 9:198
- Yoshida N, Nakasato M, Ohmura N, Ando A, Saolo J, Ishii M, Igarashi Y (2006) *Acidianus manzaensis* sp. nov., a novel thermoacidophilic Archaeon growing autotrophically by the oxidation of H₂ with the reduction of Fe³⁺. *Curr Microbiol* 53:406–411
- Zhang R, Zhang CT (2003) Multiple replication origins of the archaeon *Halobacterium* species NRC-1. *Biochem Biophys Res Comm* 302:728–734