Correction to: *Aspergillus loretoensis*, a single isolate from marine sediment of Loreto Bay, Baja California Sur, México resulting as a new obligate halophile species

Sophia González-Martínez¹ · Clara Galindo-Sánchez² · Edgar López-Landavery² · Carmen Paniagua-Chávez² · Amelia Portillo-López¹

Published online: 22 July 2019
© Springer Japan KK, part of Springer Nature 2019

Correction to: *Extremophiles*
<https://doi.org/10.1007/s00792-019-01107-6>

In the original publication the Figs. 3 and 4 are published inappropriately. The correct version of Figs. 3 and 4 is provided in this correction.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00792-019-01107-6>.

✉ Amelia Portillo-López
portillo@uabc.edu.mx

¹ Facultad de Ciencias, Universidad Autónoma de Baja California, Km 103, Carretera Tijuana-Ensenada, Ensenada 22860, Baja California, Mexico

² Centro de Investigación Científica y de Educación Superior de Ensenada, Ensenada 22860, Baja California, Mexico

Fig. 3 Phylogenetic tree based on the maximum likelihood method with bootstrap values expressed as a percentage of different sequences data. ITS1–2; Taxon names are preceded by the GBAN. *Aspergillus wentii* was used as an outgroup

Fig. 4 Phylogenetic tree based on the maximum likelihood method with bootstrap values expressed as a percentage of different sequences data. **a** RPB1; **b** TSR1; **c** RPB2; **d** Cct8; **e** BenA; **f** CaM. Taxon names are preceded by the GBAN. *Aspergillus wentii* was used as an outgroup