

Erratum to: ATP-association to intrabacterial nanotransportation system in *Vibrio cholerae*

Yuji Matsuzaki¹ · Hong Wu¹ · Takashi Nakano¹ · Takashi Nakahari^{2,3} · Kouichi Sano¹

Published online: 2 July 2015
© The Japanese Society for Clinical Molecular Morphology 2015

Erratum to: Med Mol Morphol
DOI 10.1007/s00795-015-0105-4

In the original publication of the article, the numbers on the Y-axis scale in Fig. 1b were incorrectly rounded to one decimal place. The correct Fig. 1 is given in the following page.

The online version of the original article can be found under doi:[10.1007/s00795-015-0105-4](https://doi.org/10.1007/s00795-015-0105-4).

✉ Kouichi Sano
ksano@art.osaka-med.ac.jp

¹ Department of Microbiology and Infection Control, Osaka Medical College, 2-7 Daigaku-machi, Takatsuki-shi, Osaka 569-8686, Japan

² Department of Physiology, Osaka Medical College, 2-7 Daigaku-machi, Takatsuki-shi, Osaka 569-8686, Japan

³ Present Address: Department of Molecular Cell Physiology, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, Kyoto 602-8566, Japan




Fig. 1 Reductive effect of DNP on intracellular ATP level. DNP decreased intracytoplasmic ATP level in concentration-dependent manner (a). The decrease was reversed by neutralization of DNP activity (b). The decrease of ATP level was maintained after a 15-min incubation at pHs 7 and 8 of extrabacterial environment (c). Values plotted are the mean \pm SD from triplicated experiments