



## Correction: Regaining enamel color quality using enamel matrix derivative

Hiroyuki Sugaya<sup>1</sup> · Yoshihito Kurashige<sup>1</sup> · Kai Suzuki<sup>1</sup> · Sayaka Sakakibara<sup>1</sup> · Yusuke Fujita<sup>1</sup> · Syed Taufiqul Islam<sup>1</sup> · Takashi Nezu<sup>2</sup> · Shuichi Ito<sup>3</sup> · Yoshihiro Abiko<sup>4</sup> · Masato Saitoh<sup>1</sup> 

Published online: 7 August 2023  
© The Author(s) 2023

**Correction: Medical Molecular Morphology (2023)  
56:116–127**  
<https://doi.org/10.1007/s00795-022-00346-5>

In the original publication of the article, the last sentence “All of the mineralization buffers used in the present study had a Ca/P molar ratio of 1.67; the degrees of solution saturations with respect to HAp were (A)  $2.58 \times 10^7$ , (B)  $3.85 \times 10^7$ , (C)  $5.50 \times 10^7$ , and (D)  $8.32 \times 10^7$  [25].” should be deleted in “Materials and methods”, under the subheading “Mineralization buffers” in the last paragraph in page 117.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The original article can be found online at <https://doi.org/10.1007/s00795-022-00346-5>.

✉ Masato Saitoh  
msaitoh@hoku-iryo-u.ac.jp

- <sup>1</sup> Division of Pediatric, Dentistry School of Dentistry, Health Sciences University of Hokkaido, 1757 Kanazawa, Tobetsu, Ishikari, Hokkaido 061-0293, Japan
- <sup>2</sup> Division of Biomaterials and Bioengineering, School of Dentistry, Health Sciences University of Hokkaido, Tobetsu, Japan
- <sup>3</sup> Division of Dental Education Development, School of Dentistry, Health Sciences University of Hokkaido, Tobetsu, Japan
- <sup>4</sup> Division of Oral Medicine and Pathology, Dentistry School of Dentistry, Health Sciences University of Hokkaido, Tobetsu, Japan