Correction to: Inguinal hernia repair in Spain. A population-based study of 263,283 patients: factors associated with the choice of laparoscopic approach

S. Guillaumes¹ · C. Hoyuela^{1,2} · N. J. Hidalgo¹ · M. Juvany¹ · I. Bachero¹ · J. Ardid¹ · A. Martrat¹ · M. Trias¹

Published online: 27 April 2021

© Springer-Verlag France SAS, part of Springer Nature 2021

Correction to: Hernia

<https://doi.org/10.1007/s10029-021-02402-y>

While typesetting the article the Table 1 rows are misplaced.

The correct Table 1 has been copied below:

The original article has been updated.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10029-021-02402-y>.

✉ S. Guillaumes
sguillaumes@gmail.com

¹ Department of General and Digestive Surgery, Hospital Plató, c/ Plató 21, 08006 Barcelona, Spain

² Universitat Internacional de Catalunya, Barcelona, Spain

Table 1 Patient characteristics by laparoscopic or open repair

	Overall	Open <i>n</i> (%)	LAP-IHR <i>n</i> (%)	OR	CI 95%		<i>p</i> *
Age < 65 years	152,274	141,864 (57.2)	10,410 (69.1)	1.68	1.62	1.74	<0.001
Age groups (years)							
15–44	42,715	39,382 (15.9)	3333 (22.1)	1.51	1.45	1.57	<0.001
45–64	109,559	102,482 (41.3)	7077 (47.0)	1.26	1.22	1.30	<0.001
65–74	62,590	59,455 (24.0)	3135 (20.8)	0.84	0.80	0.87	<0.001
> 74	48,419	46,905 (18.9)	1514 (10.1)	0.48	0.46	0.51	<0.001
Gender							
Male	233,810	220,204 (88.7)	13,606 (90.4)	1.19	1.13	1.26	<0.001
Female	29,426	27,978 (11.3)	1448 (9.6)	0.84	0.79	0.89	<0.001
Recurrence							
Primary hernia	245,446	231,809 (93.4)	13,637 (90.6)	0.68	0.64	0.72	<0.001
Recurrent hernia	17,837	16,415 (6.6)	1422 (9.4)	1.47	1.39	1.56	<0.001
Laterality							
Unilateral repair	212,216	203,063 (93.1)	9153 (75.2)	0.23	0.22	0.23	<0.001
Bilateral repair	20,866	17,132 (6.9)	3734 (24.8)	4.45	4.27	6.63	<0.001
Comorbidities							
Hypertension	54,473	52,197 (21.0)	2276 (15.1)	0.67	0.64	0.70	<0.001
Dyslipidemia	32,619	31,197 (12.6)	1422 (9.4)	0.73	0.69	0.77	<0.001
Diabetes	16,903	16,271 (6.6)	632 (4.2)	0.62	0.58	0.68	<0.001
COPD (c)	11,813	11,369 (4.6)	444 (2.9)	0.63	0.58	0.70	<0.001
Obesity	4989	4,807 (1.9)	182 (1.2)	0.62	0.53	0.72	<0.001
Tobacco use	22,979	21,740 (8.8)	1239 (8.2)	0.93	0.88	0.99	0.025
Case characteristics							
Not complicated hernia	253,715	239,116 (96.3)	14,599 (96.9)	1.21	1.10	1.33	<0.001
Complicated hernia	9568	9108 (3.7)	460 (3.1)	0.83	0.75	0.91	<0.001
Obstruction	8788	8374 (3.4)	414 (2.7)	0.81	0.73	0.90	<0.001
Gangrene	780	734 (0.3)	46 (0.3)	1.03	0.77	1.39	ns
Hospital ownership							
Public	195,988	18,7531 (75.5)	8457 (56.2)	0.41	0.40	0.43	<0.001
Private	67,295	60,693 (24.5)	6602 (43.8)	2.41	2.33	2.49	<0.001
Insurance type							
Public	207,942	200,531 (80.8)	7411 (49.2)	0.23	0.22	0.24	<0.001
No insurance/self-pay	3742	3,349 (1.3)	393 (2.6)	1.96	1.76	2.18	<0.001
Private insurance	32,366	27,139 (10.9)	5227 (34.7)	4.33	4.18	4.49	<0.001
Other funds	19,233	17,205 (6.9)	2028 (13.5)	2.09	1.99	2.20	<0.001
Hospital size (beds)							
200 beds or fewer	115,062	106,290 (42.8)	8772 (58.3)	1.86	1.80	1.92	<0.001
201–500 beds	76,972	73,756 (29.7)	3216 (21.4)	0.62	0.59	0.64	<0.001
501–1000 beds	45,953	44,260 (17.8)	1693 (11.2)	0.58	0.55	0.61	<0.001
More than 1000 beds	24,804	23,437 (9.4)	1367 (9.1)	0.96	0.90	1.01	ns
Admission							
Inpatient	142,153	131,620 (52.9)	10,533 (69.9)	2.07	2.00	2.15	<0.001
Outpatient	121,130	116,604 (47.0)	4526 (30.1)	0.49	0.47	0.50	<0.001
Planned admission	234,454	220,772 (88.9)	13,682 (90.9)	1.24	1.17	1.31	<0.001
Unplanned admission	18,727	18,018 (7.3)	709 (4.7)	0.63	0.59	0.68	<0.001
Unknown	10,102	9434 (3.8)	668 (4.6)	1.18	1.08	1.27	<0.001

OR odds ratio, CI confidence interval, COPD chronic obstructive pulmonary disease

*Chi-square test