



## Correction to: It is time for an organised, scientific approach to the application of patient-reported outcome measures in clinical studies and trials

Stephen P. McKenna<sup>1,2</sup> 

Published online: 26 June 2019

© International League of Associations for Rheumatology (ILAR) 2019

**Correction to: Clinical Rheumatology (2019) 38:611–615**  
<https://doi.org/10.1007/s10067-018-04403-2>

In the above article [1], the section labelled ‘Misuse of the ASQoL’ incorrectly lists Reference 19 [2] as having used an unauthorised language version of the Ankylosing Spondylitis Quality of Life Questionnaire (ASQoL).

The author wishes to correct the record and clarify that the study by Gupta, Parshad and colleagues did not use an unauthorised language version of the ASQoL. The study pertained to the comparison of two procedures in the management of patients with Achalasia Cardia and one of the scales used by the authors was Achalasia disease-specific health-related quality of life.

### References

1. McKenna S (2019) It is time for an organised, scientific approach to the application of patient-reported outcome measures in clinical studies and trials. *Clin Rheum* 38:611–615
2. Gupta P, Parshad R, Balakrishna P, Saraya A, Makharia GK, Sachdeva S, Sharma R (2018) Angle of his accentuation is a viable alternative to dor fundoplication as an adjunct to laparoscopic Heller cardiomyotomy: results of a randomized clinical study. *Dig Dis Sci* 63:2395–2404

**Publisher’s note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The online version of the original article can be found at <https://doi.org/10.1007/s10067-018-04403-2>

---

✉ Stephen P. McKenna  
smckenna@galen-research.com

<sup>1</sup> Galen Research, Manchester, UK

<sup>2</sup> School of Health Sciences, University of Manchester, Manchester, UK