Correction to: The ultrasonographic study of the nail reveals differences in patients affected by inflammatory and degenerative conditions

Luca Idolazzi¹ • Alen Zabotti² • Angelo Fassio¹ • Enzo Errichetti³ • Camilla Benini¹ • Elisabetta Vantaggiato¹ • Maurizio Rossini¹ • Salvatore De Vita² • Ombretta Viapiana¹

Published online: 14 January 2020

© International League of Associations for Rheumatology (ILAR) 2020

Correction to: Clinical Rheumatology (2019) 38:913–920
<https://doi.org/10.1007/s10067-019-04437-0>

The original published version of this article contained an incorrect Table 2.

The correct Table 2 shown as follows:

Table 2 alterations regarding qualitative and semi quantitative parameters of the nail, joint and enthesis complex

		PSO	PSA	RA	OA	HC
PDUS Nail Bed Case count	Absent	14	21	17	6	38 [^]
	Mild	9	16	8	12	9
	Moderate	1	9	7	9	2
	Severe	7	5	5	7	1
PDUS enthesis Case count	Absent	20	23	35 [^]	25	48 [^]
	Present	11	28*	2	9	2

PDUS: power Doppler ultrasonography signal

*residual value significant with a trend which differs from the standard behaviour towards a pathological feature

[^] residual value significant with a trend which differs from the standard behaviour towards a normal feature

The online version of the original article can be found at <https://doi.org/10.1007/s10067-019-04437-0>

✉ Luca Idolazzi
luca.idolazzi@univr.it

¹ Rheumatology Unit, Department of Medicine, University of Verona, Piazzale A. L. Scuro 1, 37138 Verona, Italy

² Department of Medical and Biological Sciences, Rheumatology Clinic, University of Udine, Udine, Italy

³ Department of Experimental and Clinical Medicine, Institute of Dermatology, University of Udine, Udine, Italy