

Role of spiramycin/cotrimoxazole association in the mother-to-child transmission of toxoplasmosis infection in pregnancy

P. Valentini · M. L. Annunziata · D. F. Angelone ·
L. Masini · M. De Santis · A. Testa · R. L. Grillo ·
D. Speziale · O. Ranno

Published online: 20 May 2009
© Springer-Verlag 2009

Erratum to: Eur J Clin Microbiol Infect Dis
DOI 10.1007/s10096-008-0612-5

In the published original version of this article, page 298,
23rd line, the data in brackets (3×10^6 International Unit =
3.480 g) is not given correct.

The line should read:

spiramycin, 3×10^6 IU (3×10^6 International Unit = 870 mg)

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10096-008-0612-5>

P. Valentini (✉) · M. L. Annunziata · D. F. Angelone · A. Testa ·
O. Ranno
Department of Pediatrics, Pediatric Infectious Diseases Unit,
Catholic University of the Sacred Heart,
L.go A. Gemelli, 8,
00168 Rome, Italy
e-mail: pvalentini@rm.unicatt.it

L. Masini · M. De Santis
Department of Obstetrics and Gynecology,
Catholic University of the Sacred Heart,
Rome, Italy

R. L. Grillo · D. Speziale
Institute of Microbiology, Catholic University of the Sacred Heart,
Rome, Italy