



Correction to: Effectiveness of second-look endoscopy after gastric endoscopic submucosal dissection in patients taking antithrombotic agents: a multicenter propensity score matching analysis

Taro Iwatsubo¹ · Toshihisa Takeuchi² · Akitoshi Hakoda¹ · Yasuhiro Fujiwara³ · Yasuaki Nagami³ · Yuji Naito⁴ · Osamu Dohi⁴ · Tetsuya Tatsuta⁵ · Manabu Sawaya⁶ · Xiaoyi Jin⁷ · Tomoyuki Koike⁷ · Mitsushige Sugimoto⁸ · Masaki Murata⁹ · Kenta Hamada¹⁰ · Hiroyuki Okada¹⁰ · Hideki Kobara¹¹ · Taiga Chiyo¹¹ · Norimasa Yoshida¹² · Naoya Tomatsuri¹³ · Tomoki Inaba¹⁴ · Shigenao Ishikawa¹⁴ · Akihito Nagahara¹⁵ · Hiroya Ueyama¹⁵ · Eriko Koizumi¹⁶ · Katsuhiko Iwakiri¹⁶ · Kazuhiro Mizukami¹⁷ · Kazunari Murakami¹⁷ · Takahisa Furuta¹⁸ · Takahiro Suzuki¹⁸ · Naotaka Ogasawara¹⁹ · Kunio Kasugai¹⁹ · Hajime Isomoto²⁰ · Koichiro Kawaguchi²⁰ · Kotaro Shibagaki²¹ · Hiromi Kataoka²² · Takaya Shimura²² · Hidekazu Suzuki²³ · Toshihiro Nishizawa²⁴ · Kazuhide Higuchi¹

Published online: 30 July 2022

© The Author(s) under exclusive licence to The International Gastric Cancer Association and The Japanese Gastric Cancer Association 2022

Correction to: Gastric Cancer

<https://doi.org/10.1007/s10120-022-01303-y>

In the original publication of the article, the Table 4 has been set incorrectly. The correct Table 4 is given in this

The original article can be found online at <https://doi.org/10.1007/s10120-022-01303-y>.

✉ Toshihisa Takeuchi
toshihisa.takeuchi@ompu.ac.jp

¹ Second Department of Internal Medicine, Osaka Medical and Pharmaceutical University Hospital, Osaka, Japan

² Endoscopic Center, Osaka Medical and Pharmaceutical University Hospital, 2-7, Daigakumachi Takatsuki, Osaka 569-8686, Japan

³ Department of Gastroenterology, Osaka City University Graduate School of Medicine, Osaka, Japan

⁴ Molecular Gastroenterology and Hepatology, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, Kyoto, Japan

⁵ Departments of Gastroenterology and Hematology, Hirosaki University Graduate School of Medicine, Aomori, Japan

⁶ Departments of Gastroenterology and Hematology, Hirosaki National Hospital, Aomori, Japan

⁷ Division of Gastroenterology, Tohoku University Graduate School of Medicine, Miyagi, Japan

⁸ Department of Gastroenterological Endoscopy, Tokyo Medical University Hospital, Tokyo, Japan

⁹ Department of Gastroenterology, National Hospital Organization Kyoto Medical Center, Kyoto, Japan

¹⁰ Departments of Gastroenterology and Hepatology, Okayama University Graduate School of Medicine, Okayama, Japan

¹¹ Department of Gastroenterology and Neurology, Faculty of Medicine, Kagawa University, Kagawa, Japan

¹² Department of General Internal Medicine and Gastroenterology, Saiseikai Kyoto Hospital, Kyoto, Japan

¹³ Department of Gastroenterology, Japanese Red Cross Kyoto Daiichi Hospital, Kyoto, Japan

¹⁴ Department of Gastroenterology, Kagawa Prefectural Central Hospital, Kagawa, Japan

¹⁵ Department of Gastroenterology, Juntendo University School of Medicine, Tokyo, Japan

¹⁶ Department of Gastroenterology, Graduate School of Medicine, Nippon Medical School, Tokyo, Japan

¹⁷ Department of Gastroenterology, Faculty of Medicine, Oita University, Oita, Japan

¹⁸ Center for Clinical Research, Hamamatsu University School of Medicine, Shizuoka, Japan

¹⁹ Department of Gastroenterology, Aichi Medical University Hospital, Aichi, Japan

²⁰ Division of Gastroenterology and Nephrology, Department of Multidisciplinary Internal Medicine, Faculty of Medicine, Tottori University, Tottori, Japan

²¹ Department of Gastroenterology, Faculty of Medicine, Shimane University, Shimane, Japan

²² Department of Gastroenterology and Metabolism, Nagoya City University Graduate School of Medical Sciences, Aichi, Japan

²³ Department of Gastroenterology and Hepatology, Tokai University School of Medicine, Kanagawa, Japan

²⁴ Department of Gastroenterology and Hepatology, International University of Health and Welfare, Narita Hospital, Tokyo, Japan

Table 4 Incidence of delayed bleeding in the matched cohort between the SLE and non-SLE groups, according to risk category

BEST-J score									
Predicted bleeding rate	> Intermediate (6.1%)			> High (11.4%)			> Very high (29.7%)		
	Delayed bleeding, <i>n</i> (%)	OR (95% CI)	<i>p</i> value	Delayed bleeding, <i>n</i> (%)	OR (95% CI)	<i>p</i> value	Delayed bleeding, <i>n</i> (%)	OR (95% CI)	<i>p</i> value
Non-SLE group	16/158 (10.1)	Reference		13/107 (12.1)	Reference		6/31 (19.3)	Reference	
SLE group	10/175 (5.7)	0.538 (0.211–1.309)	0.154	7/126 (5.6)	0.426 (0.138–1.205)	0.100	4/30 (13.3)	0.645 (0.119–3.103)	0.731
SAMURAI model									
Predicted bleeding rate	> 5%			> 10%			> 25%		
	Delayed bleeding, <i>n</i> (%)	OR (95% CI)	<i>p</i> value	Delayed bleeding, <i>n</i> (%)	OR (95% CI)	<i>p</i> value	Delayed bleeding, <i>n</i> (%)	OR (95% CI)	<i>p</i> value
Non-SLE group	15/117 (12.8)	Reference		12/76 (15.8)	Reference		4/21 (19.0)	Reference	
SLE group	7/122 (5.7)	0.415 (0.137–1.134)	0.073	6/76 (7.9)	0.459 (0.133–1.416)	0.209	5/20 (25.0)	1.404 (0.249–8.494)	0.719

CI confidence interval, OR odds ratio, SLE second-look endoscopy.

correction. In addition, the ORCID iD of the first author, Taro Iwatsubo (0000-0001-7721-8292) is given in this correction.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.